

## On the Sets of Strongly $f$ -Lacunary Summable Sequences

Ibrahim S. Ibrahim<sup>a,b\*</sup>, Rifat Çolak<sup>a</sup>

<sup>a</sup>Department of Mathematics, Faculty of Science, Firat University, Elazig,  
Türkiye

<sup>b</sup>Department of Mathematics, Faculty of Science, University of Zakho, Zakho,  
Iraq

E-mail: [ibrahimmath95@gmail.com](mailto:ibrahimmath95@gmail.com)

E-mail: [rftcolak@gmail.com](mailto:rftcolak@gmail.com)

**ABSTRACT.** The statistical convergence with respect to a modulus function has various applications in both mathematics and statistics. The main purpose of this research paper is to establish the relations between the sets of strongly  $f$ -lacunary summable and strongly  $g$ -lacunary summable sequences, strongly  $f$ -lacunary summable and  $g$ -lacunary statistically convergent sequences, where  $f$  and  $g$  are different modulus functions under certain conditions. Furthermore, for some special modulus functions, we establish the relations between the sets of strongly  $f$ -lacunary summable and strongly lacunary summable sequences.

**Keywords:** Modulus function, Statistical convergence,  $f$ -lacunary statistical convergence, Strong  $f$ -lacunary summability.

**2010 Mathematics subject classification:** 40A05, 40C05, 46A45.

### 1. INTRODUCTION

The notion of statistical convergence was first based on the first edition of the monograph of Zygmund [22], and its definition was introduced by Fast [10] in a short note and later reintroduced by Schoenberg [21] independently

---

\*Corresponding Author

Received 09 August 2020; Accepted 31 July 2022

©2025 Academic Center for Education, Culture and Research TMU

with some of the basic properties of statistical convergence. In recent decades, statistical convergence has been confirmed in many several fields and under various names, such as measure theory, approximation theory, Banach spaces, hopfield neural network, locally convex spaces, trigonometric series, number theory, summability theory, ergodic theory, turnpike theory, Fourier analysis, and optimization. Subsequently, Šalát [20], Fridy [12], Connor [8], Rath and Tripathy [18], Et [9], and many others were further explored from the point of perspective of sequence spaces and related to summability theory.

The idea of a modulus function was structured by Nakano [17]. Some authors such as Ruckle [19] and Maddox [15] have introduced some sequence spaces by using a modulus function. Other than them, to establish a number of sequence spaces, Gosh and Srivastava [13], Bhardwaj and Singh [3], and some others have used a modulus function. More information on this principle, as well as its applications, can be found in [2, 5, 6, 7, 14].

We denote by  $\mathbb{N}$  the set of natural numbers. The number  $\delta(A)$  of a set  $A \subset \mathbb{N}$  is called the natural density of  $A$  and is defined by

$$\delta(A) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : k \in A\}|$$

where  $|\{k \leq n : k \in A\}|$  denotes the number of elements of  $A$  which are less than or equal to  $n$ . One easily may see that  $\delta(A) = 0$  if  $A \subset \mathbb{N}$  is a finite set and  $\delta(\mathbb{N}) = 1$  and also  $\delta(\mathbb{N} - A) = \delta(\mathbb{N}) - \delta(A) = 1 - \delta(A)$ .

In this paper, the spaces of bounded and convergent sequences are symbolized by  $\ell_\infty$  and  $c$ , respectively, as well as the set of all complex numbers is symbolized by  $\mathbb{C}$ .

A sequence  $(x_k)$  in  $\mathbb{C}$  is named statistically convergent to the number  $l \in \mathbb{C}$  if  $\delta(\{k \in \mathbb{N} : |x_k - l| \geq \varepsilon\}) = 0$  for each  $\varepsilon > 0$ .

We imply an increasing sequence  $\theta = (k_r)$  of non-negative integer numbers with  $k_0 = 0$  by a lacunary sequence such that  $h_r = k_r - k_{r-1} \rightarrow \infty$  as  $r \rightarrow \infty$ . The intervals put by  $\theta$  shall be represented by  $I_r = (k_{r-1}, k_r]$  and the ratio  $\frac{k_r}{k_{r-1}}$  can be shortened by  $q_r$ .

Fridy and Orhan [11] defined lacunary statistical convergence as the following expression.

Let  $\theta = (k_r)$  be a lacunary sequence. A sequence  $(x_k)$  of numbers is named lacunary statistically convergent (or  $S_\theta$ -convergent) to the number  $l$ , if

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} |\{k \in I_r : |x_k - l| \geq \varepsilon\}| = 0$$

for each  $\varepsilon > 0$ . In this case, we write  $S_\theta - \lim x_k = l$ . Throughout the paper, the class of  $S_\theta$ -convergent sequences will be symbolized by  $S_\theta$ .

A function  $f : [0, \infty) \rightarrow [0, \infty)$  is called a modulus function (or a modulus) if

$$(1) \quad f(t) = 0 \text{ if and only if } t = 0,$$

- (2)  $f(t_1 + t_2) \leq f(t_1) + f(t_2)$  for every  $t_1, t_2 \in [0, \infty)$ ,
- (3)  $f$  is increasing,
- (4)  $f$  is continuous from the right at 0.

From these properties, it is clear that a modulus function must be continuous everywhere on  $[0, \infty)$ . A modulus function may be either unbounded or bounded. For instance,  $f(t) = \log(t + 1)$  is unbounded, but  $f(t) = \frac{t}{t+1}$  is bounded.

**Lemma 1.1.** [16] For any modulus  $f$ ,  $\lim_{t \rightarrow \infty} \frac{f(t)}{t}$  exists and  $\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \inf_{t \in (0, \infty)} \frac{f(t)}{t}$ .

The following definition was presented by Aizpuru et al. [1].

The number  $\delta_f(A)$  of a set  $A \subset \mathbb{N}$  is defined by

$$\delta_f(A) = \lim_{n \rightarrow \infty} \frac{f(|\{k \leq n : k \in A\}|)}{f(n)}$$

and is called the  $f$ -density of  $A$ , where  $f$  is an unbounded modulus function.

A sequence  $(x_k)$  in  $\mathbb{C}$  is named  $f$ -statistically convergent to  $l \in \mathbb{C}$  if for each  $\varepsilon > 0$ ,

$$\delta_f(A) = (\{k \in \mathbb{N} : |x_k - l| \geq \varepsilon\}) = 0.$$

Throughout the paper,  $S^f$  symbolizes the class of all  $f$ -statistically convergent sequences.

**Definition 1.2.** Suppose  $\theta = (k_r)$  is a lacunary sequence and  $f$  is an unbounded modulus. Then, the sequence  $(x_k)$  in  $\mathbb{C}$  is named  $f$ -lacunary statistically convergent (or  $S_\theta^f$ -convergent) to  $l \in \mathbb{C}$ , if

$$\lim_{r \rightarrow \infty} \frac{1}{f(h_r)} f(|\{k \in I_r : |x_k - l| \geq \varepsilon\}|) = 0$$

for every  $\varepsilon > 0$ . We write  $S_\theta^f - \lim x_k = l$  if  $(x_k)$  is  $S_\theta^f$ -convergent to  $l$ . Throughout the paper,  $S_\theta^f$  symbolizes the class of all sequences which are  $S_\theta^f$ -convergent, that is,

$$S_\theta^f = \left\{ (x_k) : \lim_{r \rightarrow \infty} \frac{1}{f(h_r)} f(|\{k \in I_r : |x_k - l| \geq \varepsilon\}|) = 0, \text{ for every } \varepsilon > 0 \right\}.$$

We write  $S_\theta$  instead of  $S_\theta^f$  in the case  $f(t) = t$  and for  $\theta = (2^r)$ , we write  $S^f$  instead of  $S_\theta^f$  and also in the particular case  $f(t) = t$  and  $\theta = (2^r)$ , we write  $S$  instead of  $S_\theta^f$ .

**Lemma 1.3.** The  $S_\theta^f$ -limit of every  $S_\theta^f$ -convergent sequence is unique.

## 2. MAIN RESULTS

In this section, we first give the definition of  $N_\theta^f$  and then establish the relations between  $N_\theta^f$  and  $N_\theta^g$ ,  $N_\theta$  and  $N_\theta^f$ ,  $N_\theta^f$  and  $S_\theta^g$ ,  $\ell_\infty \cap S_\theta^f$  and  $N_\theta^g$ , for different modulus functions  $f$  and  $g$  under some conditions on the considered

modulus functions. However, the relations between the sets  $N_\theta$  and  $S_\theta$ ,  $S_\theta$  and  $S$  are known already in [11].

**Definition 2.1.** Suppose  $\theta = (k_r)$  is a lacunary sequence and  $f$  is a modulus function. Then, the sequence  $(x_k)$  in  $\mathbb{C}$  is named strongly  $f$ -lacunary summable (or strongly  $N_\theta^f$ -summable) to  $l \in \mathbb{C}$ , if

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} \sum_{k \in I_r} f(|x_k - l|) = 0.$$

If the sequence  $(x_k)$  is strongly  $N_\theta^f$ -summable to  $l$ , we write  $N_\theta^f - \lim x_k = l$ . The class of strongly  $N_\theta^f$ -summable sequences will be symbolized by  $N_\theta^f$ , that is,

$$N_\theta^f = \left\{ (x_k) : \lim_{r \rightarrow \infty} \frac{1}{h_r} \sum_{k \in I_r} f(|x_k - l|) = 0 \text{ for some number } l \right\}.$$

Note that the modulus function  $f$  need not be unbounded in this definition.

The strong  $N_\theta^f$ -summability will reduce to the strong  $N_\theta$ -summability if we take  $f(t) = t$ .

**Theorem 2.2.** Suppose  $f$  and  $g$  are modulus functions and  $\theta = (k_r)$  is a lacunary sequence. If  $\sup_{t \in (0, \infty)} \frac{f(t)}{g(t)} < \infty$ , then  $N_\theta^g \subset N_\theta^f$ .

*Proof.* Assume that  $\alpha = \sup_{t \in (0, \infty)} \frac{f(t)}{g(t)} < \infty$ , then we have  $\frac{f(t)}{g(t)} \leq \alpha$ , and so that  $f(t) \leq \alpha g(t)$  for every  $t \in [0, \infty)$ . Now, it is clear that  $\alpha > 0$  and if  $(x_k)$  is strongly  $N_\theta^g$ -summable to some  $l \in \mathbb{C}$ , we may write

$$\frac{1}{h_r} \sum_{k \in I_r} f(|x_k - l|) \leq \alpha \frac{1}{h_r} \sum_{k \in I_r} g(|x_k - l|).$$

Taking the limits on both sides as  $r \rightarrow \infty$ , we obtain that  $(x_k) \in N_\theta^g$  implies  $(x_k) \in N_\theta^f$ .  $\square$

**Remark 2.3.** The following example shows that the inclusion  $N_\theta^g \subset N_\theta^f$  is strict at least for some special modulus functions  $f$  and  $g$ .

**EXAMPLE 2.4.** Let the lacunary sequence  $\theta = (k_r)$  be given and consider the sequence  $(x_k)$  such that  $x_k$  to be  $[\sqrt{h_r}]$  at the first  $[\sqrt{h_r}]$  integers in  $I_r$ , and  $x_k = 0$  otherwise, where  $[t]$  denotes the integer part of a real number  $t$ . Now, if we take modulus functions  $f(t) = \frac{t}{t+1}$  and  $g(t) = t$ , then  $\sup_{t \in (0, \infty)} \frac{f(t)}{g(t)} = 1 < \infty$

and so that  $N_\theta^g \subset N_\theta^f$ . By using the equality  $f(0) = 0$ , we have

$$\frac{1}{h_r} \sum_{k \in I_r} f(|x_k|) = \frac{1}{h_r} [\sqrt{h_r}] f([\sqrt{h_r}]) = \frac{[\sqrt{h_r}] [\sqrt{h_r}]}{h_r ([\sqrt{h_r}] + 1)}.$$

By taking the limits as  $r \rightarrow \infty$ , we get that  $N_\theta^f - \lim x_k = 0$  and so  $(x_k) \in N_\theta^f$ . But since

$$\frac{1}{h_r} \sum_{k \in I_r} g(|x_k|) = \frac{1}{h_r} \left[ \sqrt{h_r} \right] g \left( \left[ \sqrt{h_r} \right] \right) = \frac{\left[ \sqrt{h_r} \right] \left[ \sqrt{h_r} \right]}{h_r}$$

and  $\frac{\left[ \sqrt{h_r} \right] \left[ \sqrt{h_r} \right]}{h_r} \rightarrow 1$  as  $r \rightarrow \infty$ , we get that  $(x_k) \notin N_\theta^g$ . Hence,  $(x_k) \in N_\theta^f - N_\theta^g$  and the inclusion  $N_\theta^g \subset N_\theta^f$  is strict.

**Theorem 2.5.** Suppose  $f$  and  $g$  are modulus functions and  $\theta = (k_r)$  is a lacunary sequence. If  $\inf_{t \in (0, \infty)} \frac{f(t)}{g(t)} > 0$ , then  $N_\theta^f \subset N_\theta^g$  and the inclusion is strict.

*Proof.* Assume that  $\beta = \inf_{t \in (0, \infty)} \frac{f(t)}{g(t)} > 0$ . Then, we have  $\frac{f(t)}{g(t)} \geq \beta$  and so that  $\beta g(t) \leq f(t)$  for every  $t \in [0, \infty)$ . Now, if  $(x_k)$  is strongly  $N_\theta^f$ -summable to some  $l \in \mathbb{C}$ , we may write

$$\frac{1}{h_r} \sum_{k \in I_r} g(|x_k - l|) \leq \frac{1}{h_r} \sum_{k \in I_r} \frac{1}{\beta} f(|x_k - l|).$$

Taking the limits on both sides as  $r \rightarrow \infty$ , we obtain that  $(x_k) \in N_\theta^f$  implies  $(x_k) \in N_\theta^g$ .

For the strict inclusion, if we take the sequence of Example 2.4 with modulus functions  $f(t) = t$  and  $g(t) = \frac{t}{t+1}$ , then the strict inclusion will happen.  $\square$

The result below is obtained from Theorem 2.2 and Theorem 2.5.

**Corollary 2.6.** Suppose  $f$  and  $g$  are modulus functions and  $\theta = (k_r)$  is a lacunary sequence. If

$$0 < \inf_{t \in (0, \infty)} \frac{f(t)}{g(t)} \leq \sup_{t \in (0, \infty)} \frac{f(t)}{g(t)} < \infty,$$

then  $N_\theta^f = N_\theta^g$ .

**Theorem 2.7.** For any modulus function  $f$  and lacunary sequence  $\theta = (k_r)$ , we have  $N_\theta \subset N_\theta^f$ .

*Proof.* Assume that  $(x_k) \in N_\theta$ . So that

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} \sum_{k \in I_r} |x_k - l| = 0$$

for some  $l \in \mathbb{C}$ . Given  $\varepsilon > 0$  and choose  $\delta$  with  $0 < \delta < 1$  such that  $f(t) < \varepsilon$  for  $t \in (0, \delta]$ . Now, consider

$$\sum_{k \in I_r} f(|x_k - l|) = \sum_{\substack{k \in I_r \\ |x_k - l| \leq \delta}} f(|x_k - l|) + \sum_{\substack{k \in I_r \\ |x_k - l| > \delta}} f(|x_k - l|).$$

Since  $f(|x_k - l|) < \varepsilon$  for  $|x_k - l| \leq \delta$ , then

$$\sum_{\substack{k \in I_r \\ |x_k - l| \leq \delta}} f(|x_k - l|) < \varepsilon h_r,$$

and also for  $|x_k - l| > \delta$ , we get

$$|x_k - l| < \frac{|x_k - l|}{\delta} < 1 + \left\lceil \frac{|x_k - l|}{\delta} \right\rceil,$$

Since  $f$  is a modulus, we have

$$f(|x_k - l|) \leq f\left(1 + \left\lceil \frac{|x_k - l|}{\delta} \right\rceil\right) \leq f(1) \left(1 + \left\lceil \frac{|x_k - l|}{\delta} \right\rceil\right) \leq 2f(1) \frac{|x_k - l|}{\delta}.$$

So, we get

$$\sum_{\substack{k \in I_r \\ |x_k - l| > \delta}} f(|x_k - l|) \leq \frac{2f(1)}{\delta} \sum_{\substack{k \in I_r \\ |x_k - l| > \delta}} |x_k - l|.$$

Thus,

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} f(|x_k - l|) &\leq \varepsilon + \frac{2f(1)}{\delta} \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |x_k - l| > \delta}} |x_k - l| \\ &\leq \varepsilon + \frac{2f(1)}{\delta} \frac{1}{h_r} \sum_{k \in I_r} |x_k - l|. \end{aligned}$$

Since  $(x_k) \in N_\theta$ , we obtain that  $(x_k) \in N_\theta^f$ .  $\square$

**Corollary 2.8.** Suppose  $f$  is any modulus function and  $\theta = (k_r)$  is a lacunary sequence. If  $\inf_{t \in (0, \infty)} \frac{f(t)}{t} > 0$ , then  $N_\theta^f = N_\theta$ .

*Proof.* Since  $N_\theta \subset N_\theta^f$  for any modulus function  $f$  by Theorem 2.7, taking  $g(t) = t$  in Theorem 2.5, we get  $N_\theta^f \subset N_\theta$  if  $\inf_{t \in (0, \infty)} \frac{f(t)}{t} > 0$ . Therefore,  $N_\theta^f = N_\theta$  if  $\inf_{t \in (0, \infty)} \frac{f(t)}{t} > 0$ .  $\square$

**Theorem 2.9.** Assume that  $f$  and  $g$  are unbounded modulus functions and  $\theta = (k_r)$  is a lacunary sequence. If  $\inf_{t \in (0, \infty)} \frac{f(t)}{g(t)} > 0$  and  $\lim_{t \rightarrow \infty} \frac{g(t)}{t} > 0$ , then every strongly  $N_\theta^f$ -summable sequence is  $S_\theta^g$ -statistically convergent.

*Proof.* Suppose that  $\beta = \inf_{t \in (0, \infty)} \frac{f(t)}{g(t)} > 0$ . Then, we have  $\frac{f(t)}{g(t)} \geq \beta$  and so that  $\beta g(t) \leq f(t)$  for every  $t \in [0, \infty)$ . Now, if  $(x_k)$  is strongly  $N_\theta^f$ -summable to

some  $l \in \mathbb{C}$ , then we may write

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} f(|x_k - l|) &\geq \beta \frac{1}{h_r} \sum_{k \in I_r} g(|x_k - l|) \\ &= \beta \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |x_k - l| \geq \varepsilon}} g(|x_k - l|) + \beta \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |x_k - l| < \varepsilon}} g(|x_k - l|) \\ &\geq \beta \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |x_k - l| \geq \varepsilon}} g(|x_k - l|) \\ &\geq \beta \frac{1}{h_r} |\{k \in I_r : |x_k - l| \geq \varepsilon\}| g(\varepsilon). \end{aligned}$$

Since  $|\{k \in I_r : |x_k - l| \geq \varepsilon\}|$  is a positive integer and  $g$  is a modulus, we have

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} f(|x_k - l|) &\geq \beta \frac{1}{h_r} g(|\{k \in I_r : |x_k - l| \geq \varepsilon\}|) \frac{g(\varepsilon)}{g(1)} \\ &= \frac{g(|\{k \in I_r : |x_k - l| \geq \varepsilon\}|)}{g(h_r)} \frac{g(h_r)}{h_r} \frac{g(\varepsilon)}{g(1)} \beta. \end{aligned}$$

Taking the limits on both sides as  $r \rightarrow \infty$ , we obtain that  $(x_k) \in N_\theta^f$  implies  $(x_k) \in S_\theta^g$  since  $\lim_{t \rightarrow \infty} \frac{g(t)}{t} > 0$ . This fulfills the proof.  $\square$

*Remark 2.10.* The converse of the above theorem is not true, in general. It can be shown in the following example.

**EXAMPLE 2.11.** Let  $\theta$  be given and select the sequence  $(x_k)$  as in Example 2.4 and also consider the modulus function  $g(t) = f(t) = t$ . Then, we have  $\inf_{t \in (0, \infty)} \frac{f(t)}{g(t)} > 0$  and  $\lim_{t \rightarrow \infty} \frac{g(t)}{t} > 0$ . Given any  $\varepsilon > 0$ , we have

$$\lim_{r \rightarrow \infty} \frac{1}{g(h_r)} g(|\{k \in I_r : |x_k| \geq \varepsilon\}|) = \lim_{r \rightarrow \infty} \frac{g(\lfloor \sqrt{h_r} \rfloor)}{h_r} = 0.$$

So that  $S_\theta^g - \lim x_k = 0$  and thus  $(x_k) \in S_\theta^g$ . On the other hand,

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} \sum_{k \in I_r} f(|x_k|) = \lim_{r \rightarrow \infty} \frac{\lfloor \sqrt{h_r} \rfloor \lfloor \sqrt{h_r} \rfloor}{h_r} = 1.$$

Hence,  $N_\theta^f - \lim x_k \neq 0$  and thus  $(x_k) \notin N_\theta^f$ .

The following result is obtained by taking  $g(t) = f(t)$  in Theorem 2.9.

**Corollary 2.12.** Assume that  $f$  is an unbounded modulus function and  $\theta = (k_r)$  is a lacunary sequence. If  $\lim_{t \rightarrow \infty} \frac{f(t)}{t} > 0$ , then every strongly  $N_\theta^f$ -summable sequence is  $S_\theta^f$ -statistically convergent.

The following result is obtained by taking  $g(t) = t$  in Theorem 2.9.

**Corollary 2.13.** Assume that  $f$  is an unbounded modulus function and  $\theta = (k_r)$  is a lacunary sequence. If  $\inf_{t \in (0, \infty)} \frac{f(t)}{t} > 0$ , then every strongly  $N_\theta^f$ -summable sequence is  $S_\theta$ -statistically convergent.

The following result is obtained by taking  $f(t) = t$  in Corollary 2.13, which is also the first part of Theorem 1 of [11].

**Corollary 2.14.** *A strongly  $N_\theta$ -summable sequence is  $S_\theta$ -statistically convergent.*

**Theorem 2.15.** *Suppose  $\theta = (k_r)$  is a lacunary sequence. Then, for any unbounded modulus functions  $f$  and  $g$ , we have  $\ell_\infty \cap S_\theta^f \subset N_\theta^g$ .*

*Proof.* Let  $f$  and  $g$  be unbounded modulus functions and  $\theta = (k_r)$  be given. Since  $S_\theta^f \subset S_\theta$  for every modulus  $f$  by Theorem 11 of [4], and since  $\ell_\infty \cap S_\theta \subset N_\theta$  by the second part of Theorem 1 of [11], then we have  $\ell_\infty \cap S_\theta^f \subset \ell_\infty \cap S_\theta \subset N_\theta$ , that is,  $\ell_\infty \cap S_\theta^f \subset N_\theta$ . On the other hand,  $N_\theta \subset N_\theta^g$  for any modulus  $g$  by Theorem 2.7. Therefore,  $\ell_\infty \cap S_\theta^f \subset N_\theta^g$ .  $\square$

*Remark 2.16.* The following example shows that the inclusion  $\ell_\infty \cap S_\theta^f \subset N_\theta^g$  is strict at least for some special modulus functions  $f$  and  $g$ .

**EXAMPLE 2.17.** As an example, let the lacunary sequence  $\theta = (k_r)$  be provided and consider the sequence  $(x_k)$  such that  $x_k$  to be  $\sqrt[3]{h_r}$  at the first  $[\sqrt{h_r}]$  integers in  $I_r$ , and  $x_k = 0$  otherwise. Now, if we take modulus functions  $f(t) = \frac{t}{t+1}$  and  $g(t) = t$ , then for every  $\varepsilon > 0$ , we have

$$\lim_{r \rightarrow \infty} \frac{1}{f(h_r)} f(|\{k \in I_r : |x_k| \geq \varepsilon\}|) = \lim_{r \rightarrow \infty} \frac{f([\sqrt{h_r}])}{f(h_r)} = 1.$$

So that  $(x_k) \notin S_\theta^f$ , but since

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} \sum_{k \in I_r} g(|x_k|) = \lim_{r \rightarrow \infty} \frac{[\sqrt{h_r}] g(\sqrt[3]{h_r})}{h_r} = 0,$$

we get  $(x_k) \in N_\theta^g$ .

The outcome below is a result of Theorem 2.15.

**Corollary 2.18.** *Suppose  $\theta = (k_r)$  is a lacunary sequence and  $f$  is any unbounded modulus function. Then,*

- (1)  $\ell_\infty \cap S_\theta^f \subset N_\theta^f$ .
- (2)  $\ell_\infty \cap S_\theta^f \subset N$ .
- (3)  $\ell_\infty \cap S \subset N_\theta^f$ .

#### ACKNOWLEDGMENTS

The authors wish to thank the referees for their kindness and generosity suggestions and comments.

#### REFERENCES

1. A. Aizpuru, M. C. Listán-García, F. Rambla-Barreno, Density by Moduli and Statistical Convergence, *Quaestiones Mathematicae*, **37**(4), (2014), 525–530.
2. M. Altınok, M. Küçükaslan, A-Statistical Supremum-infinum and A-statistical Convergence, *Azerbaijan Journal of Mathematics*, **4**(2), (2014), 31–42.



3. V. K. Bhardwaj, N. Singh, On Some Sequence Spaces Defined by a Modulus, *Indian J. Pure Appl. Math.*, **30**(8), (1999), 809–817.
4. V. K. Bhardwaj, S. Dhawan, Density by Moduli and Lacunary Statistical Convergence, *Hindawi*, **2016**, (2016), 1–11.
5. B. T. Bilalov, T. Y. Nazarova, Statistical Convergence of Functional Sequences, *Rocky Mountain J. Math.*, **45**(5), (2015), 1413–1423.
6. B. T. Bilalov, T. Y. Nazarova, On Statistical Convergence in Metric Spaces, *Journal of Mathematics Research*, **7**(1), (2015), 37–43.
7. B. T. Bilalov, T. Y. Nazarova, On Statistical Type Convergence in Uniform Spaces, *Bull. Iranian Math. Soc.*, **42**(4), (2016), 975–986.
8. J. Connor, The Statistical and Strong  $p$ -Cesaro Convergence of Sequences, *Analysis*, **8**(1–2), (1988), 47–64.
9. M. Et, Strongly Almost Summable Difference Sequences of Order  $m$  Defined by a Modulus, *Studia Scientiarum Mathematicarum Hungarica*, **40**(4), (2003), 463–476.
10. H. Fast, Sur la Convergence Statistique, *Colloquium Mathematicae*, **2**(3–4), (1951), 241–244.
11. J. A. Fridy, C. Orhan, Lacunary Statistical Convergence, *Pacific Journal of Mathematics*, **160**(1), (1993), 43–51.
12. J. A. Fridy, On Statistical Convergence, *Analysis*, **5**(4), (1985), 301–314.
13. D. Ghosh, P. D. Srivastava, On Some Vector Valued Sequence Spaces Defined Using a Modulus Function, *Indian J. Pure Appl. Math.*, **30**(8), (1999), 819–826.
14. I. S. Ibrahim, R. Çolak, On Strong Lacunary Summability of Order  $\alpha$  with Respect to Modulus Functions, *Annals of the University of Craiova-Mathematics and Computer Science Series*, **1**(48), (2021), 127–136.
15. I. J. Maddox, Sequence Spaces Defined by a Modulus, *Mathematical Proceedings of the Cambridge Philosophical Society*, **100**(1), (1986), 161–166.
16. I. J. Maddox, Inclusions Between FK Spaces and Kuttner's Theorem, *Mathematical Proceedings of the Cambridge Philosophical Society*, **101**(3), (1987), 523–527.
17. H. Nakano, Concave Modulans, *Journal of the Mathematical Society of Japan*, **5**(1), (1953), 29–49.
18. D. Rath, B. C. Tripathy, On Statistically Convergent and Statistically Cauchy Sequences, *Indian Journal of Pure and Applied Mathematics*, **25**, (1994), 381–381.
19. W. H. Ruckle, FK Spaces in which the Sequence of Coordinate Vectors is Bounded, *Canadian Journal of Mathematics*, **25**(5), (1973), 973–978.
20. T. Šalát, On Statistically Convergent Sequences of Real Numbers, *Mathematica Slovaca*, **30**(2), (1980), 139–150.
21. I. J. Schoenberg, The Integrability of Certain Functions and Related Summability Methods, *The American Mathematical Monthly*, **66**(5), (1959), 361–775.
22. Zygmund. A, *Trigonometric Series*, 2nd edn, Cambridge University Press, 1979.