

## Strongly Almost Ideal Convergent Sequences in a Locally Convex Space Defined by Musielak-Orlicz Function

Bipan Hazarika\*

Department of Mathematics, Rajiv Gandhi University, Rono Hills,  
Doimukh-791112, Arunachal Pradesh, India.

E-mail: bh\_rgu@yahoo.co.in

ABSTRACT. In this article, we introduce a new class of ideal convergent sequence spaces using an infinite matrix, Musielak-Orlicz function and a new generalized difference matrix in locally convex spaces. We investigate some linear topological structures and algebraic properties of these spaces. We also give some relations related to these sequence spaces.

**Keywords:**  $I$ -convergence, Difference space, Musielak-Orlicz function.

**2000 Mathematics subject classification:** 40A05, 40B50, 46A19, 46A45.

### 1. INTRODUCTION

Kostyrko et al., [25] introduced the notion of  $I$ -convergence ( $I$  denotes the ideal of the subsets of the set  $\mathbb{N}$  of positive integers), which is a generalization of statistical convergence (see [14, 35]) and further studied by many others (see [6, 19, 20, 38, 39, 40]). Recently, Hazarika [21] introduced the notion generalized difference ideal convergent sequences and studied some interesting results. Quite recently, Esi [11] introduced strongly almost ideal convergent sequence spaces in 2-normed spaces defined by an Orlicz function and prove some results related to this notion.

---

\*Corresponding Author

Before proceeding let us recall a few concepts, which we shall use throughout this paper.

Let  $X$  be a non-empty set, then a family of sets  $I \subset 2^X$  (the class of all subsets of  $X$ ) is called an *ideal* if and only if for each  $A, B \in I$  we have  $A \cup B \in I$  and for each  $A \in I$  and each  $B \subset A$  we have  $B \in I$ . A non-empty family of sets  $F \subset 2^X$  is a *filter* on  $X$  if and only if  $\phi \notin F$  for each  $A, B \in F$  we have  $A \cap B \in F$  and each  $A \in F$  and each  $B \supset A$  we have  $B \in F$ . An ideal  $I$  is called non-trivial ideal if  $I \neq \phi$  and  $X \notin I$ . Clearly  $I \subset 2^X$  is a non-trivial ideal if and only if  $F = F(I) = \{X - A : A \in I\}$  is a filter on  $X$ . A non-trivial ideal  $I \subset 2^X$  is called admissible if and only if  $\{\{x\} : x \in X\} \subset I$ . A non-trivial ideal  $I$  is maximal if there cannot exist any non-trivial ideal  $J \neq I$  containing  $I$  as a subset. Further details on ideals of  $2^X$  can be found in Kostyrko et al., [25]. Recall that a sequence  $x = (x_k)$  of points in  $\mathbb{R}$  is said to be  $I$ -convergent to a real number  $\ell$  if  $\{k \in \mathbb{N} : |x_k - \ell| \geq \varepsilon\} \in I$  for every  $\varepsilon > 0$  ([25]). In this case we write  $I - \lim x_k = \ell$ .

Throughout the article  $w, \ell_\infty, c, c_0$ , denote the classes of *all, bounded, convergent, null* sequences of complex numbers, respectively.

The notion of difference sequence space was introduced by Kizmaz [24], who studied the difference sequence spaces  $\ell_\infty(\Delta), c(\Delta), c_0(\Delta)$ . The notion was further generalized by Et and Colak [12] introducing the sequence spaces  $\ell_\infty(\Delta^p), c(\Delta^p), c_0(\Delta^p)$ . For a non negative integer  $p$ , the generalized difference sequence spaces are defined as follows. For a given sequence space  $Z$  we have

$$Z(\Delta^p) = \{x = (x_k) \in w : (\Delta^p x_k) \in Z\},$$

where  $\Delta^p x_k = \Delta^{p-1} x_k - \Delta^{p-1} x_{k+1}$ ,  $\Delta^0 x_k = x_k$ , for all  $k \in \mathbb{N}$ , the difference operator is equivalent to the following binomial representation:

$$\Delta^p x_k = \sum_{\nu=0}^p (-1)^\nu \binom{p}{\nu} x_{k+\nu} \text{ for all } k \in \mathbb{N}.$$

Taking  $p = 1$ , the spaces  $\ell_\infty(\Delta), c(\Delta), c_0(\Delta)$ , introduced and studied by Kizmaz [24].

Tripathy and Esi [36] introduced and studied the new type of generalized difference sequence spaces

$$Z(\Delta_i) = \{(x_k) \in w : \Delta_i x_k \in Z\},$$

for  $Z = \ell_\infty, c, c_0$  where  $\Delta_i x = (\Delta_i x_k) = (x_k - x_{k+i})$  for all  $k, i \in \mathbb{N}$ .

Tripathy et al., [37] further generalized this notion and introduced the following sequence spaces. For  $p \geq 1$  and  $i \geq 1$ ,

$$Z(\Delta_i^p) = \{(x_k) \in w : \Delta_i^p x_k \in Z\},$$

for  $Z = \ell_\infty, c, c_0$ . This generalized difference has the following binomial representation,

$$\Delta_i^p x_k = \sum_{\nu=0}^n (-1)^\nu \binom{p}{\nu} x_{k+i\nu} \quad \text{for all } k \in \mathbb{N}.$$

Dutta [10] introduced the following difference sequence spaces

$$Z(\Delta_{(i)}^p) = \{(x_k) \in w : \Delta_{(i)}^p x_k \in Z\} \quad \text{for all } p, i \in \mathbb{N},$$

for  $Z = \ell_\infty, \bar{c}, \bar{c}_0$  where  $\bar{c}, \bar{c}_0$  are the sets of statistically convergent and statistically null sequences, respectively, and  $\Delta_{(i)}^p x = (\Delta_{(i)}^p x_k) = (\Delta_{(i)}^{p-1} x_k - \Delta_{(i)}^{p-1} x_{k-i})$  and  $\Delta_{(i)}^0 x_k = x_k$  for all  $k \in \mathbb{N}$ , which is equivalent to the following binomial representation:

$$\Delta_{(i)}^p x_k = \sum_{\nu=0}^p (-1)^\nu \binom{p}{\nu} x_{k-i\nu}.$$

Basar and Altay [3] introduced the generalized difference matrix  $B(r, s) = (b_{pk}(r, s))$  which is a generalization of  $\Delta_{(1)}^1$ -difference operator as follows:

$$b_{pk}(r, s) = \begin{cases} r, & \text{if } k = p; \\ s, & \text{if } k = p - 1; \\ 0, & \text{if } 0 \leq k < p - 1 \text{ or } k > p. \end{cases}$$

for all  $k, p \in \mathbb{N}, r, s \in \mathbb{R} - \{0\}$ .

Basarir and Kayikci [4] have defined the generalized difference matrix  $B^p$  of order  $p$ , which reduced the difference operator  $\Delta_{(1)}^p$  in case  $r = 1, s = -1$  and the binomial representation of this operator is

$$B^p x_k = \sum_{\nu=0}^p \binom{p}{\nu} r^{p-\nu} s^\nu x_{k-\nu},$$

where  $r, s \in \mathbb{R} - \{0\}$  and  $p \in \mathbb{N}$ .

Recently Basarir et al., [5] introduced the following generalized difference sequence spaces

$$Z(B_{(i)}^p) = \{(x_k) \in w : B_{(i)}^p x_k \in Z\} \quad \text{for all } p, i \in \mathbb{N},$$

for  $Z = \ell_\infty, \bar{c}, \bar{c}_0$  where  $\bar{c}, \bar{c}_0$  are the sets of statistically convergent and statistically null sequences, respectively, and  $B_{(i)}^p x = (B_{(i)}^p x_k) = (rB_{(i)}^{p-1} x_k +$

$sB_{(i)}^{p-1}x_{k-i}$ ) and  $B_{(i)}^0x_k = x_k$  for all  $k \in \mathbb{N}$ , which is equivalent to the following binomial representation:

$$B_{(i)}^p x_k = \sum_{\nu=0}^p \binom{p}{\nu} r^{p-\nu} s^\nu x_{k-i\nu}.$$

Let  $X$  and  $Y$  be two nonempty subsets of the space  $w$  of complex sequences. Let  $A = (a_{nk}), (n, k = 1, 2, 3, \dots)$  be an infinite matrix of complex numbers. We write  $Ax = (A_n(x))$  if  $A_n(x) = \sum_{k=1}^{\infty} a_{nk}x_k$  converges for each  $n$ . If  $x = (x_k) \in X \Rightarrow Ax = (A_n(x)) \in Y$  we say that  $A$  defines a (matrix) transformation from  $X$  to  $Y$  and we denote it by  $A : X \rightarrow Y$ .

A sequence  $x = (x_k) \in \ell_\infty$  is said to be almost convergent if all of its Banach limits coincide. Let  $\widehat{c}$  denotes the space of all almost convergent sequences.

Lorentz [29] introduced the following sequence space.

$$\widehat{c} = \left\{ x \in \ell_\infty : \lim_k t_{m,k}(x) \text{ exists uniformly in } m \right\}$$

where  $t_{m,k}(x) = \frac{x_k + x_{k+1} + \dots + x_{k+m}}{m+1}$ .

The following space of strongly almost convergent sequences was introduced by Maddox [30],

$$[\widehat{c}] = \left\{ x \in \ell_\infty : \lim_k t_{m,k}(|x - Le|) \text{ exists uniformly in } m, \text{ for some } L \right\}$$

where  $e = (1, 1, 1, \dots)$ .

It is clear that

$$t_{m,k}(x) = \begin{cases} \frac{1}{m+1} \sum_{i=1}^m x_{k+i} & \text{for } m \geq 1; \\ x_k & \text{for } m = 0 \end{cases}$$

An Orlicz function is a function  $M : [0, \infty) \rightarrow [0, \infty)$ , which is continuous, non-decreasing and convex with  $M(0) = 0, M(x) > 0$  as  $x > 0$  and  $M(x) \rightarrow \infty$  as  $x \rightarrow \infty$  (see [26]).

An Orlicz function  $M$  can always be represented in the following integral form:

$$M(x) = \int_0^x p(t)dt$$

where  $p$  is the known kernel of  $M$ , right differentiable for  $t \geq 0, p(0) = 0, p(t) > 0$  for  $t > 0$  and  $p(t) \rightarrow \infty$  as  $t \rightarrow \infty$ .

If convexity of Orlicz function is replaced by  $M(x+y) \leq M(x) + M(y)$  then this function is called the modulus function and characterized by Ruckle [34]. An Orlicz function  $M$  is said to satisfy  $\Delta_2$ -condition for all values of  $u$ , if there exists a constant  $K > 0$  such that  $M(2u) \leq KM(u)$ ,  $u \geq 0$ .

Let  $M$  be an Orlicz function which satisfies  $\Delta_2$ -condition and let  $0 < \delta < 1$ . Then for each  $t \geq \delta$ , we have  $M(t) < K\delta^{-1}M(2)$  for some constant  $K > 0$ .

Two Orlicz functions  $M_1$  and  $M_2$  are said to be *equivalent* if there exist positive constants  $\alpha, \beta$  and  $x_0$  such that

$$M_1(\alpha) \leq M_2(x) \leq M_1(\beta)$$

for all  $x$  with  $0 \leq x < x_0$ .

Lindenstrauss and Tzafriri [28] studied some Orlicz type sequence spaces defined as follows:

$$\ell_M = \left\{ (x_k) \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty, \text{ for some } \rho > 0 \right\}.$$

The space  $\ell_M$  with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}$$

becomes a Banach space which is called an Orlicz sequence space. The space  $\ell_M$  is closely related to the space  $\ell_p$  which is an Orlicz sequence space with  $M(t) = |t|^p$  for  $1 \leq p < \infty$ .

A sequence  $\mathbf{M} = (M_k)$  of Orlicz functions is called a *Musielak-Orlicz function* (for details see [9, 18, 22, 23]). Also a Musielak-Orlicz function  $\phi = (\phi_k)$  is called a *complementary function* of a Musielak-Orlicz function  $\mathbf{M}$  if

$$\phi_k(t) = \sup\{|t|s - M_k(s) : s \geq 0\}, \text{ for } k = 1, 2, 3, \dots$$

For a given Musielak-Orlicz function  $\mathbf{M}$ , the Musielak-Orlicz sequence space  $l_{\mathbf{M}}$  and its subspace  $h_{\mathbf{M}}$  are defined as follows:

$$l_{\mathbf{M}} = \{x = (x_k) \in w : I_{\mathbf{M}}(cx) < \infty, \text{ for some } c > 0\};$$

$$h_{\mathbf{M}} = \{x = (x_k) \in w : I_{\mathbf{M}}(cx) < \infty, \text{ for all } c > 0\},$$

where  $I_{\mathbf{M}}$  is a convex modular defined by

$$I_{\mathbf{M}} = \sum_{k=1}^{\infty} M_k(x_k), x = (x_k) \in l_{\mathbf{M}}.$$

We consider  $l_M$  equipped with the Luxemburg norm

$$\|x\| = \inf \left\{ k > 0 : I_M \left( \frac{x}{k} \right) \leq 1 \right\}$$

or equipped with the Orlicz norm

$$\|x\|^0 = \inf \left\{ \frac{1}{k} (1 + I_M(kx)) : k > 0 \right\}.$$

The following well-known inequality will be used throughout the article. Let  $p = (p_k)$  be any sequence of positive real numbers with  $0 \leq p_k \leq \sup_k p_k = G$ ,  $D = \max\{1, 2^{G-1}\}$  then

$$|a_k + b_k|^{p_k} \leq D(|a_k|^{p_k} + |b_k|^{p_k})$$

for all  $k \in \mathbb{N}$  and  $a_k, b_k \in \mathbb{C}$ . Also  $|a|^{p_k} \leq \max\{1, |a|^G\}$  for all  $a \in \mathbb{C}$ .

Subsequently Orlicz function was used to define sequence spaces by Parashar and Choudhary [33] and many others (see [2, 27, 31, 41]).

**Remark 1.1.** It is well known if  $M$  is a convex function and  $M(0) = 0$ , then  $M(\lambda x) \leq \lambda M(x)$ , for all  $\lambda$  with  $0 < \lambda < 1$ .

**Definition 1.2.** A sequence space  $E$  is said to be *solid (or normal)* if  $(\alpha_k x_k) \in E$ , whenever  $(x_k) \in E$  and for all sequence  $(\alpha_k)$  of scalars with  $|\alpha_k| \leq 1$ , for all  $k \in \mathbb{N}$ .

Let  $K = \{k_1 < k_2 < \dots\} \subseteq \mathbb{N}$  and  $E$  be a sequence space. A  $K$ -step space of  $E$  is a sequence space  $\lambda_K^E = \{(x_{k_n}) \in w : (k_n) \in E\}$ .

A canonical preimage of a sequence  $\{(x_{k_n})\} \in \lambda_K^E$  is a sequence  $\{y_n\} \in w$  defined as

$$y_k = \begin{cases} x_k, & \text{if } k \in K \\ 0, & \text{otherwise.} \end{cases}$$

A canonical preimage of a step space  $\lambda_K^E$  is a set of canonical preimages of all elements in  $\lambda_K^E$ , i.e.  $y$  is in canonical preimage of  $\lambda_K^E$  if and only if  $y$  is canonical preimage of some  $x \in \lambda_K^E$ .

**Definition 1.3.** A sequence space  $E$  is said to be *monotone* if it contains the canonical preimages of its step spaces.

**Lemma 1.1.** *Every normal space is monotone.*

Throughout this paper  $X$  we denote a locally convex Hausdorff topological linear space whose topology is determined by a set  $Q$  of continuous seminorms  $q$ . Also we denote  $I$  is an non-trivial admissible ideal of  $\mathbb{N}$ .

## 2. IDEAL CONVERGENCE IN A LOCALLY CONVEX SPACE

In this section we define  $I$ -convergence and almost  $I$ -convergence in a locally convex space  $X$  and investigate some basic properties.

**Definition 2.1.** A sequence  $x = (x_k)$  in  $X$  is said to be  $I$ -convergent to  $\ell \in X$  if for all  $q \in Q$  and all  $\varepsilon > 0$ ,

$$\{k \in \mathbb{N} : q(x_k - \ell) \geq \varepsilon\} \in I.$$

In this case we can write  $I_q\text{-}\lim x_k = \ell$ . We denote  $I_q = \{k \in \mathbb{N} : q(x_k - \ell) \geq \varepsilon\}$ .

Further, since  $X$  is Hausdorff, the limit of ideal convergent sequence is unique.

**Remark 2.1.** We can introduced this concept in TVS-cone Normed Spaces (for details on TVS-cone Normed Spaces see [32]) and in 2-inner Product Spaces (for details on 2-inner Product Spaces see [1]).

**Definition 2.2.** A sequence  $x = (x_k)$  in  $X$  is said to be almost  $I$ -convergent to  $\ell \in X$  if for all  $q \in Q$  and all  $\varepsilon > 0$ ,

$$\{k \in \mathbb{N} : q(t_{m,k}(x) - \ell) \geq \varepsilon\} \in I \text{ for all } m \in \mathbb{N}.$$

In this case we can write  $\widehat{I}_q\text{-}\lim t_{m,k}(x) = \ell$ . We denote  $\widehat{I}_q = \{k \in \mathbb{N} : q(t_{m,k}(x) - \ell) \geq \varepsilon\}$  for all  $m \in \mathbb{N}$ .

**Definition 2.3.** Let  $\mathbf{M}$  be a Musielak-Orlicz function. We say that a sequence  $x = (x_k)$  in  $\widehat{w}^I(\mathbf{M})$  if and only if there exists  $\ell \in X$  such that for all  $q \in Q$  and for every  $\varepsilon > 0$ ,

$$\left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{\rho} \right) \right] \geq \varepsilon \right\} \in I \text{ for } \rho > 0, \text{ for all } m \in \mathbb{N}. \quad (2.1)$$

When (2.1) holds we write

$$x_k \rightarrow \ell((\widehat{w}^I(\mathbf{M}))).$$

The condition (2.1) provides a definition of strong ideal summability for a sequence in a locally convex space.

**Theorem 2.1.** *Let  $A = (a_{nk})$  be a non-negative regular matrix and  $u = (u_k)$  be a bounded sequence of strictly positive real numbers. Let  $\mathbf{M}$  be a Musielak-Orlicz function. Then  $x_k \rightarrow \ell(\widehat{w}(\mathbf{M}, A, u))$  implies that  $x_k \rightarrow \ell(\widehat{I}_q(A))$ .*

*Proof.* Let  $q \in Q$ . Assume that  $x_k \rightarrow \ell(\widehat{w}(\mathbf{M}, A, u))$ , then for  $\rho > 0$  we have

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{\rho} \right) \right]^{u_k} = 0 \text{ for } \ell \in \mathbb{C}, \text{ for all } m \in \mathbb{N}.$$

Let  $\varepsilon > 0$  be given. For all  $m \in \mathbb{N}$ . We define

$$K(\varepsilon) = \{k \in \mathbb{N} : q(t_{m,k}(x) - \ell) \geq \varepsilon\}$$

and we write

$$\begin{aligned} & \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{t_k} \\ = & \sum_{k \in K(\varepsilon)} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k} + \sum_{k \notin K(\varepsilon)} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k} \\ & \geq \left( \sum_{k \in K(\varepsilon)} a_{nk} \right) \left[ M_k \left( \frac{\varepsilon}{r} \right) \right]^{u_k}. \end{aligned}$$

Then we have  $x_k \rightarrow \ell(\widehat{I}_q(A))$ .  $\square$

**Theorem 2.2.** *Let  $A = (a_{nk})$  be a non-negative regular matrix and  $u = (u_k)$  be a bounded sequence of strictly positive real numbers. Let  $\mathbf{M}$  be a Musielak-Orlicz function. If  $x = (x_k) \in \ell_{\infty}$  and  $x_k \rightarrow \ell(\widehat{I}_q(A))$ , then  $x_k \rightarrow \ell(\widehat{w}(\mathbf{M}, A, u))$ .*

*Proof.* Suppose that  $x = (x_k) \in \ell_{\infty}$  and  $x_k \rightarrow \ell(\widehat{I}_q(A))$ . Then there is a set  $K \in F(\widehat{I}_q)$  such that

$$\lim_{k \in K} q(t_{m,k}(x) - \ell) = 0 \text{ for all } m \in \mathbb{N}.$$

Now

$$\begin{aligned} & \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k} \\ = & \sum_{k \in K(\varepsilon)} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k} + \sum_{k \notin K(\varepsilon)} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k} \\ = & \sum_{k=1}^{\infty} a_{nk} \chi_K(k) \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k} + \sum_{k=1}^{\infty} a_{nk} \chi_{K^c}(k) \left[ M_k \left( \frac{q(t_{m,k}(x) - \ell)}{r} \right) \right]^{u_k}. \end{aligned}$$

If we consider the regularity of  $A$ ,  $K^c \in \widehat{I}_q$  and boundedness of  $(x_k)$ , the right side tends to zero. Hence  $x_k \rightarrow \ell(\widehat{w}(\mathbf{M}, A, u))$ .  $\square$



## 3. STRONGLY IDEAL CONVERGENT SEQUENCES IN A LOCALLY CONVEX SPACE

In this section we define some new classes of strongly  $I$ -convergent sequences by using infinite matrix in a locally convex space  $X$  and investigate their linear topological structures. Also we find out some relations related to these spaces.

Recall that a mapping  $g : X \rightarrow \mathbb{R}$  is called a *paranorm* on  $X$  if it satisfies the following conditions:

- (i)  $g(\theta) = 0$  where  $\theta$  is the zero element of the space;
- (ii)  $g(x) = g(-x)$ ;
- (iii)  $g(x + y) \leq g(x) + g(y)$ ;
- (iv)  $\lambda^n \rightarrow \lambda (n \rightarrow \infty)$  and  $g(x^n - x) \rightarrow 0 (n \rightarrow \infty)$  imply  $g(\lambda^n x^n - \lambda x) \rightarrow 0 (n \rightarrow \infty)$  for all  $x, y \in X$ . The ordered pair  $(X; g)$  is called a paranormed space with respect to the paranorm  $g$ .

The main aim of this article is to introduce the following sequence spaces and examine some properties of the resulting sequence spaces.

Let  $I$  be an admissible ideal of  $\mathbb{N}$ ,  $u = (u_k)$  be a bounded sequence of strictly positive real numbers and  $A = (a_{nk})$  be an infinite matrix. Let  $\mathbf{M}$  be a Musielak-Orlicz function. Further  $w(X)$  denotes the space of all  $X$ -valued sequences. For each  $\varepsilon > 0$ , for all  $q \in Q$  and for  $\rho > 0$  we define the following sequence spaces.

$$\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q) =$$

$$\left\{ (x_k) \in w(X) : \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - \ell)}{\rho} \right) \right]^{u_k} \geq \varepsilon \right\} \in I \text{ for } \ell \in X, \text{ for all } m \in \mathbb{N} \right\},$$

$$\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q) =$$

$$\left\{ (x_k) \in w(X) : \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} \geq \varepsilon \right\} \in I \text{ for all } m \in \mathbb{N} \right\},$$

$$\widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q) =$$

$$\left\{ (x_k) \in w(X) : \exists K > 0 \text{ s.t. } \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} \geq K \right\} \in I \text{ for all } m \in \mathbb{N} \right\},$$

$$\widehat{w}_\infty(A, B_{(i)}^p, \mathbf{M}, u, q) =$$

$$\left\{ (x_k) \in w(X) : \sup_{n \in \mathbb{N}} \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} < \infty \text{ for all } m \in \mathbb{N} \right\}.$$

Some classes are obtained by specializing  $p$ ,  $A$ ,  $\mathbf{M}$  and  $u = (u_k)$  for all  $k \in \mathbb{N}$ . Here are some examples.

- (i) If  $p = 1$ , then above spaces are denoted by  $\widehat{w}^I(A, B_{(i)}, \mathbf{M}, u, q)$ ,  $\widehat{w}_0^I(A, B_{(i)}, \mathbf{M}, u, q)$ ,  $\widehat{w}_\infty^I(A, B_{(i)}, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty(A, B_{(i)}, \mathbf{M}, u, q)$ .

- (ii) If  $i = 1$  then above spaces are denoted by  $\widehat{w}^I(A, B^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_0^I(A, B^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_\infty^I(A, B^p, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty(A, B^p, \mathbf{M}, u, q)$ .
- (iii) If  $M_k(x) = x$  for all  $x \in [0, \infty)$ ,  $k \in \mathbf{N}$  then we obtain the above spaces as  $\widehat{w}^I(A, B_{(i)}^p, u, q)$ ,  $\widehat{w}_0^I(A, B_{(i)}^p, u, q)$ ,  $\widehat{w}_\infty^I(A, B_{(i)}^p, u, q)$  and  $\widehat{w}_\infty(A, B_{(i)}^p, u, q)$ .
- (iv) If  $u = (u_k) = (1, 1, 1, \dots)$ , then above spaces are denoted by  $\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, q)$ ,  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, q)$ ,  $\widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, q)$  and  $\widehat{w}_\infty(A, B_{(i)}^p, \mathbf{M}, q)$ .
- (v) If we take  $A = (C, 1)$ , i.e., the Cesàro matrix, then the above classes of sequences are denoted by  $\widehat{w}^I(B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_0^I(B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_\infty^I(B_{(i)}^p, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty(B_{(i)}^p, \mathbf{M}, u, q)$ .
- (vi) If we take  $A = (a_{nk})$  is a de la Vallée Poussin mean, i.e.,

$$a_{nk} = \begin{cases} \frac{1}{\lambda_n}, & \text{if } k \in I_n = [n - \lambda_n + 1, n]; \\ 0, & \text{otherwise.} \end{cases}$$

where  $(\lambda_n)$  is a non-decreasing sequence of positive numbers tending to  $\infty$  and  $\lambda_{n+1} \leq \lambda_n + 1$ ,  $\lambda_1 = 1$ , then the above classes of sequences are denoted by  $\widehat{w}^I(\lambda, B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_0^I(\lambda, B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_\infty^I(\lambda, B_{(i)}^p, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty(\lambda, B_{(i)}^p, \mathbf{M}, u, q)$ .

- (vii) By a lacunary sequence  $\theta = (k_r)$ , where  $k_0 = 0$ , we shall mean an increasing sequence of non-negative integers with  $k_r - k_{r-1} \rightarrow \infty$  as  $r \rightarrow \infty$ . The intervals determined by  $\theta$  will be denoted by  $J_r = (k_{r-1}, k_r]$  and we let  $h_r = k_r - k_{r-1}$ . As a final illustration let

$$a_{nk} = \begin{cases} \frac{1}{h_r}, & \text{if } k \in I_r = (k_{r-1}, k_r]; \\ 0, & \text{otherwise.} \end{cases}$$

Then the above classes of sequences are denoted by  $\widehat{w}^I(\theta, B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_0^I(\theta, B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_\infty^I(\theta, B_{(i)}^p, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty(\theta, B_{(i)}^p, \mathbf{M}, u, q)$ .

**Theorem 3.1.**  $\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ ,  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  are topological linear spaces.

*Proof.* We will prove the result for the space  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  only and the others can be proved in similar way. Let  $x = (x_k)$  and  $y = (y_k)$  be two elements in  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . Then there exist  $\rho_1 > 0$  and  $\rho_2 > 0$  such that

$$A_{\frac{\varepsilon}{2}} = \left\{ n \in \mathbf{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho_1} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \in I$$

and

$$B_{\frac{\varepsilon}{2}} = \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(y)) \right)}{\rho_2} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \in I.$$

Let  $\alpha, \beta$  be two scalars in  $\mathbb{R}$ . Since  $B_{(i)}^p$  is linear and the continuity of the Musielak-Orlicz function  $\mathbf{M}$ , the following inequality holds:

$$\begin{aligned} & \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(\alpha x + \beta y)) \right)}{|\alpha|\rho_1 + |\beta|\rho_2} \right) \right]^{u_k} \\ & \leq D \sum_{k=1}^{\infty} a_{nk} \left[ \frac{|\alpha|}{|\alpha|\rho_1 + |\beta|\rho_2} M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho_1} \right) \right]^{u_k} \\ & \quad + D \sum_{k=1}^{\infty} a_{nk} \left[ \frac{|\beta|}{|\alpha|\rho_1 + |\beta|\rho_2} M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(y)) \right)}{\rho_2} \right) \right]^{u_k} \\ & \leq DK \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho_1} \right) \right]^{p_k} \\ & \quad + DK \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(y)) \right)}{\rho_2} \right) \right]^{u_k}, \end{aligned}$$

where  $K = \max\left\{1, \left(\frac{|\alpha|\rho_1}{|\alpha|\rho_1 + |\beta|\rho_2}\right), \left(\frac{|\beta|\rho_2}{|\alpha|\rho_1 + |\beta|\rho_2}\right)\right\}$ .

From the above relation we get

$$\begin{aligned} & \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(\alpha x + \beta y)) \right)}{(|\alpha|\rho_1 + |\beta|\rho_2)} \right) \right]^{u_k} \geq \varepsilon \right\} \\ & \subseteq \left\{ n \in \mathbb{N} : DK \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho_1} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \\ & \cup \left\{ n \in \mathbb{N} : DK \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(y)) \right)}{\rho_2} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\}. \quad (3.1) \end{aligned}$$

Since both of the sets on the right hand of (3.1) are belong to  $I$ , this completes the proof.  $\square$

**Remark 3.2.** It is easy to verify that the space  $\widehat{w}_\infty(A, B_{(i)}^p, \mathbf{M}, u, q)$  is a linear space.

**Theorem 3.3.** Let  $\mathbf{S} = (S_k)$  and  $\mathbf{T} = (T_k)$  be Musielak-Orlicz functions. Then the following holds:

$$\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{S}, u, q) \cap \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{T}, u, q) \subseteq \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{S} + \mathbf{T}, u, q).$$

*Proof.* Let  $x = (x_k) \in \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{S}, u, q) \cap \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{T}, u, q)$ . Then the result follows from the inequality

$$\begin{aligned} & \sum_{k=1}^{\infty} a_{nk} \left[ (S_k + T_k) \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} \\ \leq & D \sum_{k=1}^{\infty} a_{nk} \left[ S_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} + D \sum_{k=1}^{\infty} a_{nk} \left[ T_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{p_k}. \end{aligned}$$

□

**Theorem 3.4.** Let  $\mathbf{S} = (S_k)$  and  $\mathbf{T} = (T_k)$  be Musielak-Orlicz functions. Then the following holds:

$$\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{T}, u, q) \subseteq \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{S}\mathbf{T}, u, q)$$

provided  $h = \inf u_k > 0$ .

*Proof.* For a given  $\varepsilon > 0$ , we first choose  $\varepsilon_0 > 0$  such that  $\sup_n (\sum_{k=1}^n a_{nk}) \max\{\varepsilon_0^h, \varepsilon_0^H\} < \varepsilon$ . Using the continuity of  $\mathbf{M}$ , choose  $0 < \delta < 1$  such that  $0 < \delta < t$  implies that  $S_k(t) < \varepsilon_0$  for all  $k \in \mathbb{N}$ . Let  $x = (x_k) \in \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{T}, u, q)$ . For some  $\rho > 0$  we denote

$$A_5 = \left\{ n \in \mathbb{N} : \sum_{k=1}^n a_{nk} \left[ T_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} \geq \delta^H \right\} \in I, m \in \mathbb{N}.$$

If  $n \notin A_5$ , then we have

$$\begin{aligned} & \sum_{k=1}^n a_{nk} \left[ T_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} < \delta^H \\ \text{i.e. } & \left[ T_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) \right]^{u_k} < \delta^H \text{ for all } k, m \in \mathbb{N} \\ \text{i.e. } & T_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)))}{\rho} \right) < \delta \text{ for all } k, m \in \mathbb{N} \end{aligned}$$

$$i.e. S_k \left( T_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho} \right) \right) < \varepsilon_0 \text{ for all } k, m \in \mathbb{N}.$$

Consequently, we get

$$\sum_{k=1}^n a_{nk} \left[ S_k \left( T_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho} \right) \right) \right]^{u_k} < \sup_n \left( \sum_{k=1}^n a_{nk} \right) \max\{\varepsilon_0^h, \varepsilon_0^H\} < \varepsilon$$

,  $m \in \mathbb{N}$ . i.e.

$$\sum_{k=1}^n a_{nk} \left[ S_k \left( T_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho} \right) \right) \right]^{u_k} < \varepsilon, m \in \mathbb{N}.$$

This shows that

$$\left\{ n \in \mathbb{N} : \sum_{k=1}^n a_{nk} \left[ S_k \left( T_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p(x)) \right)}{\rho} \right) \right) \right]^{u_k} \geq \varepsilon \right\} \subset A_5 \in I.$$

This completes the proof.  $\square$

**Theorem 3.5.** *The inclusions  $Z(A, B_{(i)}^{p-1}, \mathbf{M}, u, q) \subset Z(A, B_{(i)}^p, \mathbf{M}, u, q)$ , are strict for  $p \geq 1$ . In general  $Z(A, B_{(i)}^j, \mathbf{M}, u, q) \subset Z(A, B_{(i)}^p, \mathbf{M}, u, q)$ , for  $j = 0, 1, 2, \dots, p-1$  and the inclusions are strict, where  $Z = \widehat{w}_0^I, \widehat{w}^I, \widehat{w}_\infty^I$ .*

*Proof.* We shall give the proof for  $\widehat{w}_0^I(A, B_{(i)}^{p-1}, \mathbf{M}, u, q)$  only. The others can be proved by similar arguments. Let  $x = (x_k)$  be any element in the space  $\widehat{w}_0^I(A, B_{(i)}^{p-1}, \mathbf{M}, u, q)$ . Let  $\varepsilon > 0$  be given. Then there exists  $\delta > 0$  such that the set

$$\left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_k) \right)}{\rho} \right) \right]^{p_k} \geq \varepsilon \right\} \in I.$$

Since  $\mathbf{M}$  is non-decreasing and convex, it follows that

$$\begin{aligned} & \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p x_k) \right)}{2\rho} \right) \right]^{p_k} \\ &= \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_{k+1} - B_{(i)}^{p-1} x_k) \right)}{2\rho} \right) \right]^{p_k} \\ &\leq D \sum_{k=1}^{\infty} \left[ \frac{1}{2} M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_{k+1}) \right)}{\rho} \right) \right]^{p_k} \end{aligned}$$

$$\begin{aligned}
& +D \sum_{k=1}^{\infty} a_{nk} \left[ \frac{1}{2} M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_k) \right)}{\rho} \right) \right]^{p_k} \\
& \leq DH \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_{k+1}) \right)}{\rho} \right) \right]^{p_k} \\
& +DH \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_k) \right)}{\rho} \right) \right]^{p_k},
\end{aligned}$$

where  $H = \max\{1, (\frac{1}{2})^G\}$ . Thus we have

$$\begin{aligned}
& \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p x_k) \right)}{2\rho} \right) \right]^{p_k} \geq \varepsilon \right\} \\
& \subseteq \left\{ n \in \mathbb{N} : DH \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_{k+1}) \right)}{\rho} \right) \right]^{p_k} \geq \frac{\varepsilon}{2} \right\} \\
& \cup \left\{ n \in \mathbb{N} : DH \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^{p-1} x_k) \right)}{\rho} \right) \right]^{p_k} \geq \frac{\varepsilon}{2} \right\} \quad (3.2)
\end{aligned}$$

Since both the sets in the right side of (3.2) belongs to  $I$ , we get

$$\left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q \left( t_{m,k}(B_{(i)}^p x_k) \right)}{2\rho} \right) \right]^{p_k} \geq \varepsilon \right\} \in I.$$

□

If follow from the following example that the inclusion is strict.

**Example 3.1.** Let  $A = (C, 1)$ ,  $M_k(x) = x$ , for all  $x \in [0, \infty)$ ,  $u_k = 1$  for all  $k \in \mathbb{N}$  and  $r = 1$ ,  $s = -1$ . Consider a sequence  $x = (x_k) = (k^p)$ . Then  $x = (x_k)$  belongs to  $w_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  but does not belong to  $w_0^I(A, B_{(i)}^{p-1}, M, u, q)$ , because  $B_{(i)}^p x_k = 0$  and  $B_{(i)}^{p-1} x_k = (-1)^{p-1}(p-1)!$ .

**Theorem 3.6.** (a) Let  $0 < \inf u_k \leq u_k \leq 1$ , then  $\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q) \subset \widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, q)$ ;  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q) \subset \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, q)$ .

(b) If  $1 < u_k \leq \sup u_k < \infty$ , then  $\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, q) \subset \widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ ;  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, q) \subset \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ .

*Proof.* (a) Let  $x = (x_k) \in \widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . Since  $0 < \inf u_k \leq u_k \leq 1$ , we have

$$\sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right] \leq \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right]^{p_k}$$

and therefore

$$\begin{aligned} & \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right] \geq \varepsilon \right\} \\ & \subseteq \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right]^{p_k} \geq \varepsilon \right\} \in I. \end{aligned}$$

(b) Let  $1 < u_k \leq \sup u_k < \infty$  and let  $x = (x_k) \in \widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, q)$ . Then for each  $0 < \varepsilon < 1$  there exists a positive integer  $N$  such that

$$\sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right] \leq \varepsilon < 1$$

for all  $n \geq N$ . This implies that

$$\sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right]^{p_k} \leq \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right].$$

Thus we have

$$\begin{aligned} & \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right]^{p_k} \geq \varepsilon \right\} \\ & \subseteq \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k) - \ell)}{\rho} \right) \right] \geq \varepsilon \right\} \in I. \end{aligned}$$

This completes the proof.  $\square$

**Corollary 3.7.** *Let  $A = (C, 1)$  Cesáro matrix and let  $M$  be an Orlicz function.*

- (a) *If  $0 < \inf u_k \leq u_k \leq 1$ , then*
- (i)  $\widehat{w}^I(B_{(i)}^p, \mathbf{M}, u, q) \subset \widehat{w}^I(B_{(i)}^p, \mathbf{M}, q)$ ;
  - (ii)  $\widehat{w}_0^I(B_{(i)}^p, \mathbf{M}, u, q) \subset \widehat{w}_0^I(B_{(i)}^p, \mathbf{M}, q)$ .
- (b) *If  $1 < u_k \leq \sup u_k < \infty$ , then*
- (i)  $\widehat{w}^I(B_{(i)}^p, \mathbf{M}, q) \subset \widehat{w}^I(B_{(i)}^p, \mathbf{M}, u, q)$ ;
  - (ii)  $\widehat{w}_0^I(B_{(i)}^p, \mathbf{M}, q) \subset \widehat{w}_0^I(B_{(i)}^p, \mathbf{M}, u, q)$ .

**Theorem 3.8.** *Let  $0 < u_k \leq v_k$  for all  $k \in \mathbb{N}$  and  $\left(\frac{v_k}{u_k}\right)$  is bounded, then  $\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, v, q) \subseteq \widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ .*

*Proof.* The proof of the theorem is straightforward, so it is omitted.  $\square$

**Theorem 3.9.** *If  $\lim_k u_k > 0$  and  $x = (x_k) \rightarrow x_0(\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q))$ , then  $x_0$  is unique.*

*Proof.* Let  $\lim_k u_k = u_0 > 0$ . Suppose that  $(x_k) \rightarrow x_0(\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q))$  and  $(x_k) \rightarrow y_0(\widehat{w}^I(A, B_{(i)}^p, \mathbf{M}, u, q))$ .

Then there exist  $\rho_1, \rho_2 > 0$  such that for all  $m \in \mathbb{N}$

$$B_1 = \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - x_0)}{\rho_1} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \in I \quad (3.3)$$

and

$$B_2 = \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - y_0)}{\rho_1} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \in I. \quad (3.4)$$

Let  $\rho = \max\{2\rho_1, 2\rho_2\}$ . Then we have

$$\sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(x_0 - y_0)}{\rho} \right) \right]^{u_k} \leq D \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - x_0)}{\rho_1} \right) \right]^{u_k} + D \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - y_0)}{\rho_1} \right) \right]^{u_k}.$$

Thus from (3.3) and (3.4) we have for all  $m \in \mathbb{N}$

$$\begin{aligned} & \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(x_0 - y_0)}{\rho} \right) \right]^{u_k} \geq \varepsilon \right\} \\ & \subseteq \left\{ n \in \mathbb{N} : D \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - x_0)}{\rho_1} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \\ & \cup \left\{ n \in \mathbb{N} : D \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{m,k}(B_{(i)}^p(x)) - y_0)}{\rho_1} \right) \right]^{u_k} \geq \frac{\varepsilon}{2} \right\} \subseteq B_1 \cup B_2 \in I. \end{aligned}$$

Also we have

$$\left[ M_k \left( \frac{q(x_0 - y_0)}{\rho} \right) \right]^{u_k} \rightarrow \left[ M_k \left( \frac{q(x_0 - y_0)}{\rho} \right) \right]^{u_0} \text{ as } k \rightarrow \infty.$$



Therefore we have

$$\left[ M_k \left( \frac{q(x_0 - y_0)}{\rho} \right) \right]^{u_k} \rightarrow \left[ M_k \left( \frac{q(x_0 - y_0)}{\rho} \right) \right]^{u_0} = 0.$$

Hence  $x_0 = y_0$ .  $\square$

**Theorem 3.10.** *The sequence spaces  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  and  $\widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  are normal as well as monotone.*

*Proof.* We give the proof for only  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . Let  $x = (x_k) \in \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  and  $\alpha = (\alpha_k)$  be a sequence of scalars such that  $|\alpha_k| \leq 1$  for all  $k \in \mathbb{N}$ . Then for given  $\varepsilon > 0$ , for all  $m \in \mathbb{N}$  we have

$$\begin{aligned} & \left\{ n \in \mathbb{N} : \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p(\alpha_k x_k)))}{\rho} \right) \right]^{u_k} \geq \varepsilon \right\} \\ & \subseteq \left\{ n \in \mathbb{N} : E \sum_{k=1}^{\infty} a_{nk} \left[ M_k \left( \frac{q(t_{mk}(B_{(i)}^p x_k))}{\rho} \right) \right]^{u_k} \geq \varepsilon \right\} \in I, \end{aligned}$$

where  $E = \max\{1, |\alpha_k|^G\}$ .

Hence  $(\alpha_k x_k) \in \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . Thus the space  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  is normal. Also from Lemma 1.1, it follows that  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  is monotone.  $\square$

**Theorem 3.11.** *Let  $\mathbf{M} = (M_k)$  be a Musielak-Orlicz function. Then the following statements are equivalent:*

- (i)  $\widehat{w}_\infty^I(A, B_{(i)}^p, u, q) \subseteq \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$
- (ii)  $\widehat{w}_0^I(A, B_{(i)}^p, u, q) \subseteq \widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q)$
- (iii)  $\sup_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{t}{\rho} \right) \right]^{u_k} < \infty$  ( $t, \rho > 0$ ).

*Proof.* (i) $\Rightarrow$ (ii) is obvious, because  $\widehat{w}_0^I(A, B_{(i)}^p, u, q) \subseteq \widehat{w}_\infty^I(A, B_{(i)}^p, u, q)$ .

(ii) $\Rightarrow$ (iii). Suppose  $\widehat{w}_0^I(A, B_{(i)}^p, u, q) \subseteq \widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . We assume that (iii) is not satisfied. Then for some  $t, \rho > 0$

$$\sup_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{t}{\rho} \right) \right]^{u_k} = \infty,$$

and therefore there exists a sequence  $(n_j)$  of positive integers such that

$$\sum_{k=1}^{n_j} a_{n_j k} \left[ M_k \left( \frac{j^{-1}}{\rho} \right) \right]^{u_k} > j, j = 1, 2, 3, \dots \quad (3.5)$$

Define a sequence  $x = (x_k)$  by

$$B_{(i)}^p x_k = \begin{cases} \frac{1}{j}, & \text{if } 1 \leq k \leq n_j, j = 1, 2, 3, \dots; \\ 0, & \text{if } k > n_j \end{cases}$$

Then  $x = (x_k) \in \widehat{w}_0^I(A, B_{(i)}^p, u, q)$  but by equation (3.5) we have  $x = (x_k) \notin \widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  which contradicts (ii). Hence (iii) must hold.

(iii) $\Rightarrow$ (i) Suppose (iii) is satisfied and  $x \in \widehat{w}_\infty^I(A, B_{(i)}^p, u, q)$ . Suppose that  $x \notin \widehat{w}_\infty^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . Then

$$\sup_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{q \left( t_{mk}(B_{(i)}^p x_k) \right)}{\rho} \right) \right]^{u_k} = \infty, \text{ for all } m \in \mathbb{N}. \quad (3.6)$$

Put  $t = q \left( t_{mk}(B_{(i)}^p x_k) \right)$  for all  $k, m \in \mathbb{N}$ . Then by the equation (3.6) we have

$$\sup_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{t}{\rho} \right) \right]^{u_k} = \infty$$

which contradicts (iii). Hence (i) must hold.  $\square$

**Theorem 3.12.** *Let  $\mathbf{M} = (M_k)$  be a Musielak-Orlicz function. Let  $1 \leq u_k \leq \sup_k u_k < \infty$ . Then the following statements are equivalent:*

- (i)  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, q) \subseteq \widehat{w}_0^I(A, B_{(i)}^p, u, q)$
- (ii)  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q) \subseteq \widehat{w}_\infty^I(A, B_{(i)}^p, u, q)$
- (iii)  $\inf_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{t}{\rho} \right) \right]^{u_k} > 0$  ( $t, \rho > 0$ ).

*Proof.* (i) $\Rightarrow$ (ii) is obvious.

(ii) $\Rightarrow$ (iii). Suppose  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q) \subseteq \widehat{w}_\infty^I(A, B_{(i)}^p, u, q)$ . We assume that (iii) does not hold. Then for some  $t, \rho > 0$

$$\inf_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{t}{\rho} \right) \right]^{u_k} = 0.$$

We can choose an index sequence  $(n_j)$  of positive integers such that

$$\sum_{k=1}^{n_j} a_{n_j k} \left[ M_k \left( \frac{i}{\rho} \right) \right]^{u_k} > \frac{1}{j}, j = 1, 2, 3, \dots \quad (3.7)$$

Define a sequence  $x = (x_k)$  by

$$B_{(i)}^p x_k = \begin{cases} j, & \text{if } 1 \leq k \leq n_j, j = 1, 2, 3, \dots; \\ 0, & \text{if } k > n_j \end{cases}$$

Then by equation (3.7) we have  $x = (x_k) \in \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$  but  $x = (x_k) \notin \widehat{w}_\infty^I(A, B_{(i)}^p, u, q)$  which contradicts (ii). Hence (iii) must hold.

(iii) $\Rightarrow$ (i) Let (iii) hold and  $x \in \widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, u, q)$ . Then for every  $\varepsilon > 0$ , for all  $m \in \mathbb{N}$  we have

$$\left\{ n \in \mathbb{N} : \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{q \left( t_{mk}(B_{(i)}^p x_k) \right)}{\rho} \right) \right]^{u_k} \geq \varepsilon \right\} \in I. \quad (3.8)$$

Suppose that  $x \notin \widehat{w}_0^I(A, B_{(i)}^p, u, q)$ . Then for some integer  $\varepsilon_0 > 0$  for all  $m \in \mathbb{N}$  we have

$$\left\{ n \in \mathbb{N} : \sum_{k=1}^n a_{nk} \left[ q \left( t_{mk}(B_{(i)}^p x_k) \right) \right]^{u_k} \geq \varepsilon_0 \right\} \notin I.$$

Therefore we have

$$\left[ M_k \left( \frac{\varepsilon_0}{\rho} \right) \right]^{u_k} \leq \left[ M_k \left( \frac{q \left( t_{mk}(B_{(i)}^p x_k) \right)}{\rho} \right) \right]^{u_k}$$

and consequently by the relation (3.8) we have

$$\inf_n \sum_{k=1}^n a_{nk} \left[ M_k \left( \frac{\varepsilon_0}{\rho} \right) \right]^{u_k} = 0$$

which contradicts (iii). Hence  $\widehat{w}_0^I(A, B_{(i)}^p, \mathbf{M}, q) \subseteq \widehat{w}_0^I(A, B_{(i)}^p, u, q)$ .  $\square$

#### ACKNOWLEDGMENTS

The author would like to thank the referees for a careful reading and several constructive comments that have improved the presentation of the results.

#### REFERENCES

1. A. A. Arefijamaal, G. Sadeghi, Frames in 2-inner Product Spaces, *Iranian Jour. Math. Sci. Inform.*, **8** (2), (2013), 123-130.
2. C. Aydin, F. Basar, Some new difference sequence spaces, *Appl. Math. Comput.*, **157**(3), (2004), 677-693.
3. F. Başar, B. Altay, On the space of sequences of  $p$ -bounded variation and related matrix mappings, *Ukrainian Math. Jour.*, **55**(1), (2003), 136-147.
4. M. Başarir, M. Kayikçi, On the generalized  $B^m$ -Riesz difference sequence spaces and  $\beta$ -property, *Jour. Ineq. Appl.*, **2009**, (2009), Article ID 385029, 18 pages.
5. M. Başarir, Ş. Konca, E. E. Kara, Some generalized difference statistically convergent sequence spaces in 2-normed space, *Jour. Ineq. Appl.*, **2013**, 357
6. H. Çakalli, B. Hazarika, Ideal quasi-Cauchy sequences, *Jour. Ineq. Appl.*, (2012), pages 11, doi:10.1186/1029-242X-2012-234
7. R. Çolak, M. Et, On Some Generalized Difference Sequence Spaces and Related Matrix Transformation, *Hokkaido Math. J.*, **26** (3), (1997), 483-492.

8. R. Colak, H. Altinok, M. Et, Generalized difference sequences of fuzzy numbers, *Chaos, Solitons, Fractals*, **40**, (2009), 1106-1117.
9. Y. A. Cui, *On some geometric properties in Musielak-Orlicz sequence spaces*, Lecture Notes in Pure and Applied Mathematics (Marcel Dekker. Inc., New York and Basel, 2003), pp 213.
10. H. Dutta, Some statistically convergent difference sequence spaces defined over real 2-normed linear space, *Appl. Sci.*, **12**, (2010), 37-47.
11. A. Esi, Strongly almost summable sequence spaces in 2-normed spaces defined by ideal convergence and an Orlicz function, *Stud. Univ. Babeş-Bolyai Math.*, **57(1)**(2012), 75–82
12. M. Et, R. Çolak, On generalized difference sequence spaces, *Soochow J. Math.*, **21** (4), (1995), 377-386.
13. M. Et, Y. Altin, B. Choudhary, B. C. Tripathy, On some classes of sequences defined by sequences of Orlicz functions, *Math. Ineq. Appl.*, **9** (2),(2006) 335-342.
14. H. Fast, Sur la convergence statistique, *Colloq. Math.*, **2**, (1951), 241-244.
15. J. A. Fridy, On statistical convergence, *Analysis*, **5**, (1985), 301-313.
16. M. Gungor, M. Et,  $\Delta^m$ -strongly almost summable sequences defined by Orlicz functions, *Indian J. Pure Appl. Math.*, **34** (8), (2003), 1141-1151.
17. M. Gurdal, On ideal convergent sequences in 2-normed spaces, *Thai J. Math.*, **4** (1), (2006), 85-91.
18. B. Hazarika, Some lacunary difference sequence spaces defined by Musielak-Orlicz functions, *Asia-European Jour. Math.*, **4** (4), (2011), 613-626, DOI: 10.1142/S1793557111000484.
19. B. Hazarika, On Paranormed Ideal Convergent Generalized Difference Strongly Summable Sequence Spaces Defined over  $n$ -Normed Spaces, *ISRN Math. Anal.*, **17**, (2011), doi:10.5402/2011/317423.
20. B. Hazarika, E. Savas, Some  $I$ -convergent lambda-summable difference sequence spaces of fuzzy real numbers defined by sequence of Orlicz functions, *Math. Compu. Modell.*, **54** (11-12), (2011), 2986-2998.
21. B. Hazarika, On generalized difference ideal convergence in random 2-normed spaces, *Filomat*, **26** (6), (2012), 1265-1274.
22. B. Hazarika, On fuzzy real valued generalized difference  $I$ -convergent sequence spaces defined by Musielak-Orlicz function, *Jour. Intell. Fuzzy Systems*, **25** (1), (2013), 9-15, DOI: 10.3233/IFS-2012-0609.
23. V. A. Khan, Q. M. D. Lohani, Some new difference sequence spaces defined by Musielak-Orlicz function, *Thai J. Math.*, **6** (1), (2008), 215-223.
24. H. Kizmaz, On certain sequence spaces, *Canad. Math. Bull.*, **24**(2), (1981), 169-176.
25. P. Kostyrko, T. Šalát, W. Wilczyński,  $I$ -convergence, *Real Analysis Exchange*, **26** (2), (2000-2001), 669-686.
26. M. A. Krasnoselski, Y. B. Rutitsky, *Convex Functions and Orlicz Functions*, P. Noordhoff, Groningen, Netherlands, 1961.
27. K Lindberg, On subspaces of Orlicz sequence spaces, *Studia Math.*, **45**, (1973), 119-146.
28. J. Lindenstrauss, L. Tzafriri, On Orlicz sequence spaces, *Israel J. Math.*, **10**, (1971), 379-390.
29. G. G. Lorentz, A contribution to the theory of divergent series, *Acta Math.*, **80**, (1948), 167-190.
30. I. J. Maddox, Spaces of strongly summable sequence, *The Quarterly Jour. Math.*, **18**, (1967), 345-355.
31. M. Mursaleen, M. A. Khan, Qamaruddin, Difference sequence spaces defined by Orlicz functions, *Demonstratio. Math.*, **32**, (1999), 145-150.

32. A. Niknam, S. S. Gamchi , M. Janfada, Some Results on TVS-cone Normed Spaces and Algebraic Cone Metric Spaces, *Iranian Jour. Math. Sci. Inform.*, **9** (1), (2014), 71-80.
33. S. D. Parashar, B. Choudhary, Sequence spaces defined by Orlicz functions, *Indian J. Pure Appl. Math.*, **25**(4), (1994), 419-428.
34. W. H. Ruckle, FK-spaces in which the sequence of coordinate vectors is bounded, *Canad. J. Math.*, **25**, (1973), 973-978.
35. I. J. Schoenberg, The integrability of certain functions and related summability methods, *Amer. Math. Monthly*, **66**, (1959), 361-375.
36. B. C. Tripathy, A. Esi, A new type of difference sequence spaces, *Int. Jour. Sci. Tech.*, **1**(1), (2006), 11-14.
37. B.C. Tripathy, A. Esi, B. K. Tripathy, On a new type of generalized difference Cesaro sequence spaces, *Soochow J. Math.*, **31**, (2005), 333-340.
38. B. C. Tripathy, B. Hazarika, Paranorm  $I$ -convergent sequence spaces, *Math. Slovaca*, **59** (4), (2009), 485-494.
39. B. C. Tripathy, B. Hazarika, Some  $I$ -convergent sequence spaces defined by Orlicz functions, *Acta Math. Appl. Sinica*, **27** (1), (2011), 149-154.
40. B. C. Tripathy, B. Hazarika,  $I$ -monotonic and  $I$ -convergent sequences, *Kyungpook Math. Jour.*, **51**(2), (2011), 233-239 DOI 10.5666/KMJ.2011.51.2.233.
41. B. C. Tripathy, M. Et, Y. Altin, Generalized difference sequence spaces defined by Orlicz function in a locally convex space, *J. Anal. Appl.*, **3** (1), (2003), 175-192.