

Sufficient Inequalities for Univalent Functions

Rahim Kargar^{a,*}, Ali Ebadian^b and Janusz Sokół^c

^aYoung Researchers and Elite Club, Urmia Branch, Islamic Azad University,
Urmia, Iran.

^bDepartment of Mathematics, Payame Noor University, Tehran, Iran.

^cUniversity of Rzeszów, Faculty of Mathematics and Natural Sciences, ul.
Prof. Pigoń 1, 35-310 Rzeszów, Poland

E-mail: rkargar1983@gmail.com

E-mail: ebadian.ali@gmail.com

E-mail: jsokol@ur.edu.pl

ABSTRACT. In this work, applying a lemma due to Nunokawa et al. (J. Ineq. Appl. 2015. 1(2015) 7), we obtain some sufficient inequalities for some certain subclasses of univalent functions.

Keywords: Analytic, Univalent, Starlike functions, Convex functions.

2000 Mathematics subject classification: 30C45.

1. INTRODUCTION

Let \mathcal{A} denote the class of functions $f(z)$ of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1.1)$$

which are analytic in the open unit disk $\Delta = \{z \in \mathbb{C} : |z| < 1\}$. The subclass of \mathcal{A} consisting of all univalent functions $f(z)$ in Δ is denoted by \mathcal{S} . A function $f \in \mathcal{S}$ is called starlike (with respect to 0), denoted by $f \in \mathcal{S}^*$, if $tw \in f(\Delta)$ whenever $w \in f(\Delta)$ and $t \in [0, 1]$. A function $f \in \mathcal{S}$ that maps Δ onto a convex

*Corresponding Author

domain, denoted by $f \in \mathcal{K}$, is called a convex function. A function $f(z)$ in \mathcal{A} is said to be starlike of order $0 \leq \gamma < 1$ if it satisfies

$$\Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \gamma \quad z \in \Delta.$$

We denote by $\mathcal{S}^*(\gamma)$ the subclass of \mathcal{A} consisting of all starlike functions of order γ in Δ . Put $\mathcal{S}^*(0) \equiv \mathcal{S}^*$. A function $f(z)$ in \mathcal{A} is said to be convex of order $0 \leq \gamma < 1$ if it satisfies

$$\Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \gamma \quad z \in \Delta.$$

We denote by $\mathcal{K}(\gamma)$ the subclass of \mathcal{A} consisting of all convex functions of order γ in Δ . Let \mathcal{R}_γ denote the subclass of \mathcal{A} consisting of functions $f(z)$ which satisfy

$$\Re\{f'(z)\} > \gamma \quad z \in \Delta,$$

for some $0 \leq \gamma < 1$. The class \mathcal{R}_γ is considered in [2]. We remark that the classes $\mathcal{S}^*(\gamma)$, $\mathcal{K}(\gamma)$ and \mathcal{R}_γ ($0 \leq \gamma < 1$) are subclasses of univalent functions.

Nunokawa, Cho, Kwon and Sokół [4], obtained the following result.

Lemma 1.1. *Let $B(z)$ and $C(z)$ be analytic in Δ with*

$$|\Im\{C(z)\}| < \Re\{B(z)\}.$$

If $p(z)$ is analytic in Δ with $p(0) = 1$, and if

$$|\arg\{B(z)zp'(z) + C(z)p(z)\}| < \pi/2 + t(z), \tag{1.2}$$

where

$$t(z) = \begin{cases} \arg\{C(z) + iB(z)\} & \text{when } \arg\{C(z) + iB(z)\} \in [0, \pi/2], \\ \arg\{C(z) + iB(z)\} - \pi/2 & \text{when } \arg\{C(z) + iB(z)\} \in (\pi/2, \pi], \end{cases}$$

then we have

$$\Re\{p(z)\} > 0 \quad z \in \Delta.$$

In this paper, applying the Lemma 1.1, we obtain sufficient conditions for some certain subclasses of univalent and analytic functions. The study of analytic functions and such results, also for other subclass were investigated in [1, 3, 7].

2. MAIN RESULTS

One of our main results is contained in the following.

Theorem 2.1. *Assume that $f \in \mathcal{A}$, $f(z)/z$ is bounded and not vanishing in the unit disc. If*

$$\left| \arg \left\{ \sqrt[n]{\left(\frac{z}{f(z)}\right)^{n-1}} f'(z) - \gamma \right\} \right| < \frac{\pi}{2} + \arctan n, \tag{2.1}$$

then

$$\Re \left\{ \sqrt[n]{\frac{f(z)}{z}} \right\} > \gamma \quad z \in \Delta, \quad (2.2)$$

where $\sqrt{1} = 1$, $\gamma < 1$ and $n = 1, 2, 3, \dots$

Proof. Let $f(z) \neq 0$ for $z \neq 0$ and let $p(z)$ be defined by

$$p(z) = \frac{1}{1-\gamma} \left(\sqrt[n]{\frac{f(z)}{z}} - \gamma \right) \quad z \in \Delta,$$

where $\gamma < 1$. Then $p(z)$ is analytic in Δ , $p(0) = 1$ and

$$(1-\gamma)p(z) + n(1-\gamma)zp'(z) = \sqrt[n]{\left(\frac{z}{f(z)}\right)^{n-1}} f'(z) - \gamma.$$

If we put $B(z) = n(1-\gamma)$ and $C(z) = 1-\gamma$, from (2.1) and applying Lemma 1.1, we obtain (2.2) immediately. \square

Putting $n = 1$, in Theorem 2.1, we have:

Corollary 2.2. (see [5]) *Let $f \in \mathcal{A}$, $\gamma < 1$ and $f(z)/z$ is bounded and not vanishing in the unit disc. If*

$$|\arg \{f'(z) - \gamma\}| < \frac{3\pi}{4},$$

then

$$\Re \left\{ \frac{f(z)}{z} \right\} > \gamma \quad z \in \Delta.$$

If we take $n = 2$, in Theorem 2.1, we have:

Corollary 2.3. *Let $f \in \mathcal{A}$, $\gamma < 1$ and $f(z)/z$ is bounded and not vanishing in the unit disc. If*

$$\left| \arg \left\{ \sqrt{\frac{z}{f(z)}} f'(z) - \gamma \right\} \right| < \frac{\pi}{2} + \arctan 2 \approx 153.4349\dots,$$

then

$$\Re \left\{ \sqrt{\frac{f(z)}{z}} \right\} > \gamma \quad z \in \Delta.$$

In [6], Obradović, Ponnusamy and Tuneski introduced the class $\mathcal{P}(1/2)$ as follows:

$$\mathcal{P}(1/2) := \{f \in \mathcal{A} : \Re(f(z)/z) > 1/2 \text{ for } z \in \Delta\}.$$

We remark that $\mathcal{K}(0) \subset \mathcal{P}(1/2)$ (see [6]).

Setting $n = 1$ and $\gamma = 1/2$, in Theorem 2.1, we have:

Corollary 2.4. Let $f \in \mathcal{A}$, $f(z)/z$ is bounded and not vanishing in the unit disc. If

$$\left| \arg \left\{ f'(z) - \frac{1}{z} \right\} \right| < \frac{3\pi}{4} \quad z \in \Delta,$$

then $f \in \mathcal{P}(1/2)$.

Next, we derive the following.

Theorem 2.5. Suppose that $f \in \mathcal{A}$, $f(z)/z$ is bounded and not vanishing in the unit disc. If

$$\left| \arg \left\{ \left[\frac{z}{f(z)} \right]^{\frac{n-1}{n(1-\gamma)}} \left(\frac{f(z)}{z^\gamma} \right)^{\frac{1}{1-\gamma}} \left(\frac{f'(z)}{f(z)} - \frac{\gamma}{z} \right) - \gamma(1-\gamma) \right\} \right| < \frac{\pi}{2} + \arctan n, \quad (2.3)$$

then

$$\Re \left\{ \left(\frac{f(z)}{z} \right)^{\frac{1}{n(1-\gamma)}} \right\} > \gamma \quad z \in \Delta,$$

where $\gamma < 1$ and $n = 1, 2, 3, \dots$

Proof. Let us define the function $p(z)$ by

$$p(z) = \left(\frac{f(z)}{z^\gamma} \right)^{\frac{1}{1-\gamma}} = z + \dots \quad z \in \Delta, \quad (2.4)$$

for $f \in \mathcal{A}$. Then $p(z) \in \mathcal{A}$ and

$$p'(z) = \frac{1}{1-\gamma} \left(\frac{f(z)}{z^\gamma} \right)^{\frac{1}{1-\gamma}} \left(\frac{f'(z)}{f(z)} - \frac{\gamma}{z} \right) \quad z \in \Delta, \quad (2.5)$$

which gives

$$\begin{aligned} & \left(\frac{z}{p(z)} \right)^{\frac{n-1}{n}} p'(z) - \gamma \\ &= \frac{1}{1-\gamma} \left\{ \left[\frac{z}{f(z)} \right]^{\frac{n-1}{n(1-\gamma)}} \left(\frac{f(z)}{z^\gamma} \right)^{\frac{1}{1-\gamma}} \left(\frac{f'(z)}{f(z)} - \frac{\gamma}{z} \right) - \gamma(1-\gamma) \right\} \quad z \in \Delta. \end{aligned} \quad (2.6)$$

Applying Theorem 2.1, we have

$$\Re \left\{ \sqrt[n]{\frac{p(z)}{z}} \right\} > \gamma \quad z \in \Delta,$$

that is

$$\Re \left\{ \left(\frac{f(z)}{z} \right)^{\frac{1}{n(1-\gamma)}} \right\} > \gamma \quad z \in \Delta.$$

So the proof is completed. \square

If in Theorem 2.5 we take $n = 2$ and $\gamma = 1/2$, then we get:

Corollary 2.6. *Suppose that $f \in \mathcal{A}$, $f(z)/z$ is bounded and not vanishing in the unit disc. If*

$$\left| \arg \left\{ f'(z) - \frac{f(z)}{2z} - \frac{1}{4} \right\} \right| < \frac{\pi}{2} + \arctan 2 \approx 153.4349\dots \quad z \in \Delta, \quad (2.7)$$

then

$$\Re \left\{ \frac{f(z)}{z} \right\} > \frac{1}{2} \quad z \in \Delta,$$

i.e. $f \in \mathcal{P}(1/2)$.

In the following, we will provide some sufficient inequalities for the functions belonging to the classes $\mathcal{S}^*(\gamma)$, $\mathcal{K}(\gamma)$ and $\mathcal{R}(\gamma)$.

Theorem 2.7. *Assume that $f \in \mathcal{A}$ and $0 \leq \gamma < 1$. If*

$$\left| \arg \left\{ \frac{zf'(z)}{f(z)} \left(2 - \frac{zf'(z)}{f(z)} \right) + \frac{z^2 f''(z)}{f(z)} - \gamma \right\} \right| < \frac{3\pi}{4}, \quad (2.8)$$

then $f \in \mathcal{S}^*(\gamma)$.

Proof. For $0 \leq \gamma < 1$ let $p(z)$ be defined by

$$p(z) = \frac{1}{1-\gamma} \left(\frac{zf'(z)}{f(z)} - \gamma \right) \quad z \in \Delta.$$

Then $p(z)$ is analytic in Δ , $p(0) = 1$ and

$$(1-\gamma)p(z) + (1-\gamma)zp'(z) = \frac{zf'(z)}{f(z)} \left(2 - \frac{zf'(z)}{f(z)} \right) + \frac{z^2 f''(z)}{f(z)} - \gamma.$$

Put $B(z) = 1 - \gamma = C(z)$. Now, from (2.8) and applying Lemma 1.1, we can obtain the result. \square

Corollary 2.8. *Let $f \in \mathcal{A}$. If*

$$\left| \arg \left\{ \frac{zf'(z)}{f(z)} \left(2 - \frac{zf'(z)}{f(z)} \right) + \frac{z^2 f''(z)}{f(z)} \right\} \right| < \frac{3\pi}{4},$$

then f is starlike function.

Theorem 2.9. *Assume that $f \in \mathcal{A}$. If*

$$\left| \arg \left\{ \frac{2zf''(z) + z^2 f'''(z)}{f'(z)} + 1 - \left(\frac{zf''(z)}{f'(z)} \right)^2 - \gamma \right\} \right| < \frac{3\pi}{4}, \quad (2.9)$$

then $f \in \mathcal{K}(\gamma)$, where $0 \leq \gamma < 1$.

Proof. It is enough that $p(z)$ to be defined as follows:

$$p(z) = \frac{1}{1-\gamma} \left(1 + \frac{zf''(z)}{f'(z)} - \gamma \right) \quad z \in \Delta.$$

Then applying Lemma 1.1, the result is obtained. \square

Corollary 2.10. *Let $f \in \mathcal{A}$. If*

$$\left| \arg \left\{ \frac{2zf''(z) + z^2f'''(z)}{f'(z)} + 1 - \left(\frac{zf''(z)}{f'(z)} \right)^2 \right\} \right| < \frac{3\pi}{4}, \quad (2.10)$$

then f is convex function.

Theorem 2.11. *Assume that $f \in \mathcal{A}$. If*

$$|\arg\{f'(z) + zf''(z) - \gamma\}| < \frac{3\pi}{4}, \quad (2.11)$$

then $f \in \mathcal{R}_\gamma$ where $0 \leq \gamma < 1$.

Proof. Suppose that $p(z)$ be defined by

$$(1 - \gamma)p(z) + \gamma = f'(z) \quad z \in \Delta.$$

Then applying Lemma 1.1, the result is obtained. \square

ACKNOWLEDGMENTS

The authors express their sincerest thanks to the referee for various useful suggestions.

REFERENCES

1. R. Aghalary, Application of the Norm Estimates for Univalence of Analytic Functions, *Iranian Journal of Mathematical Sciences and Informatics*, **9**(2), (2014), 101-108.
2. A.W. Goodman, *Univalent Functions*, Vols. 1 and 2, Mariner, Tampa, FL, 1983.
3. R. Kargar, A. Ebadian, J. Sokół, Certain Coefficient Inequalities for p -valent Functions, *Malaya J. Mat.*, **4**(1), (2016), 37-41.
4. M. Nunokawa, N.E. Cho, O.S. Kwon, J. Sokół, On Differential Subordinations in the Complex Plane, *Journal of Inequalities and Applications*, **2015**(1), (2015), 7.
5. M. Nunokawa, S. Owa, K. Kuroki, J. Sokół, Some Generalization Properties of Analytic Functions Concerned with Sakaguchi Result, *Tamkang Journal of Mathematics*, **46**(2), (2015), 143-150.
6. M. Obradović, S. Ponusamy, N. Tuneski, Radius of Univalence of Certain Combination of Univalent of Analytic functions, *Bull. Malays. Math. Sci. Soc.*, **35**(2), (2012), 325-334.
7. A. Taghavi, R. Hosseinzadeh, Uniform Boundedness Principle for Operators on Hyper-vector spaces, *Iranian Journal of Mathematical Sciences and Informatics*, **7**(2), (2012), 9-16.