

A Graphical Characterization for *SPAP*-Rings

Esmail Rostami

Department of Pure Mathematics, Faculty of Mathematics and
Computer, Shahid Bahonar University of Kerman, Kerman, Iran

E-mail: e_rostami@uk.ac.ir

ABSTRACT. Let R be a commutative ring and I an ideal of R . The zero-divisor graph of R with respect to I , denoted by $\Gamma_I(R)$, is the simple graph whose vertex set is $\{x \in R \setminus I \mid xy \in I, \text{ for some } y \in R \setminus I\}$, with two distinct vertices x and y are adjacent if and only if $xy \in I$. In this paper, we state a relation between zero-divisor graph of R with respect to an ideal and almost prime ideals of R . We then use this result to give a graphical characterization for *SPAP*-rings.

Keywords: *SPAP*-ring, Almost prime ideal, Zero-divisor graph with respect to an ideal.

2000 Mathematics subject classification: 13A25, 05C17.

1. INTRODUCTION

Throughout this paper, all rings are assumed to be commutative with identity. A graph (simple graph) G is an ordered pair of disjoint sets (V, E) such that $V = V(G)$ is the vertex set of G and $E = E(G)$ is its edge set. A graph F is called a subgraph of a graph G if $V(F) \subseteq V(G)$ and $E(F) \subseteq E(G)$. A subgraph F of G is said to be an induced subgraph of G if each edge of G having its ends in $V(F)$ is also an edge of F . A graph in which each pair of distinct vertices is joined by an edge is called complete.

There have been several studies concerning the assignment a graph to a ring, a group, a semigroup or a module, for more information see [1], [8] and [12]. The

Received 5 August 2015; Accepted 28 June 2016

©2018 Academic Center for Education, Culture and Research TMU

concept of the zero-divisor graph of a commutative ring R was first introduced by Beck [6]. The zero-divisor graph of a commutative ring R is defined to be the graph $\Gamma(R)$, whose vertices are the non-zero zero-divisors of R , and where x is adjacent to y if $xy = 0$. In [10] Redmond has generalized the notion of the zero-divisor graph. For a given ideal I of a commutative ring R , he defined the zero-divisor graph of R with respect to I , denoted by $\Gamma_I(R)$, is the simple graph whose vertex set is $\{x \in R \setminus I \mid xy \in I, \text{ for some } y \in R \setminus I\}$, with two distinct vertices x and y joined by an edge when $xy \in I$. Clearly $\Gamma_0(R) = \Gamma(R)$. Bhatwadekar and Sharma [7] defined a proper ideal I of an integral domain R to be almost prime if for $a, b \in R$, $ab \in I \setminus I^2$, then either $a \in I$ or $b \in I$. Anderson and Bataineh [3], use this definition for an arbitrary commutative ring and stated a necessary and sufficient condition for a commutative Noetherian ring under which every proper ideal of R is a product of almost prime ideals. Then Rostami and Nekooei [11], considered *SPAP*-rings and characterized the structure of *SPAP*-rings, in special cases. Also, they showed that *SPAP*-rings are *quasi-Frobenius* (a Noetherian self-injective ring), and *SPAP*-rings are an applicative class of rings in Coding Theory, for more information see [11]. In the next section, we state a relation between zero-divisor graph with respect to an ideal of R and almost prime ideals of R . Then we state the concept of the intersection graph of ideals of R , and we give a graphical characterization for *SPAP*-rings.

2. MAIN RESULTS

A proper ideal I in a ring R is called almost prime if for all $a, b \in R$, $ab \in I \setminus I^2$ either $a \in I$ or $b \in I$. Also, a proper ideal I of a ring R is called weakly prime if for all $a, b \in R$ with $0 \neq ab \in I$, either $a \in I$ or $b \in I$. Clearly, every weakly prime ideal is almost prime. The following lemma which plays an important role in this paper gives a graphical characterization for almost prime ideals.

Lemma 2.1. *Let I be a proper ideal of R . Then I is an almost prime ideal of R if and only if $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$.*

Proof. Let I be an almost prime ideal of R and $x \in V(\Gamma_I(R))$, then $x \in R \setminus I$, and there exists $y \in R \setminus I$ such that $xy \in I$. Thus $x, y \notin I^2$. Now if $xy \notin I^2$, then we have $xy \in I \setminus I^2$, this gives $x \in I$ or $y \in I$, a contradiction. Thus $xy \in I^2$, and so $x \in V(\Gamma_{I^2}(R))$. Now let $x, y \in V(\Gamma_I(R))$ be adjacent in $\Gamma_I(R)$, so $xy \in I$, if $xy \notin I^2$, then we have $xy \in I \setminus I^2$, this gives $x \in I$ or $y \in I$, a contradiction. Therefore, x and y are adjacent in $\Gamma_{I^2}(R)$. Thus $E(\Gamma_I(R)) \subseteq E(\Gamma_{I^2}(R))$. Clearly, each edge of $\Gamma_{I^2}(R)$ having its ends in $\Gamma_I(R)$ is also an edge of $\Gamma_I(R)$. Therefore, $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$. Conversely, let $\Gamma_I(R)$ be an induced subgraph of $\Gamma_{I^2}(R)$ and $ab \in I \setminus I^2$, if

$a, b \notin I$ then, a and b are adjacent in $\Gamma_I(R)$ and so a and b are adjacent in $\Gamma_{I^2}(R)$, thus $ab \in I^2$, a contradiction. Therefore, either $a \in I$ or $b \in I$. \square

The following lemma is a similar result for weakly prime ideals.

Lemma 2.2. *Let I be a proper ideal of R . Then I is a weakly prime ideal of R if and only if $\Gamma_I(R)$ is an induced subgraph of $\Gamma(R)$.*

Proof. Let I be a weakly prime ideal of R and $x \in V(\Gamma_I(R))$. Then $x \in R \setminus I$, and there exists $y \in R \setminus I$ such that $xy \in I$. If $xy \neq 0$, we have $0 \neq xy \in I$, this gives $x \in I$ or $y \in I$, a contradiction. Thus $xy = 0$, and so $x \in V(\Gamma(R))$. Now, let $x, y \in V(\Gamma_I(R))$ be adjacent in $\Gamma_I(R)$, thus $xy \in I$. Repeating the previous argument leads to $xy = 0$. Hence x, y are adjacent in $\Gamma(R)$. Clearly, each edge of $\Gamma(R)$ having its ends in $\Gamma_I(R)$ is also an edge of $\Gamma_I(R)$. Therefore $\Gamma_I(R)$ is an induced subgraph of $\Gamma(R)$. Conversely, let $\Gamma_I(R)$ be an induced subgraph of $\Gamma(R)$ and $0 \neq ab \in I$ for $a, b \in R$, if $a \notin I$ and $b \notin I$ then, a and b are adjacent in $\Gamma_I(R)$, thus a and b are adjacent in $\Gamma(R)$. This gives $ab = 0$, a contradiction. Thus either $a \in I$ or $b \in I$. \square

Lemma 2.3. *Let I be a proper ideal of R . Then I is a prime ideal of R if and only if $\Gamma_I(R) = \emptyset$.*

Proof. The proof is straightforward. \square

Now let I be a prime ideal of R . Thus $\Gamma_I(R) = \emptyset$ and so $\Gamma_I(R) = \emptyset$ is an induced subgraph of $\Gamma_{I^2}(R)$ and $\Gamma(R)$, this is a graphical verification for the fact that “prime ideals are almost prime and weakly prime”.

Definition 2.4. A local ring (R, m) is called special product of almost prime ideals ring (*SPAP-ring*), if for each $x \in m \setminus m^2$, $\langle x^2 \rangle = m^2$ and $m^3 = 0$.

SPAP-rings were first introduced in [3] by D. D. Anderson and M. Bataineh. In [3], D. D. Anderson and M. Bataineh used *SPAP-rings* to characterize Noetherian rings whose proper ideals are a product of almost prime ideals. In general, an *SPAP-ring* is not Noetherian, see [3, Example 20]. For an *SPAP-ring* (R, m) , m is the unique prime ideal of R , thus R is a Noetherian ring if and only if R is an Artinian ring if and only if m is a finitely generated ideal of R .

Before proceeding, we mention the definition of the intersection graph of ideals of a ring which helps us to give a characterization for *SPAP-rings*.

Definition 2.5. Let R be a ring, the intersection graph of ideals of R , denoted by $G(R)$, is the graph whose vertices are proper non-trivial ideals of R and two distinct vertices are adjacent if and only if the corresponding ideals of R have a non-trivial (non-zero) intersection.

Lemma 2.6. [5, Theorem 2.11.] *Let (R, m) be an Artinian local ring. Then the intersection graph of ideals of R is complete if and only if R has a unique minimal ideal.*

For more information about intersection graph of ideals of R , see [2, 5]. In the remainder of this section, we characterize Artinian local rings which $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$ for all non-minimal ideals I of R , and the intersection graph of ideal of R is complete.

Lemma 2.7. *Let (R, m) be an Artinian local ring and $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, for every non-minimal ideal I of R . Then m^2 is a minimal ideal of R or $m^2 = 0$.*

Proof. Let m^2 be a non-minimal ideal of R . Then by Lemma 2.1, m^2 is an almost prime ideal of R . We show that m^2 must be zero in this case. For this purpose, we show that $m^2 = m^3 = m^4$ and the Nakayama's Lemma gives $m^2 = 0$. If for all $x, y \in m$, $xy \in m^4$, we have $m^2 \subseteq m^4$, thus $m^2 = m^3 = m^4$. Now let there exist $x, y \in m$ such that $xy \notin m^4$, so $xy \in m^2 \setminus m^4 = m^2 \setminus (m^2)^2$, since m^2 is almost prime, only one of the following cases happens;

$x \in m^2$ and $y \notin m^2$ or $x \notin m^2$ and $y \in m^2$. Suppose $x \in m^2$ and $y \notin m^2$. Since $y^2 \in m^2$, $y \notin m^2$ and m^2 is almost prime, we must have $y^2 \in m^4$. Repeating the previous argument and $y, x+y \notin m^2$ and $y(x+y) \in m^2$ leads to $y(x+y) \in m^4$. Thus $xy + y^2 = y(x+y)$, $y^2 \in m^4$, so $xy \in m^4$, a contradiction. Thus m^2 is zero or a minimal ideal. \square

Now we mention the definition of a class of rings which are important in the rest of this paper.

Definition 2.8. A commutative ring R is called special principal ideal ring (*SPIR*), if it is a principal ideal ring with unique prime ideal and that prime ideal is nilpotent.

Mori [9] has shown that a ring has the property that every ideal is a product of prime ideals if and only if it is a finite direct product of Dedekind domains and special principal ideal rings (*SPIRs*) (For more information about special principal ideal ring see [9]). In the next lemma, we state a relation between *SPAP*-rings and *SPIR* rings.

Lemma 2.9. *Let (R, m) be an *SPIR* ring such that $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, for every non-minimal ideal I and m^2 is the unique minimal ideal of R . Then (R, m) is an *SPAP*-ring.*

Proof. Since R is an *SPIR* ring, $m = \langle x \rangle$ for some $x \in m$. Now let $0 \neq J \neq m^2$ be an ideal of R . If $J = J^2$, Nakayama's Lemma gives $J = 0$, a contradiction. So $J \neq J^2$, thus we can select $y \in J \setminus J^2 \subseteq m$ such that $J = \langle y \rangle$. Thus $y = rx \in J \setminus J^2$, for some $r \in R$. Since $J \neq m^2$, Lemma

2.1 gives J is an almost prime ideal of R and since $y = rx \in J \setminus J^2$, we have $x \in J$ or $r \in J$. If $x \in J$, then $J = m$ and if $r \in J \subseteq m$, then we have $J = \langle y \rangle = \langle rx \rangle \subseteq m^2$ and since m^2 is the unique minimal ideal of R , $J = 0$ or $J = m^2$, a contradiction. This means, the set of all ideals of R is $\{0, m^2, m, R\}$.

Now if $m = m^2$, we have $m = 0$, a contradiction. Thus $m \neq m^2$. If $a \in m \setminus m^2$, since the set of all ideals of R is $\{0, m^2, m, R\}$, we have $m = \langle a \rangle$, so $m^2 = \langle a^2 \rangle$. Now if $m^3 \neq 0$, we have $m^2 = m^3$, and Nakayama's Lemma gives $m = 0$, a contradiction. Thus $m^3 = 0$. This completes the proof. \square

D. D. Anderson and M. Bataineh in [3], by using *SPAP*-rings, characterized Noetherian rings whose proper ideals are a product of almost prime ideals. Actually, they stated the following theorem.

Theorem 2.10. [3, Theorem 22]. *Let R be a Noetherian ring. Then every proper ideal of R is a product of almost prime ideals if and only if R is a finite direct product of Dedekind domains, SPIRs, and (Noetherian) SPAP-rings.*

Proposition 2.11. *Let (R, m) be an Artinian local ring such that $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, for every non-minimal ideal I of R and the intersection graph of ideal of R is complete. If $m^2 \neq 0$ then R is an SPAP-ring.*

Proof. Since the intersection graph of ideals of R is complete, by Lemma 2.6, R has a unique minimal ideal. Since $m^2 \neq 0$, Lemma 2.7 gives, m^2 is the unique minimal ideal of R . Now let I be an arbitrary proper ideal of R , if I is a non-minimal ideal of R , then $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, so I is an almost prime ideal of R , by Lemma 2.1, and if I is a minimal ideal of R , then $I = m^2$. Therefore, in all cases I is finite product of almost prime ideals (note that m is prime and so is almost prime), thus by Theorem 2.10, R is a finite direct product of Dedekind domains, SPIR rings, and SPAP-rings. Since R is a local ring, this direct product must have a single ring.

Let R be a Dedekind domain. Since m^2 is a minimal ideal of R , we have $m^3 = 0$ or $m^2 = m^3$, in both cases, we have $m^2 = 0$. Thus R is not a Dedekind domain and Lemma 2.9, completes the proof. \square

An R -module M is said to be a multiplication R -module if for each submodule N of M there exists an ideal I of R such that $N = IM$. Clearly, every cyclic module is multiplication module, see [4] for more information. After stating the main result, we require the following three lemmas.

Lemma 2.12. *Let (R, m) be an SPAP-ring. If $m^2 \neq 0$, then m^2 is a minimal ideal of R .*

Proof. If $m = m^2$, then $m^2 = m^3 = 0$, a contradiction. Therefore $m \neq m^2$, thus there exists $y \in m \setminus m^2$. So $m^2 = \langle y^2 \rangle$. Therefore, m^2 is a cyclic R -module and so it is a multiplication R -module. Now if J is a submodule (ideal of R) of m^2 , there exists an ideal K of R , such that $J = Km^2$. If $K = R$, then $J = m^2$ and if $K \neq R$ then $J = Km^2 \subseteq m^3 = 0$, hence $J = 0$. Therefore m^2 is a minimal ideal of R . \square

Lemma 2.13. *Let (R, m) be an SPAP-ring. If $m^2 \neq 0$ and I is a proper ideal of R , then $I = 0$ or $I = m^2$ or $I^2 = m^2$.*

Proof. Since $m^2 \neq 0$, by Lemma 2.12, m^2 is a minimal ideal of R . Now let I be a proper ideal of (R, m) . If $I \subseteq m^2$, then $I = 0$ or $I = m^2$. If $I \not\subseteq m^2$, then there exists $y \in I \setminus m^2$. So $m^2 = \langle y^2 \rangle$, hence $m^2 = \langle y^2 \rangle \subseteq I^2 \subseteq m^2$. Thus $I^2 = m^2$. \square

By combining the above two lemmas, we have the following lemma.

Lemma 2.14. *Let (R, m) be an SPAP-ring. If $m^2 \neq 0$, then m^2 is the unique minimal ideal of R .*

Now we can state the main result of this paper.

Theorem 2.15. *Let (R, m) be an Artinian local ring with $m^2 \neq 0$. Then $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, for every non-minimal ideal I of R and the intersection graph of ideals of R is complete if and only if R is an SPAP-ring.*

Proof. Let R be an SPAP-ring by Lemma 2.14, m^2 is the unique minimal ideal of R , so by Lemma 2.6, the intersection graph of ideals of R is complete. Now let I be a proper ideal of R , Lemma 2.13 gives $I = 0$ or $I = m^2$ or $I^2 = m^2$. If I is a non-zero non-minimal ideal of R and $ab \in I \setminus I^2$, for $a, b \in R$, then $ab \notin I^2 = m^2$, so a or b is not in m , thus a or b is unit. Thus a or b must be in I . This shows that I is an almost prime ideal of R . Hence, by Lemma 2.1, $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$. In general, the zero ideal is an almost prime of R . Thus every non-minimal ideal of R is almost prime and so $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, for every non-minimal ideal I of R .

The converse of theorem is valid by Proposition 2.11. \square

EXAMPLE 2.16. Let k be an ordered field. Then for a non-empty set $\{x_\alpha\}_{\alpha \in \Delta}$ of indeterminates. Define $R = k[[\{x_\alpha\}_{\alpha \in \Delta}]]$, $m = \langle \{x_\alpha\}_{\alpha \in \Delta} \rangle$, and $J = \langle \{x_\alpha x_\beta, x_\alpha^2 - x_\beta^2\}_{\alpha \neq \beta}, \{x_\alpha^3\}_\alpha \rangle$. Let $\bar{R} = \frac{R}{J}$. Then \bar{R} is an SPAP-ring with $\bar{m}^2 \neq 0$ and \bar{m} is not principal for $|\Delta| > 1$, see [3, Example 20]. If Δ is a finite set, then \bar{R} is a Noetherian SPAP-ring with $\bar{m}^2 \neq 0$, and thus $\Gamma_{\bar{I}}(\bar{R})$ is an induced subgraph of $\Gamma_{\bar{I}^2}(\bar{R})$, for every non-minimal ideal \bar{I} of \bar{R} and the intersection graph of ideals of \bar{R} is complete.

Theorem 2.17. *Let (R, m) be an Artinian local ring with $m^2 \neq 0$, such that $\Gamma_I(R)$ is an induced subgraph of $\Gamma_{I^2}(R)$, for every non-minimal ideal I of R and the intersection graph of ideals of R is complete. If $\text{char}(R) \neq p^2$, for any prime number p and $\text{char}(\frac{R}{m}) \neq 2$, then there exists a regular local ring (S, n) , a positive integer number h , and subset $\{x_a\}_{a=1, \dots, h}$ of n such that $R \cong \frac{S}{K}$ in which K is minimally generated by the elements $\{x_i x_j\}_{1 \leq i < j \leq h}$, $\{x_j^2\}_{2 \leq j \leq \tau}$ and $\{x_i^2 u_i x_1^2\}_{\tau+1 \leq i \leq h}$, where the u_i are unit in R and τ is the Cohen- Macaulay type of R .*

Proof. By Theorem 2.15 and [11, Proposition 6.3]. □

ACKNOWLEDGMENTS

The author would like to express its sincere thanks to the referees for their valuable suggestions and comments.

REFERENCES

1. A. Abbasi, H. Roshan-Shekalgourabi, D. Hassanzadeh-Lelekaami, Associated Graphs of Modules Over Commutative Rings, *Iran. J. Math. Sci. Inform.*, **10**(1), (2015), 45-58.
2. S. Akbari, R. Nikandish, M. J. Nikmehr, Some results on the intersection graphs of ideals of rings, *J. Algebra Appl.*, **12**(4), (2013), 125–200. [13 pages] DOI: 10.1142/S0219498812502003.
3. D. D. Anderson, M. Bataineh, Generalization of Prime Ideals, *Comm. Algebra*, **36**, (2008), 686-696.
4. A. Barnard, Multiplication modules, *J. Algebra*, **71**, (1981), 174-178.
5. I. Chakrabarty, S. Ghosh, T. K. Mukherjee, M. K. Sen, Intersection graphs of ideals of rings, *Discrete Math.*, **309**, (2009), 5381-5392.
6. I. Beck, Coloring of commutative rings, *J. Algebra*, **116**, (1988), 208-226.
7. S. M. Bhatwadekar, P. K . Sharma, Unique factorization and birth of almost primes, *Comm. Algebra*, **33**, (2005), 43-49.
8. H. R. Maimani, Median and center of zero-divisor graph of commutative semogroups, *Iran. J. Math. Sci. Inform.*, **3**(2), (2008), 69-76.
9. S. Mori, Allgemeine Z.P.I.-Ringe, *J. Sci. Hiroshima Univ. Ser. A.*, **10**, (1940), 117-136.
10. S. P. Redmond, An ideal-based zero-divisor graph of a commutative ring, *Comm. Algebra*, **31**, (2003), 4425-4443 .
11. E. Rostami, R. Nekooei, On SPAP-rings, *Bull. Iranian Math. Soc.*, **41**(4), (2015), 907-921.
12. A. Tehranian, H. R. Maimani, A study of the Total graph, *Iran. J. Math. Sci. Inform.*, **6**(2), (2011), 75-80.