Iranian Journal of Mathematical Sciences and Informatics
Vol. 18, No. 1 (2023), pp 41-54
DOI: 10.52547/ijmsi.18.1.41

The Hyper-Zagreb Index of Trees and Unicyclic Graphs

Hassan Rezapour[a], Ramin Nasiri[b*], Seyedahmad Mousavi[c]

[a]Department of Mathematics, Faculty of Basic Sciences, University of Qom, Qom, Iran
[b]Department of Mathematics, Faculty of Sciences, Imam Hossein Comprehensive University, Tehran, Iran
[c]Department of Mathematics and Statistics, University of Maryland, Baltimore County, Baltimore, MD 21250, U.S.A.

E-mail: hassan.rezapour@gmail.com
E-mail: rnasiri@ihu.ac.ir
E-mail: smousav1@umbc.edu

Abstract. Topological indices are widely used as mathematical tools to analyze different types of graphs emerged in a broad range of applications. The Hyper-Zagreb index (HM) is an important tool because it integrates the first two Zagreb indices. In this paper, we characterize the trees and unicyclic graphs with the first four and first eight greatest HM-value, respectively.

Keywords: Hyper-Zagreb index, Vertex degree, Unicyclic graphs, Trees.

2000 Mathematics subject classification: 05C05, 05C07, 05C35.

1. Introduction

A nonnegative number is assigned to a graph G to define an associated topological index if it is the same for every isomorphic graph of G, i.e., it is graph invariant. Topological indices are appropriate tools to mathematically investigate and properly comprehend molecular structures and their properties

*Corresponding Author

Received 06 July 2018; Accepted 08 May 2019
©2023 Academic Center for Education, Culture and Research TMU

[Downloaded from ijmsi.ir on 2026-10-14]
[DOI: 10.52547/ijmsi.18.1.41]

such as complexity [9, 10]. The first topological index is proposed by Wiener [24] in order to examine chemical features of paraffin. Since trees demonstrate a remarkable importance in various applications, authors in [4] specifically investigate this index for this setting. Moreover, In [20], the extremal unicyclic graphs with respect to Wiener index is studied. The Hyper-Wiener index for acyclic structures is due to Randic, where later [15] extends this notion so that it applies applied for any connected graphs. An interested reader can explore some chemical applications of the Hyper-Wiener index in [12]. Zagreb indices were first suggested by Gutman et al. [13] in the 1970s, which absorbed attention of many scientists in different fields. The reader is encouraged to consult with [1, 3, 11, 14, 21, 25, 27] for more useful information. A comprehensive study on relations between the mentioned indices is found in [26].

All graphs in this paper are simple, finite and undirected. The vertex and edge sets of a graph G are shown by $V(G)$ and $E(G)$, respectively. Also, $n(G)$ denotes the number of vertices of G, which is called its order.

For a graph G, the *Hyper-Zagreb index* of G is defined as the following

$$HM(G) = \sum_{xy \in E(G)} \left(d_G(x) + d_G(y)\right)^2, \tag{1.1}$$

where $d_G(x)$ is the degree of vertex x. For the edge $xy \in E(G)$, if consider $h_G(xy) := \left(d_G(x) + d_G(y)\right)^2$. Then, the above formulation can be equivalently written as

$$HM(G) = \sum_{xy \in E(G)} h_G(xy).$$

This was initially presented by Shirdel et al. [23] in 2013. They consider two simply connected graphs and compute this distance-based index for the resulted Cartesian product, composition, join and disjunction graphs. Gao et al. [7] discuss acyclic, unicyclic, and bicyclic graphs and find sharp bounds for their Hyper-Zagreb index. The degree of vertices is the main part of some other graph invariants such as irregularity and total irregularity, see [6, 17, 18, 19]. There is an extensive literature on this topic including [2, 5, 8, 22, 16].

2. Preliminaries and lemmas

In this section, we first bring several notations and definitions. Then, we propose different propositions which are essential for the subsequent section.

Unicylcic graph G of order n with circuit $C_m = x_1x_2 \dots x_mx_1$ of length m is denoted by $C_m^{u_1,u_2,\dots,u_k}(T_1, T_2, \dots, T_k)$ in which trees T_i's for $i = 1, 2, \dots, k$ are all nontrivial components of $G - E(C_m)$ and u_i $(i = 1, 2, \dots, k)$ is the common vertex of T_i and C_m. Specially, $G = C_n$ for $k = 0$. For convenience, we denote $C_m^{u_1,u_2,\dots,u_k}(T_1, T_2, \dots, T_k)$ by $C_m(T_1, T_2, \dots, T_k)$, for any integer number $k \geq 1$. Let $n(T_i) = l_i + 1$, $i = 1, 2, \dots, k$, then $l = \sum_{i=1}^{k} l_i = n - m$. Also, if

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]



FIGURE 1. Tree $T^x(T_1, T_2, \ldots, T_\Delta)$.

a tree T_i is the star S_{l_i+1} then we replace it by l_i, for example we denote $C_4(T_1, S_5, T_3, S_9)$ by $C_4(T_1, 4, T_3, 8)$.

Let T be a *tree* with n vertices ($n \geq 2$) such that $x \in V(T)$ and x has a maximum degree of vertices in graph T, i.e. $\Delta = d_T(x) = \max\{d_T(u), u \in V(T)\}$. T is shown by $T^x(T_1, T_2, \ldots, T_\Delta)$, where $T_i = T_i^* + \{y_i x\}$, $i = 1, 2, \ldots, \Delta$, and $T_1^*, T_2^*, \ldots, T_\Delta^*$ are trees with disjoint vertex sets and $n_1, n_2, \ldots, n_\Delta$ are numbers of their vertices, respectively. Therefore, we have $|V(T_i)| = |V(T_i^*)| + 1 = n_i + 1$, $i = 1, 2, \ldots, \Delta$, and $n = |V(T)| = \sum_{i=1}^{\Delta} n_i + 1$ and $y_i \in V(T_i^*)$. Moreover, $E(T_i) = E(T_i^*) \cup \{y_i x\}$ and $V(T_i) = V(T_i^*) \cup \{x\}$ (see Figure 1).

The *coalescence* of G and H is denoted by $G(u)oH(v)$ and obtained by identifying the vertex u of G with the vertex v of H.

Lemma 2.1. *Assume that $z \in V(H)$ and $\{u, w\} \subseteq V(G)$ such that the following conditions hold:*

(a) $d_G(u) \leq d_G(w)$,

(b) $$\sum_{x \in N_G(u) \setminus \{w\}} d_G(x) \leq \sum_{x \in N_G(w) \setminus \{u\}} d_G(x).$$

Moreover, let $G_1 = G(u)oH(z)$ and $G_2 = G(w)oH(z)$, where G_1 and G_2 are as shown in Figure 2 . Then, $HM(G_2) \geq HM(G_1)$, with the equality if and only if equality holds in both given conditions.

Proof. Recall that

$$\sum_{x \in N_G(w) \setminus \{u\}} h_{G_1}(xw) = \sum_{x \in N_G(w) \setminus \{u\}} \left(d_G(x) + d_G(w)\right)^2,$$

$$\sum_{x \in N_H(z)} h_{G_1}(xz) = \sum_{x \in N_H(z)} \left(d_H(z) + d_G(u) + d_H(x)\right)^2$$

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]



FIGURE 2. The transformation of two graphs.

and $h_{G_1}(uw) = h_{G_2}(uw) = (d_G(z) + d_G(w) + d_H(x))^2$. In addition, one has

$$\sum_{x\in N_H(z)} h_{G_2}(zx) = \sum_{x\in N_H(z)} (d_H(z) + d_G(w) + d_H(x))^2,$$
$$\sum_{x\in N_G(u)\setminus\{w\}} h_{G_2}(xu) = \sum_{x\in N_G(u)\setminus\{w\}} (d_G(u) + d_G(x))^2$$

and

$$\sum_{x\in N_G(w)\setminus\{u\}} h_{G_2}(xw) = \sum_{x\in N_G(w)\setminus\{x\}} (d_G(w) + d_G(x) + d_H(z))^2.$$

We consider two cases where either $uw \in E(G)$ or $uw \notin E(G)$. First, suppose that $uw \in E(G)$. For $i = 1$ and 2, we have

$$HM(G_i) = \sum_{\substack{xy\in E(G)\\ x,y\notin\{u,w\}}} h_G(xy) + \sum_{x\in N_G(u)\setminus\{w\}} h_{G_i}(xu) + \sum_{x\in N_G(w)\setminus\{u\}} h_{G_i}(xw)$$
$$+ h_{G_i}(uw) + \sum_{x,y\neq z} h_H(xy) + \sum_{x\in N_H(z)} h_{G_i}(xz).$$

On the other hand,

$$\sum_{x\in N_G(u)\setminus\{w\}} h_{G_1}(xu) = \sum_{x\in N_G(u)\setminus\{w\}} (d_G(u) + d_G(x) + d_H(x))^2.$$

Therefore,

$$\begin{aligned} &HM(G_2) - HM(G_1)\\ &= \sum_{x\in N_G(u)\setminus\{w\}} \Big((d_G(u) + d_G(x))^2 - (d_G(u) + d_G(x) + d_H(z))^2\Big)\\ &\quad + \sum_{x\in N_G(w)\setminus\{u\}} \Big((d_G(w) + d_G(x) + d_H(z))^2 - (d_G(w) + d_G(u))^2\Big)\\ &\quad + \sum_{x\in N_H(z)} \Big((d_H(z) + d_G(w) + d_H(x))^2 - (d_H(z) + d_G(u) + d_H(w))^2\Big) \end{aligned}$$

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

this implies that

$$\begin{aligned} HM(G_2) - HM(G_1) \geq & 2d_H(z)\left(d_G(u)\left(d_G(w)-1\right) - d_G(u)\left(d_G(u)-1\right)\right) \\ & + 2d_H(z)\left(\sum_{x\in N_G(w)\setminus\{u\}} d_G(u) - \sum_{x\in N_G(w)\setminus\{w\}} d_G(x)\right) \\ \geq & 0. \end{aligned}$$

Now, suppose that $uw \notin E(G)$. Then, for $i = 1$ and 2, we have

$$\begin{aligned} HM(G_i) = & \sum_{\substack{xy\in E(G)\\ x,y\notin\{u,w\}}} h_G(xy) + \sum_{x\in N_G(u)} h_{G_i}(xu) + \sum_{x\in N_G(w)} h_{G_i}(xw) \\ & + \sum_{x,y\neq z} h_H(xy) + \sum_{x\in N_{G_i}} h_{G_i}(xz). \end{aligned}$$

Also, in this case one has

$$\sum_{x\in N_G(w)\setminus\{u\}} d_G(x) = \sum_{x\in N_G(w)} d_G(x), \sum_{x\in N_G(u)\setminus\{w\}} d_G(x) = \sum_{x\in N_G(u)} d_G(x).$$

Hence, a similar approach as the previous case can be used to prove the result. □

Lemma 2.2. *Suppose u and v are vertices of graphs G_1 and G_2, respectively. Let G be the graph obtained by joining $u \in V(G_1)$ to $v \in V(G_2)$ by an edge, and G' be the graph obtained by identifying $u \in V(G_1)$ with $v \in V(G_2)$ and attaching a pendent vertex to the common vertex as shown in Figure 3. Then if $d_G(u), d_{G'}(v) \geq 2$, we have $HM(G) < HM(G')$.*



FIGURE 3. An illustration of graphs in Lemma 2.2.

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

Proof. Assume that the graph G' is obtained by identifying $u \in V(G_1)$ with $v \in V(G_2)$ and attaching a pendent vertex w to the common vertex. Then,

$$\begin{aligned} HM(G) =& h_G(uv) + \sum_{x \in N_{G_1}(u)} h_G(ux) + \sum_{x \in N_{G_2}(v)} h_G(vx) + \sum_{\substack{xy \in E(G_1) \\ u \notin \{x,y\}}} h_{G_1}(xy) \\ &+ \sum_{xy \in E(G_2), v \notin \{x,y\}} h_{G_2}(xy) \end{aligned}$$

and

$$\begin{aligned} HM(G') =& h_{G'}(uw) + \sum_{x \in N_{G_1}(u)} h_{G'}(ux) + \sum_{x \in N_{G_2}(v)} h_{G'}(vx) + \sum_{\substack{xy \in E(G_1) \\ u \notin \{x,y\}}} h_{G_1}(xy) \\ &+ \sum_{\substack{xy \in E(G_2) \\ v \notin \{x,y\}}} h_{G_2}(xy). \end{aligned}$$

Since $d_G(u) = d_{G_1}(u) + 1$, $d_G(v) = d_{G_1}(v) + 1$, $d_{G'}(w) = 1$ and $d_{G'}(u) = d_{G'}(v) = d_{G_1}(u) + d_{G_2}(v) + 1$ we have

$$\sum_{x \in N_{G_1}(u)} h_G(ux) < \sum_{x \in N_{G_1}(u)} h_{G'}(ux), \qquad \sum_{x \in N_{G_2}(v)} h_G(vx) < \sum_{x \in N_{G_2}(v)} h_{G'}(vx)$$

and $h_G(uv) = h_{G'}(uw) = (d_{G_1}(u) + d_{G_2}(v) + 2)^2$. Hence,

$$\begin{aligned} HM(G') - HM(G) =& \sum_{x \in N_{G_1}(u)} h_{G'}(ux) - \sum_{x \in N_{G_1}(u)} h_G(ux) \\ &+ \sum_{x \in N_{G_2}(v)} h_{G'}(vx) - \sum_{x \in N_{G_2}(v)} h_G(vx) \\ >& 0. \end{aligned}$$

□

Corollary 2.3. *Let T be a tree with n vertices. Then, $HM(T) \leq HM(S_n)$, with the equality if and only if $T \cong S_n$.*

Corollary 2.4. *Let $G = C_m(T_1, T_2, \ldots, T_k)$ be a unicyclic graph and $n(T_i) = l_i + 1$. Then, $HM(G) \leq HM(C_m(l_1, l_2, \ldots, l_k))$, with the equality if and only if $T_i \cong S_{l_i+1}$, $i = 1, 2, \ldots, k$.*

Lemma 2.5. *Let $G_1 = C_m(l_1, l_2, \ldots, l_k)$ be a unicyclic graph and $y_1u_i, u_iu_{i+1} \in E(C_m)$ such that $d_{G_1}(y_1), d_{G_1}(u_i) \leq d_{G_1}(u_{i+1})$, then for $G_2 = C_m(l_1, \ldots, l_{i-1}, l_{i+1} + l_i, l_{i+2}, \ldots, l_k)$ one has that $HM(G_1) < HM(G_2)$.*

Proof. Let $G = C_m(l_1, \ldots, l_{i-1}, l_{i+1}, l_{i+2}, \ldots, l_k)$, then $2 = d_G(u_i) < 3 \leq d_G(u_{i+1})$; meaning that the condition *(a)* in Lemma 2.1 holds. Hence, we now

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

show that the second condition in this Lemma is also satisfied. Suppose that $y_2u_{i+1} \in E(C_m)$. By a simple calculation one can check that

$$\begin{aligned}\sum_{x\in N_G(u_i)\setminus\{u_{i+1}\}} d_G(x) =& d_G(y_1),\\ \sum_{x\in N_G(u_{i+1})\setminus\{u_i\}} d_G(x) =& \sum_{\substack{x\in N_G(u_{i+1})\setminus\{u_i\}\\ x\in V(C_m)}} d_G(x) + \sum_{\substack{x\in N_G(u_{i+1})\setminus\{u_i\}\\ x\notin V(C_m)}} d_G(x)\\ =& d_G(y_2) + \sum_{\substack{x\in N_G(u_{i+1})\setminus\{u_i\}\\ x\notin V(C_m)}} 1\\ =& d_G(y_2) + d_G(u_{i+1}) - 2.\end{aligned}$$

Moreover, $d_{G_1}(u_{i+1}) = d_G(u_{i+1})$ and $d_G(y_1) = d_{G_1}(y_1)$. On the other hand, since $y_2 \in V(C_m)$ then $d_G(y_2) \geq 2$; implying that $d_G(y_2) - 2 \geq 0$. So, we have

$$\textstyle\sum_{x\in N_G(u_i)\setminus\{u_{i+1}\}} d_G(x) = d_G(y_1) = d_{G_1}(y_1) \leq d_{G_1}(u_{i+1}) = d_G(u_{i+1}) \leq \sum_{x\in N_G(u_{i+1})\setminus\{u_i\}} d_G(x).$$

Therefore, the condition *(b)* of Lemma 2.1 holds, which completes the proof. □

Lemma 2.6. *Let $G = C_m^{u_1,u_2,\ldots,u_k}(l_1, l_2, \ldots, l_k)$ be a unicyclic graph and $k > 1$. Then if $u_iu_{i+1} \in E(C_m)$, $i = 1, 2, \ldots, k-1$, then $HM(G) < HM(C_m(n-m))$. Otherwise, there exist positive integers $l'_1, l'_2, \ldots, l'_r$ $(r \leq k)$, such that $HM(G) < HM(G') < HM(G'')$, where $G' = C_m^{v_1,v_2,\ldots,v_r}(l'_1, l'_2, \ldots, l'_r)$,*
$G'' = C_m^{v_2,v_3,\ldots,v_r}(l'_1 + l'_2, l'_3, \ldots, l'_r)$, $d_{G'}(v_i, v_j) \geq 2$ for $1 \leq i < j \leq r$ and $\{v_1, v_2, \ldots, v_r\} \subseteq \{u_1, u_2, \ldots, u_k\}$.

Proof. $HM(G) < HM(G')$ is straightforward in light of Lemma 2.5. Now, by considering $u = v_1, w = v_2$, $H = S_{l'_1+1}$ and $G = C_m^{v_2,v_3,\ldots,v_r}(l'_2, l'_3, \ldots, l'_r)$ and using Lemma 2.1 we can conclude that $HM(G') < HM(G'')$, as desired. □

Lemma 2.7. *Let $G = C_m(l_1, l_2, \ldots, l_k)$ be a unicyclic graph of order n. Then $HM(G) \leq HM(C_m(n-m))$, with equality if and only if $k = 1$.*

Proof. The proof is obtained by applying Lemmas 2.5 and 2.6. □

Corollary 2.8. *Let $G = C_m(T_1, T_2, \ldots, T_k)$ be a unicyclic graph and $n(T_i) = l_i + 1$, $i = 1, 2, \ldots, k$. Then,*

$$HM(G) \leq HM(C_m(l_1, l_2, \ldots, l_k)) \leq HM(C_m(n-m)),$$

with left equality if and only if $T_i \cong S_{l_i+1}$, $i = 1, 2, \ldots, k$, and right equality if and only if $k = 1$.

Lemma 2.9. *Let $G_1 = C_m(n-m)$ and $G_2 = C_{m-1}(n-m+1)$, $m \geq 4$, be unicyclic graphs of order n. Then, $HM(G_1) < HM(G_2)$.*

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

Proof. By a simple calculation we have

$$\begin{aligned} HM(G_1) =& 4\,(m-2) + 2\,(n-m+4)^2 + (n-m)\,(n-m+3)^2 \\ <& 4\,(m-3) + 2\,(n-m+5)^2 + (n-m+1)\,(n-m+4)^2 \\ =& HM(G_2). \end{aligned}$$

As desired. □

Lemma 2.10. *Let $G = C_m\,(T_1, T_2, \ldots, T_k)$ be a unicyclic graph. Then,*

$$HM(G) = \sum_{i=1}^{k} \sum_{xy \in E(T_i)} h_G(xy) + \sum_{xy \in E(C_m)} h_G(xy).$$

Proof. The proof is trivial by the Hyper-Zagreb index definition (1.1). □

3. Main Results

In this section, we characterize the trees and unicyclic graphs with the first four and first eight greatest HM-value, respectively.

Theorem 3.1. *Let T be a tree with n vertices. If $T \ncong S_n, T_n^1$ or T_2^n, then*

$$HM(T) \leq HM\left(T_n^3\right) < HM\left(T_n^2\right) < HM\left(T_n^1\right) < HM\left(S_n\right),$$

with the equality if and if $T \cong T_3^n$, where T_n^1, T_n^2 and T_n^3 are given as in Figure 4.



Figure 4. Some trees with large Hyper-Zagreb values.

Graph	HM-value
S_n	$n^3 - n^2$
T_1^n	$n^3 - 4n^2 + 7n + 6$
T_2^n	$n^3 - 7n^2 + 20n + 16$
T_3^n	$n^3 - 7n^2 + 20n$

Table 1. Trees with large Hyper-Zagreb values.

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

Proof. Using Table 1, we have $HM(S_n) > HM\left(T_n^1\right) > HM\left(T_n^2\right) > HM\left(T_n^3\right)$. Hence, we need to prove that $HM(T) < HM\left(T_n^3\right)$ when $T \ncong T_n^3$. Let $T = T^x(T_1, T_2, \ldots, T_\Delta)$, where $\Delta = d_T(x)$. By Corollary 2.3, we have $HM(T_i) \leq HM(S_{n_i})$, $i = 1, 2, \ldots, \Delta$. Moreover, let $T' = T^x(T_1', T_2', \ldots T_\Delta')$, where $T_i'^* = S_{n_i}$, $i = 1, 2, \ldots, \Delta$, then we have $HM(T) \leq HM(T')$. To complete the proof, we consider three different cases as follows:

Case 1: assume that $d_T(y_i) = 1$, $i = 1, 2, \ldots, \Delta$, then $T = S_n$. This is a contradiction to the assumption.

Case 2: assume that there exists y_t for $t = 1, 2, \ldots, \Delta$ such that $d_T(y_t) \geq 2$ and $d_T(y_i) = 1$ for $i = 1, 2, \ldots, \Delta$ and $i \neq t$. In this case, there are three subcases that can happen:

(i) If $|V(T_t^*)| = 2$, then $T \cong T_1^n$. This is clearly a contradiction.

(ii) If $|V(T_t^*)| = 3$, then we must consider that $d_T(y_t) = 2$ or 3. The case $d_T(y_t) = 3$ implies that $T \cong T_n^2$, which is a contradiction. If $d_T(y_t) = 2$, then

$$\begin{aligned} HM(T) =& (n-4)(n-2)^2 + (n-1)^2 + 16 + 9 \\ =& n^3 - 7n^2 + 18n + 10 \\ <& n^3 - 7n^2 + 20n \\ =& HM\left(T_n^3\right). \end{aligned}$$

(iii) If $|V(T_t^*)| \geq 4$, then $T = T^x\left(\overbrace{S_2, \ldots, S_2}^{t-1 \text{ times}}, T_t, \overbrace{S_2, \ldots, S_2}^{\Delta - t \text{ times}}\right)$. By Corollary 2.3, the Hyper-Zagreb index for T is maximum when $T_t^* = S_{n_t}$. On the other hand, it follows from Lemma 2.1 that if $n_t = 4$ than T has maximum HM-value, i.e. in this case T has maximum HM-value when $T \cong T_n^4$ (see Figure 4). Hence, applying Lemma 2.1, it is clear that $HM(T) \leq HM\left(T_n^4\right) < HM\left(T_n^3\right)$.

Case 3: Suppose that there exist $1 \leq s, t \leq \Delta$ such that $d_T(y_s), d_T(y_t) \geq 2$. Similar to previous, applying Corollary 2.3 and Lemma 2.1, it can concluded that $HM(T) < HM\left(T_n^3\right)$. □

Theorem 3.2. *Let G be a unicylcic graph of order $n \geq 15$. If $G \ncong C_3(n-3)$, $C_3(1, n-4)$, $C_3\left(T_{n-2}^1\right)$, $C_4(n-2)$, $C_3(2, n-5)$, $C_3(1, 1, n-5)$ and $C_3\left(T_{n-2}^2\right)$. Then,*

$$\begin{aligned} HM(G) \leq& HM\left(C_3\left(T_{n-2}^3\right)\right) < HM\left(C_3\left(T_{n-2}^2\right)\right) < HM\left(C_3(1, 1, n-5)\right) \\ <& HM\left(C_3(2, n-5)\right) < HM\left(C_4(n-4)\right) < HM\left(C_3\left(T_{n-2}^1\right)\right) \\ <& HM\left(C_3(1, n-4)\right) < HM\left(C_3(n-3)\right), \end{aligned}$$

with the equality if and only if $G \cong C_3\left(T_{n-2}^3\right)$ or $G \cong C_3(P_3, 10)$ for $n = 15$ (see Figure 5).

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]



FIGURE 5. The unicyclic graphs with the first eight greatest Hyper-Zagreb.

Graph	HM-value
$C_3\,(n-3)$	$n^3-n^2+4n+18$
$C_3\,(1,n-4)$	$n^3-4n^2+11n+38$
$C_3\left(T^1_{n-2}\right)$	$n^3-4n^2+11n+20$
$C_4\,(n-4)$	$n^3-4n^2+9n+28$
$C_3\,(2,n-5)$	$n^3-7n^2+24n+68$
$C_3\,(1,1,n-5)$	$n^3-7n^2+24n+48$
$C_3\left(T^2_{n-2}\right)$	$n^3-7n^2+24n+26$
$C_3\left(T^3_{n-2}\right)$	$n^3-7n^2+24n+10$

TABLE 2. Unicyclic graphs with large Hyper-Zagreb values.

Proof. Assume that $G=C_m\,(T_1,T_2,\ldots,T_k)$ be a unicyclic graph and $n(T_i)=l_i+1$, $i=1,2,\ldots,k$. The given Table 2 provides the Hyper-Zagreb index of some graphs by which the result is trivial. It is enough to discuss about the equality case. If $G\cong C_3\left(T^3_{n-2}\right)$, then $HM(G)=HM\left(C_3\left(T^3_{n-2}\right)\right)$. Also, if $G\cong C_3\,(P_3,10)$ for $n=15$, then $HM(G)=2170=HM\left(C_3\left(T^3_{13}\right)\right)$. We now prove that $HM(G)<HM\left(C_3\left(T^3_{n-2}\right)\right)$, where $G\ncong C_3\,(n-3)$, $C_3\,(1,n-4)$, $C_3\left(T^1_{n-2}\right)$, $C_4\,(n-2)$, $C_3\,(2,n-5)$, $C_3\,(1,1,n-5)$ and $C_3\left(T^2_{n-2}\right)$. We examine three cases of $m=3,4$ and 5 for $G=C_m\,(T_1,T_2,\ldots,T_k)$ as follows:

Case 1: $m=3$. We need to discuss three subcases that $k=1,2$ and 3.

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

(i) $k = 1$, then $G = C_3(T_1)$. By assumption, we know that $T_1 \ncong S_{n-2}, T^1_{n-2}, T^2_{n-2}$ and T^3_{n-2}. So, Theorem 3.1 implies that $HM(T_1) < HM(T^3_{n-2})$. By Lemma 2.10, we get $HM(G) < HM\left(C_3\left(T^3_{n-2}\right)\right)$.

(ii) $k = 2$, then $G = C_3(T_1, T_2)$. By assumption, $G \ncong C_3(1, n-4)$ and $C_3(2, n-5)$. By Corollaries 2.3, 2.4 and Lemmas 2.1, 2.10, the maximum value of $HM(G)$ happens when $G \cong C_3(3, n-6)$ or $C_3(P_3, n-5)$. The first case yields that

$$HM(G) \leq HM(C_3(3, n-6)) = n^3 - 10n^2 + 43n + 108.$$

Hence, we have (for $n \geq 15$) that

$$\begin{aligned} HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(G) \geq & HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(C_3(3, n-6)) \\ = & \left(n^3 - 7n^2 + 24n + 10\right) \\ & - \left(n^3 - 10n^2 + 43n + 108\right) \\ = & 3n^2 - 19n - 98 \\ > & 0. \end{aligned}$$

Similarly, for the second case we have

$$\begin{aligned} HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(G) \geq & HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(C_3(P_3, n-5)) \\ = & \left(n^3 - 7n^2 + 24n + 10\right) - \left(n^3 - 7n^2 + 22n + 40\right) \\ = & 2n - 30 \\ > & 0. \end{aligned}$$

This means that in both cases $HM(G) < HM\left(C_3\left(T^3_{n-2}\right)\right)$.

(iii) $k = 3$, then $G = C_3(T_1, T_2, T_3)$. By Corollary 2.4, it is simple to see that $HM(G) \leq HM(C_3(l_1, l_2, l_3))$. On the other hand, since by assumption $G \ncong C_3(1, 1, n-5)$, the Hyper-Zagreb index attains its maximum when
$G \cong C_3(1, 2, n-6)$. Hence,

$$\begin{aligned} HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(G) \geq & HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(C_3(1, 2, n-6)) \\ = & \left(n^3 - 7n^2 + 24n + 10\right) - \left(n^3 - 10n^2 + 43n + 62\right) \\ > & 0. \end{aligned}$$

Case 2: $m = 4$. This needs to be analyzed for $k = 1, 2, 3$ and 4.

(i) $k = 1$, then $G = C_4(T_1)$. Since $G \ncong C_4(n-4)$, we have $T_1 \ncong S_{n-3}$. Note that G has a maximum value of the Hyper-Zagreb index if $T_1 \cong T^1_{n-3}$ by Theorem 3.1 and Lemma 2.10. Moreover, we have (for $n \geq 15$)

$$\begin{aligned} HM\left(C_3\left(T^3_{n-3}\right)\right) - HM(G) \geq & HM\left(C_3\left(T^3_{n-3}\right)\right) - HM\left(C_4\left(T^1_{n-3}\right)\right) \\ = & \left(n^3 - 7n^2 + 24n + 10\right) - \left(n^3 - 7n^2 + 22n + 20\right) \\ > & 0. \end{aligned}$$

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

(ii) $k=2$, then $G=C_4^{u_1,u_2}(T_1,T_2)_\alpha=C_4\left(T_1,T_2\right)_\alpha$, where $\alpha=d_G(u_1,u_2)$. By Lemma 2.1, G attains maximum HM-value if $G\cong C_4\left(l_1,l_2\right)_{\alpha=1}$. This lemma also implies that

$$HM\left(C_4\left(l_1,l_2\right)\right)_{\alpha=1}\leq HM\left(C_4\left(1,n-5\right)\right)_{\alpha=1}=n^3-7n^2+22n+38.$$

Therefore, for $n\geq 15$, we have

$$HM(G)\leq n^3-7n^2+22n+38<n^3-7n^2+24n+10=HM\left(C_3\left(T_{n-2}^3\right)\right).$$

(iii) $k=3$, then G is considered as $C_4\left(T_1,T_2,T_3\right)$. By Corollary 2.4 and Lemmas 2.1, 2.10, for $n\geq 15$ we have

$$\begin{aligned}HM\left(C_3\left(T_{n-2}^3\right)\right)-HM(G)\geq&HM\left(C_3\left(T_{n-2}^3\right)\right)-HM\left(C_4\left(l_1,l_2,l_3\right)\right)\\>&HM\left(C_3\left(T_{n-2}^3\right)\right)-HM\left(C_4\left(1,n-5\right)\right)_{\alpha=1}\\=&\left(n^3-7n^2+24n+10\right)-\left(n^3-7n^2+22n+38\right)\\>&0.\end{aligned}$$

(iv) $k=4$, then $G=C_4(T_1,T_2,T_3,T_4)$. In a similar way, one can see easily that $HM(G)<HM\left(C_3\left(T_{n-2}^3\right)\right)$; completing the proof of the second case.

Case 3: $m\geq 5$. Using Lemmas 2.9, 2.10 and Corollaries 2.3, 2.8, we conclude that (for $n\geq 15$)

$$\begin{aligned}HM\left(C_3\left(T_{n-2}^3\right)\right)-HM(G)\geq&HM\left(C_3\left(T_{n-2}^3\right)\right)-HM\left(C_m\left(l_1,l_2,\ldots,l_k\right)\right)\\\geq&HM\left(C_3\left(T_{n-2}^3\right)\right)-HM\left(C_m\left(n-m\right)\right)\\\geq&HM\left(C_3\left(T_{n-2}^3\right)\right)-HM\left(C_5\left(n-5\right)\right)\\=&\left(n^3-7n^2+24n+10\right)-\left(n^3-7n^2+20n+30\right)\\>&0.\end{aligned}$$

□

4. Conclusion

In this paper, we studied the Hyper-Zagreb index and characterized the trees and unicyclic graphs with the first four and first eight greatest HM-value. It would be of interest to investigate its behavior on other classes of graphs with simple connectivity patterns and cyclic structures.

Acknowledgments

The authors would like to thank the referees for their valuable comments.

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

References

1. S. Baby, M. K. Shafiq, H. M. A. Siddiqui, On Zagreb Indices and Zagreb Polynomials of Single-walled Titania Nanotubes, *Science International*, **28**(4), (2016).
2. B. Basavanagoud, S. Patil, The Hyper-Zagreb Index of Four Operations on Graphs, *Math. Sci. Lett.*, **6**(2), (2017), 193-198.
3. J. Braun, A. Kerber, M. Meringer, C. Rucker, Similarity of Molecular Descriptors: the Equivalence of Zagreb Indices and Walk Counts, *MATCH Commun. Math. Comput. Chem.*, **54**, (2005), 163-176.
4. A. Dobrynin, R. Entringer, I. Gutman, Wiener Index of Trees: Theory and Applications, *Acta Appl. Math.*, **66**(3), (2001), 211-249.
5. F. Falahati-Nezhad, M. Azari, Bounds on the Hyper-Zagreb Index, *J. Appl. Math. Inform.*, **34**, (2016), 319-330.
6. G. H. Fath-Tabar, I. Gutman, R. Nasiri, Extremely Irregular Trees, *Bull. Acad. Serbe Sci. Arts (Cl. Sci. Math. Natur)*, **38**, (2013), 1-8.
7. W. Gao, M. K. Jamil, A. Javed, M. R. Farahani, Sh. Wang, J. B. Liu, Sharp Bounds of the Hyper-Zagreb Index on Acyclic, Unicylic, and Bicyclic Graphs,*Discrete Dyn. Nat. Soc.*, (2017).
8. W. Gao, M. K. Jamil, M. R. Farahani, The Hyper-Zagreb Index and Some Graph Operations, *J. Appl. Math. Comput.*, **54**(1-2), (2017), 263-275.
9. W. Gao, M. K. Siddiqui, Molecular Descriptors of Nanotube, Oxide, Silicate, and Triangulene Networks, *J. Chem.*, Article ID 6540754.10 (2017).
10. W. Gao, W. Wang, M. R. Farahani, Topological Indices Study of Molecular Structure in Anticancer Drugs, *J. Chem.*, (2016).
11. I. Gutman, K. Ch. Das, The First Zagreb Index 30 Years After, *MATCH Commun. Math. Comput. Chem.*, **50**, (2004), 83-92.
12. I. Gutman, W. Linert, I. Lukovits, A. Dobrynin, Trees with Extremal Hyper-Wiener Index: Mathematical Basis and Chemical Applications, *J. Chem. Inf. Comput. Sci.*, **37**(2), (1997), 349-354.
13. I. Gutman, N. Trinajstić, Graph Theory and Molecular Orbitals. Total φ-electron Energy of Alternant Hydrocarbons, *Chem. Phys. Lett.*, **17**(4), (1972), 535-538.
14. S. M. Kang, M. Munir, A. R. Nizami, Z. Shahzadi, W. Nazeer, Some Topological Invariants of the Mbius Ladder, *Global J.Pure Appl. Math.*, **12**, (2016), 5317-5327.
15. D. J. Klein, I. Lukovits, I. Gutman, On the Definition of the Hyper-Wiener Index for Cycle-containing Structures, *J. Chem. Inf. Comput. Sci.*, **35**(1), (1995), 50-52.
16. V. R. Kulli, B. Chaluvaraju, H. S. Boregouda, Some Degree Based Connectivity Indices of Kulli Cycle Windmill Graphs, *South Asain J. Math.*, **6**(6), (2016), 263-268.
17. R. Nasiri, H. R. Ellahi, A. Gholami, G. H. Fath-Tabar, The Irregularity and Total Irregularity of Eulerian Graphs, *Iranian J. Math. Chem.*, **9**(2), (2018), 101-111.
18. R. Nasiri, A. Gholami, G. H. Fath-Tabar, H. R. Ellahi, Extremely Irregular Unicyclic Graphs, *Kragujevac J. Math.*, **43**(2), (2019), 281-292.
19. R. Nasiri, G. H. Fath-Tabar, The Second Minimum of the Irregularity of Graphs, *Electron. Notes Discret. Math.*, **45**, (2014), 133-140.
20. R. Nasiri, H. Yousefi-Azari, M. R. Darafsheh, A. R. Ashrafi, Remarks on the Wiener Index of Unicyclic Graphs, *J. Appl. Math. Comput.*, **41**, (2013), 49-59.
21. S. Nikolić, G. Kovacević, A. Milicević, N. Trinajstić, The Zagreb Indices 30 Years After, *Croat. Chem. Acta.*, **76**(2), (2003), 113-124.
22. M. K. Siddiqui, M. Imran, A. Ahmad,. On Zagreb Indices, Zagreb Polynomials of Some Nanostar Dendrimers, *Appl. Math. Comput.*, **280**, (2016), 132-139.

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]

23. G. H. Shirdel, H. Rezapour, A. M. Sayadi, The Hyper-Zagreb Index of Graph Operations, *Iranian J. Math. Chem.*, **4**(2), (2013), 213-220.
24. H. Wiener, Structural Determination of Paraffin Boiling Points, *J. Am. Chem. Soc.*, **69**(1), (1947), 17-20.
25. B. Zhou, Zagreb Indices, *MATCH Commun. Math. Comput. Chem.*, **52**, (2004), 113-118.
26. B. Zhou, I. Gutman, Relations Between Wiener, Hyper-Wiener and Zagreb Indices, *Chem Phys Lett.*, **394**(1), (2004), 93-95.
27. B. Zhou, I. Gutman, Further Properties of Zagreb Indices, *MATCH Commun. Math. Comput. Chem.*, **54**, (2005), 233-239.

[Downloaded from ijmsi.ir on 2026-10-14]

[DOI: 10.52547/ijmsi.18.1.41]