

Some Hermite-Hadamard Type Inequalities for Convex Functions Defined on Convex Bodies Via Gauss-Ostrogradsky Identity

Silvestru Sever Dragomir^{a,b}

^aApplied Mathematics Research Group, ISILC, Victoria University, PO Box 14428 Melbourne City, MC 8001, Australia

^bDepartment of Mathematical and Physical Sciences, La Trobe University, Melbourne, VIC, Australia

E-mail: sever.dragomir@vu.edu.au

ABSTRACT. In this paper, by the use of Gauss-Ostrogradsky identity, we establish some integral inequalities of Hermite-Hadamard type for functions of three variables defined on closed and bounded convex bodies of the Euclidean space $\mathbb{R}^3$. Some examples for 3-dimensional balls are also provided.

Keywords: Hermite-Hadamard inequality, Triple integral inequalities, Gauss-Ostrogradsky identity.

2000 Mathematics subject classification: 26D15.

1. INTRODUCTION

Recall the following inequalities of Hermite-Hadamard's type for convex functions defined on a ball $B(C, R)$, where $C = (a, b, c) \in \mathbb{R}^3$, $R > 0$ and

$$B(C, R) := \left\{ (x, y, z) \in \mathbb{R}^3 \mid (x - a)^2 + (y - b)^2 + (z - c)^2 \leq R^2 \right\}.$$

The following theorem holds [6].

Theorem 1.1. *Let $f : B(C, R) \rightarrow \mathbb{R}$ be a convex mapping on the ball $B(C, R)$. Then we have the inequalities:*

$$\begin{aligned} f(a, b, c) &\leq \frac{1}{V(B(C, R))} \iiint_{B(C, R)} f(x, y, z) dx dy dz \\ &\leq \frac{1}{A(B(C, R))} \iint_{S(C, R)} f(x, y, z) dS, \end{aligned} \quad (1.1)$$

where

$$S(C, R) := \left\{ (x, y, z) \in \mathbb{R}^3 \mid (x-a)^2 + (y-b)^2 + (z-c)^2 = R^2 \right\}$$

and

$$V(B(C, R)) = \frac{4\pi R^3}{3}, \quad A(B(C, R)) = 4\pi R^2.$$

Let D be a bounded convex domain from $\mathbb{R}^3$ with a piecewise smooth boundary S . We use the notations

$$A(S) := \iint_S dS, \quad V(D) = \iiint_D dV,$$

$$\overline{x_D} := \frac{1}{V(D)} \iiint_D x dV, \quad \overline{y_D} := \frac{1}{V(D)} \iiint_D y dV, \quad \overline{z_D} := \frac{1}{V(D)} \iiint_D z dV$$

and

$$\overline{x_S} := \frac{1}{A(S)} \iint_S x dS, \quad \overline{y_S} := \frac{1}{A(S)} \iint_S y dS \quad \text{and} \quad \overline{z_S} := \frac{1}{A(S)} \iint_S z dS.$$

Let us assume that the surface S is oriented with the aid of the unit normal h directed to the exterior of D

$$h = (\cos \alpha, \cos \beta, \cos \gamma).$$

The following Hermite-Hadamard type inequalities for convex functions defined on general convex domains were obtained by B. Gavrea in 2000, [9]:

Theorem 1.2. *Let $f : D \rightarrow \mathbb{R}$ be a convex function on D , a bounded convex domain from $\mathbb{R}^3$ with a piecewise smooth boundary S . Then we have the inequalities*

$$\begin{aligned} f(\overline{x_D}, \overline{y_D}, \overline{z_D}) &\leq \frac{1}{V(D)} \iiint_D f dV \leq \frac{1}{4A(S)} \iint_S f dS \\ &+ \frac{1}{4V(D)} \iint_S [(x - \overline{x_S}) \cos \alpha + (y - \overline{y_S}) \cos \beta + (z - \overline{z_S}) \cos \gamma] f(x, y, z) dS. \end{aligned} \quad (1.2)$$

For other multivariate Hermite-Hadamard type inequalities, see [2]-[4] and [11]-[18].

Motivated by the above results, we obtain in this paper other integral inequalities of Hermite-Hadamard type for convex functions defined on convex

domains from the space. Some examples for three dimensional balls are also provided.

2. SOME PRELIMINARY FACTS AND RESULTS

Following Apostol [1], consider a surface described by the vector equation

$$r(u, v) = x(u, v) \vec{i} + y(u, v) \vec{j} + z(u, v) \vec{k} \quad (2.1)$$

where $(u, v) \in [a, b] \times [c, d]$.

If x, y, z are differentiable on $[a, b] \times [c, d]$ we consider the two vectors

$$\frac{\partial r}{\partial u} = \frac{\partial x}{\partial u} \vec{i} + \frac{\partial y}{\partial u} \vec{j} + \frac{\partial z}{\partial u} \vec{k}$$

and

$$\frac{\partial r}{\partial v} = \frac{\partial x}{\partial v} \vec{i} + \frac{\partial y}{\partial v} \vec{j} + \frac{\partial z}{\partial v} \vec{k}.$$

The *cross product* of these two vectors $\frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v}$ will be referred to as the fundamental vector product of the representation r . Its components can be expressed as *Jacobian determinants*. In fact, we have [1, p. 420]

$$\begin{aligned} \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} &= \begin{vmatrix} \frac{\partial y}{\partial u} & \frac{\partial z}{\partial u} \\ \frac{\partial y}{\partial v} & \frac{\partial z}{\partial v} \end{vmatrix} \vec{i} + \begin{vmatrix} \frac{\partial z}{\partial u} & \frac{\partial x}{\partial u} \\ \frac{\partial z}{\partial v} & \frac{\partial x}{\partial v} \end{vmatrix} \vec{j} + \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} \\ \frac{\partial x}{\partial v} & \frac{\partial y}{\partial v} \end{vmatrix} \vec{k} \quad (2.2) \\ &= \frac{\partial(y, z)}{\partial(u, v)} \vec{i} + \frac{\partial(z, x)}{\partial(u, v)} \vec{j} + \frac{\partial(x, y)}{\partial(u, v)} \vec{k}. \end{aligned}$$

Let $S = r(T)$ be a parametric surface described by a vector-valued function r defined on the box $T = [a, b] \times [c, d]$. The area of S denoted A_S is defined by the double integral [1, p. 424-425]

$$\begin{aligned} A_S &= \int_a^b \int_c^d \left\| \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right\| du dv \quad (2.3) \\ &= \int_a^b \int_c^d \sqrt{\left(\frac{\partial(y, z)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(z, x)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x, y)}{\partial(u, v)} \right)^2} du dv. \end{aligned}$$

We define surface integrals in terms of a parametric representation for the surface. One can prove that under certain general conditions the value of the integral is independent of the representation.

Let $S = r(T)$ be a parametric surface described by a vector-valued differentiable function r defined on the box $T = [a, b] \times [c, d]$ and let $f : S \rightarrow \mathbb{C}$ defined

and bounded on S . The surface integral of f over S is defined by [1, p. 430]

$$\begin{aligned} \iint_S f dS &= \int_a^b \int_c^d f(x, y, z) \left\| \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right\| dudv \\ &= \int_a^b \int_c^d f(x(u, v), y(u, v), z(u, v)) \\ &\quad \times \sqrt{\left(\frac{\partial(y, z)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(z, x)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x, y)}{\partial(u, v)} \right)^2} dudv. \end{aligned} \quad (2.4)$$

If $S = r(T)$ is a parametric surface, the fundamental vector product $N = \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v}$ is normal to S at each regular point of the surface. At each such point there are two unit normals, a unit normal n_1 , which has the same direction as N , and a unit normal n_2 which has the opposite direction. Thus

$$n_1 = \frac{N}{\|N\|} \text{ and } n_2 = -n_1.$$

Let n be one of the two normals n_1 or n_2 . Let also F be a vector field defined on S and assume that the surface integral,

$$\iint_S (F \cdot n) dS,$$

called the flux surface integral, exists. Here $F \cdot n$ is the dot or inner product.

We can write [1, p. 434]

$$\iint_S (F \cdot n) dS = \pm \int_a^b \int_c^d F(r(u, v)) \cdot \left(\frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right) dudv$$

where the sign " + " is used if $n = n_1$ and the " - " sign is used if $n = n_2$.

If

$$F(x, y, z) = P(x, y, z) \vec{i} + Q(x, y, z) \vec{j} + R(x, y, z) \vec{k}$$

and

$$r(u, v) = x(u, v) \vec{i} + y(u, v) \vec{j} + z(u, v) \vec{k} \text{ where } (u, v) \in [a, b] \times [c, d]$$

then the flux surface integral for $n = n_1$ can be explicitly calculated as [1, p. 435]

$$\begin{aligned} \iint_S (F \cdot n) dS &= \int_a^b \int_c^d P(x(u, v), y(u, v), z(u, v)) \frac{\partial(y, z)}{\partial(u, v)} dudv \\ &\quad + \int_a^b \int_c^d Q(x(u, v), y(u, v), z(u, v)) \frac{\partial(z, x)}{\partial(u, v)} dudv \\ &\quad + \int_a^b \int_c^d R(x(u, v), y(u, v), z(u, v)) \frac{\partial(x, y)}{\partial(u, v)} dudv. \end{aligned} \quad (2.5)$$

The sum of the double integrals on the right is often written more briefly as [1, p. 435]

$$\int \int_S P(x, y, z) dy \wedge dz + \int \int_S Q(x, y, z) dz \wedge dx + \int \int_S R(x, y, z) dx \wedge dy.$$

Let $B \subset \mathbb{R}^3$ be a solid in 3-space bounded by an orientable closed surface S , and let n be the unit outer normal to S . If F is a continuously differentiable vector field defined on B , we have the *Gauss-Ostrogradsky identity*

$$\iiint_B (\operatorname{div} F) dV = \int \int_S (F \cdot n) dS. \quad (\text{GO})$$

If we express

$$F(x, y, z) = P(x, y, z) \vec{i} + Q(x, y, z) \vec{j} + R(x, y, z) \vec{k},$$

then (GO) can be written as

$$\begin{aligned} & \iiint_B \left(\frac{\partial P(x, y, z)}{\partial x} + \frac{\partial Q(x, y, z)}{\partial y} + \frac{\partial R(x, y, z)}{\partial z} \right) dx dy dz \\ &= \int \int_S P(x, y, z) dy \wedge dz + \int \int_S Q(x, y, z) dz \wedge dx \\ & \quad + \int \int_S R(x, y, z) dx \wedge dy. \end{aligned} \quad (2.6)$$

By taking the real and imaginary part, we can extend the above inequality for complex valued functions P, Q, R defined on B .

For the body B we consider the coordinates for the *center of gravity* $G(\bar{x}_B, \bar{y}_B, \bar{z}_B)$ defined by

$$\bar{x}_B := \frac{1}{V(B)} \iiint_B x dx dy dz, \quad \bar{y}_B := \frac{1}{V(B)} \iiint_B y dx dy dz$$

and

$$\bar{z}_B := \frac{1}{V(B)} \iiint_B z dx dy dz.$$

We have:

Lemma 2.1. *Let B be a solid in the three dimensional space $\mathbb{R}^3$ bounded by an orientable closed surface S . If $f : B \rightarrow \mathbb{C}$ is a continuously differentiable*

function defined on a open set containing B , then we have the equality

$$\begin{aligned}
& \iiint_B f(x, y, z) dx dy dz \\
&= \frac{1}{3} \iiint_B \left[(\alpha - x) \frac{\partial f(x, y, z)}{\partial x} + (\beta - y) \frac{\partial f(x, y, z)}{\partial y} \right. \\
&\quad \left. + (\gamma - z) \frac{\partial f(x, y, z)}{\partial z} \right] dx dy dz \\
&+ \frac{1}{3} \left[\int \int_S (x - \alpha) f(x, y, z) dy \wedge dz + \int \int_S (y - \beta) f(x, y, z) dz \wedge dx \right. \\
&\quad \left. + \int \int_S (z - \gamma) f(x, y, z) dx \wedge dy \right] \quad (2.7)
\end{aligned}$$

for all α, β and γ complex numbers.

In particular, we have

$$\begin{aligned}
& \iiint_B f(x, y, z) dx dy dz \\
&= \frac{1}{3} \iiint_B \left[(\overline{x_B} - x) \frac{\partial f(x, y, z)}{\partial x} + (\overline{y_B} - y) \frac{\partial f(x, y, z)}{\partial y} \right. \\
&\quad \left. + (\overline{z_B} - z) \frac{\partial f(x, y, z)}{\partial z} \right] dx dy dz \\
&+ \frac{1}{3} \left[\int \int_S (x - \overline{x_B}) f(x, y, z) dy \wedge dz + \int \int_S (y - \overline{y_B}) f(x, y, z) dz \wedge dx \right. \\
&\quad \left. + \int \int_S (z - \overline{z_B}) f(x, y, z) dx \wedge dy \right]. \quad (2.8)
\end{aligned}$$

Proof. We have

$$\begin{aligned}
\frac{\partial [(x - \alpha) f(x, y, z)]}{\partial x} &= f(x, y, z) + (x - \alpha) \frac{\partial f(x, y, z)}{\partial x}, \\
\frac{\partial [(y - \beta) f(x, y, z)]}{\partial y} &= f(x, y, z) + (y - \beta) \frac{\partial f(x, y, z)}{\partial y}
\end{aligned}$$

and

$$\frac{\partial [(z - \gamma) f(x, y, z)]}{\partial z} = f(x, y, z) + (z - \gamma) \frac{\partial f(x, y, z)}{\partial z}.$$

By adding these three equalities we get

$$\begin{aligned}
& \frac{\partial [(x - \alpha) f(x, y, z)]}{\partial x} + \frac{\partial [(y - \beta) f(x, y, z)]}{\partial y} + \frac{\partial [(z - \gamma) f(x, y, z)]}{\partial z} \\
&= 3f(x, y, z) \\
&+ (x - \alpha) \frac{\partial f(x, y, z)}{\partial x} + (y - \beta) \frac{\partial f(x, y, z)}{\partial y} + (z - \gamma) \frac{\partial f(x, y, z)}{\partial z} \quad (2.9)
\end{aligned}$$

for all $(x, y, z) \in B$.

Integrating this equality on B we get

$$\begin{aligned}
& \iiint_B \left(\frac{\partial [(x-\alpha) f(x, y, z)]}{\partial x} + \frac{\partial [(y-\beta) f(x, y, z)]}{\partial y} \right. \\
& \quad \left. + \frac{\partial [(z-\gamma) f(x, y, z)]}{\partial z} \right) dx dy dz \\
& = 3 \iiint_B f(x, y, z) dx dy dz \\
& \quad + \iiint_B \left[(x-\alpha) \frac{\partial f(x, y, z)}{\partial x} + (y-\beta) \frac{\partial f(x, y, z)}{\partial y} \right. \\
& \quad \left. + (z-\gamma) \frac{\partial f(x, y, z)}{\partial z} \right] dx dy dz. \quad (2.10)
\end{aligned}$$

Applying the *Gauss-Ostrogradsky identity* (2.6) for the functions

$$P(x, y, z) = (x - \alpha) f(x, y, z), \quad Q(x, y, z) = (y - \beta) f(x, y, z)$$

and

$$R(x, y, z) = (z - \gamma) f(x, y, z)$$

we obtain

$$\begin{aligned}
& \iiint_B \left(\frac{\partial [(x-\alpha) f(x, y, z)]}{\partial x} + \frac{\partial [(y-\beta) f(x, y, z)]}{\partial y} \right. \\
& \quad \left. + \frac{\partial [(z-\gamma) f(x, y, z)]}{\partial z} \right) dx dy dz \\
& = \int \int_S (x - \alpha) f(x, y, z) dy \wedge dz + \int \int_S (y - \beta) f(x, y, z) dz \wedge dx \\
& \quad + \int \int_S (z - \gamma) f(x, y, z) dx \wedge dy. \quad (2.11)
\end{aligned}$$

By (2.10) and (2.11) we get

$$\begin{aligned}
& 3 \iiint_B f(x, y, z) dx dy dz \\
& + \iiint_B \left[(x - \alpha) \frac{\partial f(x, y, z)}{\partial x} + (y - \beta) \frac{\partial f(x, y, z)}{\partial y} + (z - \gamma) \frac{\partial f(x, y, z)}{\partial z} \right] dx dy dz \\
& = \int \int_S (x - \alpha) f(x, y, z) dy \wedge dz + \int \int_S (y - \beta) f(x, y, z) dz \wedge dx \\
& \quad + \int \int_S (z - \gamma) f(x, y, z) dx \wedge dy,
\end{aligned}$$

which is equivalent to the desired result (2.7). $\square$

Remark 2.2. For a function f as in Lemma 2.1 above, we define the points

$$x_{B, \partial f} := \frac{\iiint_B x \frac{\partial f(x, y, z)}{\partial x} dx dy dz}{\iiint_B \frac{\partial f(x, y, z)}{\partial x} dx dy dz}, \quad y_{B, \partial f} := \frac{\iiint_B y \frac{\partial f(x, y, z)}{\partial y} dx dy dz}{\iiint_B \frac{\partial f(x, y, z)}{\partial y} dx dy dz},$$

and

$$z_{B,\partial f} := \frac{\iiint_B z \frac{\partial f(x,y,z)}{\partial z} dx dy dz}{\iiint_B \frac{\partial f(x,y,z)}{\partial z} dx dy dz}$$

provided the denominators are not zero.

If we take $\alpha = x_{B,\partial f}$, $\beta = y_{B,\partial f}$ and $\gamma = z_{B,\partial f}$ in (2.7), then we get

$$\begin{aligned} & \iiint_B f(x,y,z) dx dy dz \\ &= \frac{1}{3} \left[\int \int_S (x - x_{B,\partial f}) f(x,y,z) dy \wedge dz + \int \int_S (y - y_{B,\partial f}) f(x,y,z) dz \wedge dx \right. \\ & \quad \left. + \int \int_S (z - z_{B,\partial f}) f(x,y,z) dx \wedge dy \right], \quad (2.12) \end{aligned}$$

since, obviously,

$$\begin{aligned} & \iiint_B \left[(x_{B,\partial f} - x) \frac{\partial f(x,y,z)}{\partial x} + (y_{B,\partial f} - y) \frac{\partial f(x,y,z)}{\partial y} \right. \\ & \quad \left. + (z_{B,\partial f} - z) \frac{\partial f(x,y,z)}{\partial z} \right] dx dy dz = 0. \end{aligned}$$

We also have the following dual approach:

Remark 2.3. For a function f as in Lemma 2.1 above, we define the points

$$x_{S,f} := \frac{\int \int_S x f(x,y,z) dy \wedge dz}{\int \int_S f(x,y,z) dy \wedge dz}, \quad y_{S,f} := \frac{\int \int_S y f(x,y,z) dz \wedge dx}{\int \int_S f(x,y,z) dz \wedge dx}$$

and

$$z_{S,f} := \frac{\int \int_S z f(x,y,z) dx \wedge dy}{\int \int_S f(x,y,z) dx \wedge dy}$$

provided the denominators are not zero.

If we take $\alpha = x_{S,f}$, $\beta = y_{S,f}$ and $\gamma = z_{S,f}$ in (2.7), then we get

$$\begin{aligned} & \iiint_B f(x,y,z) dx dy dz \\ &= \frac{1}{3} \iiint_B \left[(x_{S,f} - x) \frac{\partial f(x,y,z)}{\partial x} + (y_{S,f} - y) \frac{\partial f(x,y,z)}{\partial y} \right. \\ & \quad \left. + (z_{S,f} - z) \frac{\partial f(x,y,z)}{\partial z} \right] dx dy dz \quad (2.13) \end{aligned}$$

since, obviously,

$$\begin{aligned} & \int \int_S (x - x_{S,f}) f(x,y,z) dy \wedge dz + \int \int_S (y - y_{S,f}) f(x,y,z) dz \wedge dx \\ & \quad + \int \int_S (z - z_{S,f}) f(x,y,z) dx \wedge dy = 0. \end{aligned}$$

3. INEQUALITIES FOR CONVEX FUNCTIONS

We have the following result:

Theorem 3.1. *Let B be a convex body in the three dimensional space $\mathbb{R}^3$ bounded by an orientable closed surface S and $f : B \rightarrow \mathbb{C}$ a continuously differentiable function defined on a open set containing B . If f is convex on B , then for any $(u, v, w) \in B$ we have*

$$\begin{aligned} f(u, v, w) + (\overline{x_B} - u) \frac{\partial f(u, v, w)}{\partial x} \\ + (\overline{y_B} - v) \frac{\partial f(u, v, w)}{\partial y} + (\overline{z_B} - w) \frac{\partial f(u, v, w)}{\partial z} \\ \leq \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz \\ \leq \frac{1}{4} f(u, v, w) + \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S (x - u) f(x, y, z) dy \wedge dz \right. \\ \left. + \int \int_S (y - v) f(x, y, z) dz \wedge dx + \int \int_S (z - w) f(x, y, z) dx \wedge dy \right]. \quad (3.1) \end{aligned}$$

In particular, we have

$$\begin{aligned} f(\overline{x_B}, \overline{y_B}, \overline{z_B}) &\leq \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz \\ &\leq \frac{1}{4} f(\overline{x_B}, \overline{y_B}, \overline{z_B}) + \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S (x - \overline{x_B}) f(x, y, z) dy \wedge dz \right. \\ &\left. + \int \int_S (y - \overline{y_B}) f(x, y, z) dz \wedge dx + \int \int_S (z - \overline{z_B}) f(x, y, z) dx \wedge dy \right]. \quad (3.2) \end{aligned}$$

Proof. By the gradient inequality for the convex function f on the convex set B we have

$$\begin{aligned} (x - u) \frac{\partial f(u, v, w)}{\partial x} + (y - v) \frac{\partial f(u, v, w)}{\partial y} + (z - w) \frac{\partial f(u, v, w)}{\partial z} \\ \leq f(x, y, z) - f(u, v, w) \\ \leq (x - u) \frac{\partial f(x, y, z)}{\partial x} + (y - v) \frac{\partial f(x, y, z)}{\partial y} + (z - w) \frac{\partial f(x, y, z)}{\partial z} \end{aligned}$$

for all $(u, v, w), (x, y, z) \in B$.

If we take the integral mean over the variables $(x, y, z) \in B$, we get

$$\begin{aligned}
& \frac{1}{V(B)} \iiint_B (x-u) \frac{\partial f(u, v, w)}{\partial x} dx dy dz + \frac{1}{V(B)} \iiint_B (y-v) \frac{\partial f(u, v, w)}{\partial y} dx dy dz \\
& \quad + \frac{1}{V(B)} \iiint_B (z-w) \frac{\partial f(u, v, w)}{\partial z} dx dy dz \\
& \leq \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz - \frac{1}{V(B)} \iiint_B f(u, v, w) dx dy dz \\
& \leq \frac{1}{V(B)} \iiint_B (x-u) \frac{\partial f(x, y, z)}{\partial x} dx dy dz + \frac{1}{V(B)} \iiint_B (y-v) \frac{\partial f(x, y, z)}{\partial y} dx dy dz \\
& \quad + \frac{1}{V(B)} \iiint_B (z-w) \frac{\partial f(x, y, z)}{\partial z} dx dy dz
\end{aligned}$$

namely

$$\begin{aligned}
& (\overline{x_B} - u) \frac{\partial f(u, v, w)}{\partial x} + (\overline{y_B} - v) \frac{\partial f(u, v, w)}{\partial y} + (\overline{z_B} - w) \frac{\partial f(u, v, w)}{\partial z} \\
& \leq \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz - f(u, v, w) \\
& \leq \frac{1}{V(B)} \iiint_B (x-u) \frac{\partial f(x, y, z)}{\partial x} dx dy dz + \frac{1}{V(B)} \iiint_B (y-v) \frac{\partial f(x, y, z)}{\partial y} dx dy dz \\
& \quad + \frac{1}{V(B)} \iiint_B (z-w) \frac{\partial f(x, y, z)}{\partial z} dx dy dz \quad (3.3)
\end{aligned}$$

for all $(u, v, w) \in B$, which is an inequality of interest in itself.

The first inequality in (3.3) gives now the first part of (3.1). From the identity (2.7) we get for $(\alpha, \beta, \gamma) = (u, v, w)$ that

$$\begin{aligned}
& \iiint_B f(x, y, z) dx dy dz \\
& = \frac{1}{3} \iiint_B \left[(u-x) \frac{\partial f(x, y, z)}{\partial x} + (v-y) \frac{\partial f(x, y, z)}{\partial y} \right. \\
& \quad \left. + (w-z) \frac{\partial f(x, y, z)}{\partial z} \right] dx dy dz \\
& + \frac{1}{3} \left[\int \int_S (x-u) f(x, y, z) dy \wedge dz + \int \int_S (y-v) f(x, y, z) dz \wedge dx \right. \\
& \quad \left. + \int \int_S (z-w) f(x, y, z) dx \wedge dy \right],
\end{aligned}$$

namely

$$\begin{aligned} & \frac{1}{V(B)} \iiint_B \left[(x-u) \frac{\partial f(x,y,z)}{\partial x} + (y-v) \frac{\partial f(x,y,z)}{\partial y} \right. \\ & \quad \left. + (z-w) \frac{\partial f(x,y,z)}{\partial z} \right] dx dy dz = \\ & + \frac{1}{V(B)} \left[\int \int_S (x-u) f(x,y,z) dy \wedge dz + \int \int_S (y-v) f(x,y,z) dz \wedge dx \right. \\ & \quad \left. + \int \int_S (z-w) f(x,y,z) dx \wedge dy \right] - 3 \frac{1}{V(B)} \iiint_B f(x,y,z) dx dy dz. \end{aligned}$$

From the second part of (3.3) we get

$$\begin{aligned} & \frac{1}{V(B)} \iiint_B f(x,y,z) dx dy dz - f(u,v,w) \\ & \leq \frac{1}{V(B)} \left[\int \int_S (x-u) f(x,y,z) dy \wedge dz + \int \int_S (y-v) f(x,y,z) dz \wedge dx \right. \\ & \quad \left. + \int \int_S (z-w) f(x,y,z) dx \wedge dy \right] - 3 \frac{1}{V(B)} \iiint_B f(x,y,z) dx dy dz, \end{aligned}$$

namely

$$\begin{aligned} & \frac{4}{V(B)} \iiint_B f(x,y,z) dx dy dz \leq f(u,v,w) \\ & + \frac{1}{V(B)} \left[\int \int_S (x-u) f(x,y,z) dy \wedge dz + \int \int_S (y-v) f(x,y,z) dz \wedge dx \right. \\ & \quad \left. + \int \int_S (z-w) f(x,y,z) dx \wedge dy \right], \end{aligned}$$

which gives the second part of the inequality (3.1). $\square$

Remark 3.2. The first inequality in (3.2) is the same as the one from inequality (1.2).

Corollary 3.3. *With the assumptions of Theorem 3.1 and if $(x_{S,f}, y_{S,f}, z_{S,f}) \in B$, then*

$$\begin{aligned} & f(x_{S,f}, y_{S,f}, z_{S,f}) + (\overline{x_B} - x_{S,f}) \frac{\partial f(x_{S,f}, y_{S,f}, z_{S,f})}{\partial x} \\ & + (\overline{y_B} - y_{S,f}) \frac{\partial f(x_{S,f}, y_{S,f}, z_{S,f})}{\partial y} + (\overline{z_B} - z_{S,f}) \frac{\partial f(x_{S,f}, y_{S,f}, z_{S,f})}{\partial z} \\ & \leq \frac{1}{V(B)} \iiint_B f(x,y,z) dx dy dz \leq \frac{1}{4} f(x_{S,f}, y_{S,f}, z_{S,f}). \quad (3.4) \end{aligned}$$

The proof follows by (3.1) observing that

$$\begin{aligned} \int \int_S (x - x_{S,f}) f(x, y, z) dy \wedge dz + \int \int_S (y - y_{S,f}) f(x, y, z) dz \wedge dx \\ + \int \int_S (z - z_{S,f}) f(x, y, z) dx \wedge dy = 0. \end{aligned}$$

Corollary 3.4. *With the assumptions of Theorem 3.1, we have*

$$\begin{aligned} \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz \leq \frac{1}{3} \frac{1}{V(B)} \left[\int \int_S (x - \overline{x_B}) f(x, y, z) dy \wedge dz \right. \\ \left. + \int \int_S (y - \overline{y_B}) f(x, y, z) dz \wedge dx + \int \int_S (z - \overline{z_B}) f(x, y, z) dx \wedge dy \right]. \quad (3.5) \end{aligned}$$

Proof. From (3.2) we get

$$\begin{aligned} \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz \\ \leq \frac{1}{4} f(\overline{x_B}, \overline{y_B}, \overline{z_B}) + \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S (x - \overline{x_B}) f(x, y, z) dy \wedge dz \right. \\ \left. + \int \int_S (y - \overline{y_B}) f(x, y, z) dz \wedge dx + \int \int_S (z - \overline{z_B}) f(x, y, z) dx \wedge dy \right] \\ \leq \frac{1}{4} \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz + \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S (x - \overline{x_B}) f(x, y, z) dy \wedge dz \right. \\ \left. + \int \int_S (y - \overline{y_B}) f(x, y, z) dz \wedge dx + \int \int_S (z - \overline{z_B}) f(x, y, z) dx \wedge dy \right], \end{aligned}$$

which implies that

$$\begin{aligned} \frac{3}{4V(B)} \iiint_B f(x, y, z) dx dy dz \leq \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S (x - \overline{x_B}) f(x, y, z) dy \wedge dz \right. \\ \left. + \int \int_S (y - \overline{y_B}) f(x, y, z) dz \wedge dx + \int \int_S (z - \overline{z_B}) f(x, y, z) dx \wedge dy \right] \end{aligned}$$

that is equivalent to (3.5). $\square$

Corollary 3.5. *With the assumptions of Theorem 3.1 we have*

$$\begin{aligned} & \frac{1}{A(S)} \int \int_S f(u, v, w) dS + \frac{1}{A(S)} \int \int_S \left[(\overline{x_B} - u) \frac{\partial f(u, v, w)}{\partial x} \right. \\ & \quad \left. + (\overline{y_B} - v) \frac{\partial f(u, v, w)}{\partial y} + (\overline{z_B} - w) \frac{\partial f(u, v, w)}{\partial z} \right] dS \\ & \leq \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz \\ & \leq \frac{1}{4} \frac{1}{A(S)} \int \int_S f(u, v, w) dS + \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S (x - \overline{x_S}) f(x, y, z) dy \wedge dz \right. \\ & \quad \left. + \int \int_S (y - \overline{y_S}) f(x, y, z) dz \wedge dx + \int \int_S (z - \overline{z_S}) f(x, y, z) dx \wedge dy \right], \quad (3.6) \end{aligned}$$

where

$$\overline{x_S} = \frac{1}{A(S)} \int \int_S u dS, \quad \overline{y_S} = \frac{1}{A(S)} \int \int_S v dS, \quad \overline{z_S} = \frac{1}{A(S)} \int \int_S w dS.$$

Proof. By taking the integral mean $\frac{1}{A(S)} \int \int_S (\cdot) dS$ over the variables (u, v, w) in the integral (3.1), we get

$$\begin{aligned} & \frac{1}{A(S)} \int \int_S f(u, v, w) dS + \frac{1}{A(S)} \int \int_S \left[(\overline{x_B} - u) \frac{\partial f(u, v, w)}{\partial x} \right. \\ & \quad \left. + (\overline{y_B} - v) \frac{\partial f(u, v, w)}{\partial y} + (\overline{z_B} - w) \frac{\partial f(u, v, w)}{\partial z} \right] dS \\ & \leq \frac{1}{V(B)} \iiint_B f(x, y, z) dx dy dz \\ & \leq \frac{1}{4} \frac{1}{A(S)} \int \int_S f(u, v, w) dS \\ & \quad + \frac{1}{4} \frac{1}{V(B)} \left[\int \int_S \left(x - \frac{1}{A(S)} \int \int_S u dS \right) f(x, y, z) dy \wedge dz \right. \\ & \quad \left. + \int \int_S \left(y - \frac{1}{A(S)} \int \int_S v dS \right) f(x, y, z) dz \wedge dx \right. \\ & \quad \left. + \int \int_S \left(z - \frac{1}{A(S)} \int \int_S w dS \right) f(x, y, z) dx \wedge dy \right], \end{aligned}$$

which is equivalent to (3.6). $\square$

Remark 3.6. The second inequality in (3.6) is an equivalent formulation of the second inequality in (1.2).

4. APPLICATIONS FOR THREE DIMENSIONAL BALLS

Now, let us compute the surface integral

$$K(S(C, R), f) := \iint_{S(C, R)} f(x, y, z) dS,$$

where

$$S(C, R) := \left\{ (x, y, z) \in \mathbb{R}^3 \mid (x - a)^2 + (y - b)^2 + (z - c)^2 = R^2 \right\}.$$

If we consider the parametrization of $S(C, R)$ given by:

$$S(C, R) : \begin{cases} x = R \cos \psi \cos \varphi + a \\ y = R \cos \psi \sin \varphi + b \\ z = R \sin \psi + c \end{cases} ; (\psi, \varphi) \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \times [0, 2\pi]$$

and putting

$$A := \begin{vmatrix} \frac{\partial y}{\partial \psi} & \frac{\partial z}{\partial \psi} \\ \frac{\partial y}{\partial \varphi} & \frac{\partial z}{\partial \varphi} \end{vmatrix} = -R^2 \cos^2 \psi \cos \varphi,$$

$$B := \begin{vmatrix} \frac{\partial x}{\partial \psi} & \frac{\partial z}{\partial \psi} \\ \frac{\partial x}{\partial \varphi} & \frac{\partial z}{\partial \varphi} \end{vmatrix} = R^2 \cos^2 \psi \sin \varphi,$$

and

$$C := \begin{vmatrix} \frac{\partial x}{\partial \psi} & \frac{\partial y}{\partial \psi} \\ \frac{\partial x}{\partial \varphi} & \frac{\partial y}{\partial \varphi} \end{vmatrix} = -R^2 \sin \psi \cos \psi,$$

we have that

$$A^2 + B^2 + C^2 = R^4 \cos^2 \psi \text{ for all } (\psi, \varphi) \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \times [0, 2\pi].$$

Thus,

$$\begin{aligned} K(S(C, R), f) &= \iint_{S(C, R)} f(x, y, z) dS \\ &= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} [f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c) \\ &\quad \times \sqrt{A^2 + B^2 + C^2}] d\psi d\varphi \\ &= R^2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} \cos \psi f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c) d\psi d\varphi. \end{aligned} \tag{4.1}$$

We also have

$$\begin{aligned}
L(S(C, R), f) &:= \int \int_{S(C, R)} (x - a) f(x, y, z) dy \wedge dz \\
&+ \int \int_{S(C, R)} (y - b) f(x, y, z) dz \wedge dx + \int \int_{S(C, R)} (z - c) f(x, y, z) dx \wedge dy \\
&= -R^3 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} \cos^3 \psi \cos^2 \varphi \\
&\quad \times f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c) d\psi d\varphi \\
&\quad + R^3 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} \cos^3 \psi \sin^2 \varphi \\
&\quad \times f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c) d\psi d\varphi \\
&- R^3 \int \int_S \sin^2 \psi \cos \psi f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c) d\psi d\varphi.
\end{aligned} \tag{4.2}$$

Let us consider the transformation $T_2 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by:

$$T_2(r, \psi, \varphi) := (r \cos \psi \cos \varphi + a, r \cos \psi \sin \varphi + b, r \sin \psi + c).$$

It is well known that the Jacobian of T_2 is

$$J(T_2) = r^2 \cos \psi$$

and T_2 is a one-to-one mapping defined on the interval of $\mathbb{R}^3$, $[0, R] \times [-\frac{\pi}{2}, \frac{\pi}{2}] \times [0, 2\pi]$, with values in the ball $B(C, R)$ from $\mathbb{R}^3$. Thus we have the change of variable:

$$\begin{aligned}
I(B(C, R), f) &:= \iiint_{B(C, R)} f(x, y, z) dx dy dz \\
&= \int_0^R \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} f(r \cos \psi \cos \varphi + a, r \cos \psi \sin \varphi + b, r \sin \psi + c) r^2 \cos \psi dr d\psi d\varphi.
\end{aligned} \tag{4.3}$$

Assume that f is convex on the ball $B(C, R)$. From the inequality (3.2) we get

$$\begin{aligned}
f(a, b, c) &\leq \frac{1}{V(B(C, R))} I(B(C, R), f) \\
&\leq \frac{1}{4} f(a, b, c) + \frac{1}{4V(B(C, R))} L(S(C, R), f), \tag{4.4}
\end{aligned}$$

where $V(B(C, R)) = \frac{4\pi R^3}{3}$, while from the inequality (3.5) we also have

$$\frac{1}{V(B(C, R))} I(B(C, R), f) \leq \frac{1}{3V(B(C, R))} L(S(C, R), f). \tag{4.5}$$

Further, consider

$$\begin{aligned}
 J(S(C, R), f) &:= \int \int_{S(C, R)} \left[(\overline{x_B} - u) \frac{\partial f(u, v, w)}{\partial x} \right. \\
 &\quad \left. + (\overline{y_B} - v) \frac{\partial f(u, v, w)}{\partial y} + (\overline{z_B} - w) \frac{\partial f(u, v, w)}{\partial z} \right] dS \\
 &= -R \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} \cos \psi \cos \varphi \frac{\partial f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c)}{\partial x} d\psi d\varphi \\
 &\quad - R \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} \cos \psi \sin \varphi \frac{\partial f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c)}{\partial y} d\psi d\varphi \\
 &\quad - R \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2\pi} \sin \psi \frac{\partial f(R \cos \psi \cos \varphi + a, R \cos \psi \sin \varphi + b, R \sin \psi + c)}{\partial z} d\psi d\varphi.
 \end{aligned}$$

Then from the inequality (3.6) we get the following inequalities of interest:

$$\begin{aligned}
 &\frac{1}{A(S(C, R))} K(S(C, R), f) + \frac{1}{A(S(C, R))} J(S(C, R), f) \\
 &\leq \frac{1}{V(B(C, R))} I(B(C, R), f) \\
 &\leq \frac{1}{4} \frac{1}{A(S(C, R))} K(S(C, R), f) + \frac{1}{4} \frac{1}{V(B(C, R))} L(S(C, R), f), \quad (4.6)
 \end{aligned}$$

where $A(S(C, R)) = 4\pi R^2$ is the area of the sphere.

ACKNOWLEDGMENTS

The authors would like to thank the reviewers for their thoughtful comments and efforts towards improving our manuscript.

REFERENCES

1. T. M. Apostol, *Calculus Volume II, Multi Variable Calculus and Linear Algebra, with Applications to Differential Equations and Probability*, Second Edition, John Wiley & Sons, New York, London, Sydney, Toronto, 1969.
2. A. Barani, Hermite–Hadamard and Ostrowski Type Inequalities on Hemispheres, *Mediterr. J. Math.*, **13**, (2016), 4253–4263.
3. M. Bessenyei, The Hermite–Hadamard Inequality on Simplices, *Amer. Math. Monthly*, **115**, (2008), 339–345.
4. J. de la Cal, J. Cárcamo, Multidimensional Hermite–Hadamard Inequalities and the Convex Order, *Journal of Mathematical Analysis and Applications*, **324**(1), (2006), 248–261.
5. S. S. Dragomir, On Hadamard’s Inequality on a Disk, *Journal of Ineq. Pure & Appl. Math.*, **1**(1), (2000), Article 2. [Online <https://www.emis.de/journals/JIPAM/article95.html?sid=95>].
6. S. S. Dragomir, On Hadamard’s Inequality for the Convex Mappings Defined on a Ball in the Space and Applications, *Math. Ineq. & Appl.*, **3** (2), (2000), 177–187.
7. S. S. Dragomir, Double Integral Inequalities of Hermite–Hadamard Type for h -convex Functions on Linear Spaces, *Analysis* (Berlin), **37**(1), (2017), 13–22.
8. S. S. Dragomir, Hermite–Hadamard Type Integral Inequalities for Double Integral on General Domains, Preprint *RGMA Res. Rep. Coll.*, **22**, (2019), Art. 46, 10 pp. [Online <http://rgmia.org/papers/v22/v22a46.pdf>].

9. B. Gavrea, On Hadamard's Inequality for the Convex Mappings Defined a Convex Domain in the Space, *Journal of Inequalities in Pure and Applied Mathematics*, **1**(1), (2000). [<https://www.emis.de/journals/JIPAM/article102.html?sid=102>].
10. P. O. Mohammed, M Vivas-Cortez, T Abdeljawad, Y Rangel-Oliveros, Integral Inequalities of Hermite-Hadamard Type for Quasi-convex Functions with Applications, *AIMS Math*, **5**, (2020), 7316-7331.
11. M. Matłoka, On Hadamard's Inequality for h -convex Function on a Disk, *Applied Mathematics and Computation*, **235**, (2014), 118-123.
12. F.-C. Mitroi, E. Symeonidis, The Converse of the Hermite-Hadamard Inequality on Simplices, *Expo. Math.*, **30**, (2012), 389-396.
13. E. Neuman, Inequalities Involving Multivariate Convex Functions II, *Proc. Amer. Math. Soc.*, **109**, (1990), 965-974.
14. E. Neuman, J. Pečarić, Inequalities Involving Multivariate Convex Functions, *J. Math. Anal. Appl.*, **137**, (1989), 541-549.
15. M. Vivas -Cortez, J. E. H. Hernández, L. A. Azócar, Some New Generalized Jensen and Hermite-Hadamard Inequalities for Operator h -convex Functions, *Appl. Math. Inf. Sci*, **11**(2), (2017), 383-392.
16. S. Wasowicz, A. Witkowski, On Some Inequality of Hermite-Hadamard Type, *Opusc. Math.*, **32**(3), (2012), 591-600.
17. F.-L. Wang, The Generalizations to Several-dimensions of the Classical Hadamard's Inequality, *Mathematics in Practice and Theory*, **36**(9), 370-373, 2006 (Chinese).
18. F.-L. Wang, A Family of Mappings Associated with Hadamard's Inequality on a Hypercube, *International Scholarly Research Network ISRN Mathematical Analysis*, **2011**, Article ID 594758, 9 pages doi:10.5402/2011/594758.