

Approximation by (p, q) -Lupaş Stancu Operators

Asif Khan, Vinita Sharma*

Department of Mathematics, Aligarh Muslim University, Aligarh–202002,
India.

E-mail: asifjnu07@gmail.com

E-mail: vinita.sha23@gmail.com

ABSTRACT. In this paper, (p, q) -Lupaş Bernstein Stancu operators are constructed. Statistical as well as other approximation properties of (p, q) -Lupaş Stancu operators are studied. Rate of statistical convergence by means of modulus of continuity and Lipschitz type maximal functions has been investigated.

Keywords: (p, q) -Integers, Lupaş (p, q) -Bernstein Stancu operators, Statistical approximation, Korovkin's type approximation.

2000 Mathematics subject classification: 65D17, 41A10, 41A25, 41A36.

1. INTRODUCTION AND PRELIMINARIES

In 1912, S.N. Bernstein [6] introduced his famous operators $B_n : C[0, 1] \rightarrow C[0, 1]$ defined for any $n \in \mathbb{N}$ and for any function $f \in C[0, 1]$

$$B_n(f; x) = \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} f\left(\frac{k}{n}\right), \quad x \in [0, 1]. \quad (1.1)$$

and named it Bernstein polynomials to provide a constructive proof of the Weierstrass theorem [20].

*Corresponding Authors

Further, based on q -integers, Lupaş [21] introduced the first q -Bernstein operators [6] and investigated its approximating and shape-preserving properties. Another q -analogue of the Bernstein polynomials is due to Phillips [38]. Since then several generalizations of well-known positive linear operators based on q -integers have been introduced and their approximation properties studied.

Recently, the applications of (p, q) -calculus (post quantum calculus) emerged as a new area in the field of approximation theory [20]. The development of post quantum calculus has led to the discovery of various generalizations of Bernstein polynomials involving (p, q) -integers. The aim of these generalizations is to provide appropriate and powerful tools to application areas such as numerical analysis, computer-aided geometric design [7] and solutions of differential equations.

Mursaleen *et al* [27] introduced the concept of post quantum calculus in approximation theory and constructed the (p, q) -analogue of Bernstein operators defined as follows for $0 < q < p \leq 1$:

$$B_{n,p,q}(f; x) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) f\left(\frac{[k]_{p,q}}{p^{k-n}[n]_{p,q}}\right), \quad x \in [0, 1]. \quad (1.2)$$

Note when $p = 1$, (p, q) -Bernstein Operators given by (1.2) turns out to be Phillips q -Bernstein Operators [38].

Also, we have

$$\begin{aligned} (1-x)_{p,q}^n &= \prod_{s=0}^{n-1} (p^s - q^s x) = (1-x)(p-qx)(p^2 - q^2 x) \dots (p^{n-1} - q^{n-1} x) \\ &= \sum_{k=0}^n (-1)^k p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \left[\begin{matrix} n \\ k \end{matrix} \right]_{p,q} x^k. \end{aligned}$$

Further, they applied the concept of (p, q) -calculus in approximation theory and studied approximation properties based on (p, q) -integers for Bernstein-Stancu operators, (p, q) -analogue of Bernstein-Kantorovich, (p, q) -analogue of Bernstein-Shurer operators, (p, q) -analogue of Bleimann-Butzer-Hahn operators and (p, q) -analogue of Lorentz polynomials on a compact disk in [28, 31, 32, 33, 35].

On the other hand, Khalid and Lobiyal defined (p, q) -analogue of Lupaş Bernstein operators [17] as follows :

For any $p > 0$ and $q > 0$, the linear operators $L_{p,q}^n : C[0, 1] \rightarrow C[0, 1]$ as

$$L_{p,q}^n(f; x) = \sum_{k=0}^n \frac{f\left(\frac{p^{n-k} [k]_{p,q}}{[n]_{p,q}}\right) \left[\begin{matrix} n \\ k \end{matrix} \right]_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} x^k (1-x)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-x) + q^{j-1}x\}}, \tag{1.3}$$

are (p, q) -analogue of Lupaş Bernstein operators.

Again when $p = 1$, Lupaş (p, q) -Bernstein operators turns out to be Lupaş q -Bernstein operators as given in [22, 37].

When $p = q = 1$, Lupaş (p, q) -Bernstein operators turns out to be classical Bernstein operators [6].

They studied two different techniques as de-Casteljau’s algorithm and Korovkin’s type approximation properties [17]: de-Casteljau’s algorithm and related results of degree elevation reduction for Bèzier curves and surfaces holds for all $p > 0$ and $q > 0$. However to study Korovkin’s type approximation properties for Lupaş (p, q) -analogue of the Bernstein operators, $0 < q < p \leq 1$ is needed.

Based on Korovkin’s type approximation, they proved that the sequence of (p, q) -analogue of Lupaş Bernstein operators $L_{p_n, q_n}^n(f, x)$ converges uniformly to $f(x) \in C[0, 1]$ if and only if $0 < q_n < p_n \leq 1$ such that $\lim_{n \rightarrow \infty} q_n = 1$, $\lim_{n \rightarrow \infty} p_n = 1$ and $\lim_{n \rightarrow \infty} q_n^n = 1$. On the other hand, for any $p > 0$ fixed and $p \neq 1$, the sequence $L_{p,q}^n(f, x)$ converges uniformly to $f(x) \in C[0, 1]$ if and only if $f(x) = ax + b$ for some $a, b \in \mathbb{R}$.

Furthermore, in comparison to q -Bèzier curves and surfaces based on Lupaş q -Bernstein rational functions, their generalization gives more flexibility in controlling the shapes of curves and surfaces.

Some advantages of using the extra parameter p have been discussed in the field of approximations on compact disk [35] and in computer aided geometric design [17].

For more details related to approximation theory [20], one can refer [1, 2, 3, 5, 8, 9, 12, 13, 14, 15, 18, 19, 22, 23, 24, 34, 36, 39, 40, 42, 43, 44, 45, 46, 47, 48].

Let us recall certain notations of (p, q) -calculus.

For any $p > 0$ and $q > 0$, the (p, q) integers $[n]_{p,q}$ are defined by

$$[n]_{p,q} = p^{n-1} + p^{n-2}q + p^{n-3}q^2 + \dots + pq^{n-2} + q^{n-1} = \begin{cases} \frac{p^n - q^n}{p - q}, & \text{when } p \neq q \neq 1 \\ n p^{n-1}, & \text{when } p = q \neq 1 \\ [n]_q, & \text{when } p = 1 \\ n, & \text{when } p = q = 1 \end{cases}$$

where $[n]_q$ denotes the q -integers and $n = 0, 1, 2, \dots$.

The formula for (p, q) -binomial expansion is as follows:

$$(ax + by)_{p,q}^n := \sum_{k=0}^n p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} a^{n-k} b^k x^{n-k} y^k,$$

$$(x + y)_{p,q}^n = (x + y)(px + qy)(p^2x + q^2y) \cdots (p^{n-1}x + q^{n-1}y),$$

$$(1 - x)_{p,q}^n = (1 - x)(p - qx)(p^2 - q^2x) \cdots (p^{n-1} - q^{n-1}x),$$

where (p, q) -binomial coefficients are defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = \frac{[n]_{p,q}!}{[k]_{p,q}! [n-k]_{p,q}!}.$$

Details on (p, q) -calculus can be found in [10, 11, 27].

Also, we have (p, q) -analogue of Euler's identity as:

$$(1 - x)_{p,q}^n = \prod_{s=0}^{n-1} (p^s - q^s x) = (1 - x)(p - qx)(p^2 - q^2x) \cdots (p^{n-1} - q^{n-1}x)$$

$$= \sum_{k=0}^n (-1)^k p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} x^k.$$

Again by some simple calculations and using the property of (p, q) -integers, we get (p, q) -analogue of Pascal's relation as follows:

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = q^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} + p^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} \quad (1.4)$$

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = p^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} + q^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q}. \quad (1.5)$$

We recall some results from [17] for Lupas (p, q) -Bernstein operators, which reproduces linear and constant functions.

Some auxillary results:

$$(1) L_{p,q}^n(1, \frac{u}{u+1}) = 1$$

$$(2) L_{p,q}^n(t, \frac{u}{u+1}) = \frac{u}{u+1}$$

$$(3) L_{p,q}^n(t^2, \frac{u}{u+1}) = \frac{u}{u+1} \frac{p^{n-1}}{[n]_{p,q}} + \frac{qu}{u+1} (\frac{qu}{p+qu}) \frac{[n-1]_{p,q}}{[n]_{p,q}}$$

or equivalently for $x = \frac{u}{u+1}$

$$L_{p,q}^n(1, x) = 1, \tag{1.6}$$

$$L_{p,q}^n(t, x) = x, \tag{1.7}$$

$$L_{p,q}^n(t^2, x) = \frac{p^{n-1}x}{[n]_{p,q}} + \frac{q^2 x^2}{p(1-x) + qx} \frac{[n-1]_{p,q}}{[n]_{p,q}}. \tag{1.8}$$

2. CONSTRUCTION OF (p, q) -LUPAŞ STANCU OPERATORS

In this section, we introduce (p, q) -Lupaş Stancu operators as follows:

For any $p > 0$ and $q > 0$, the linear operators $L_{p,q}^n : C[0, 1] \rightarrow C[0, 1]$

$$L_{n,p,q}^{\alpha,\beta}(f; x) = \sum_{k=0}^n f\left(\frac{p^{n-k} [k]_{p,q} + \alpha}{[n]_{p,q} + \beta}\right) b_{p,q}^{k,n}(t) \tag{2.1}$$

and $b_{p,q}^{k,n}(t)$ is given by

$$b_{p,q}^{k,n}(t) = \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} t^k (1-t)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-t) + q^{j-1}t\}}, \tag{2.2}$$

where $0 < \alpha < \beta$.

We give some equalities for operators (2.1) in the following lemma.

Lemma 4.1. The following equalities are true:

- (i) $L_{n,p,q}^{\alpha,\beta}(1; x) = 1,$
- (ii) $L_{n,p,q}^{\alpha,\beta}(t; x) = \frac{[n]_{p,q}x + \alpha}{[n]_{p,q} + \beta},$
- (iii) $L_{n,p,q}^{\alpha,\beta}(t^2; x) = \frac{1}{([n]_{p,q} + \beta)^2} \frac{q^2 [n]_{p,q} [n-1]_{p,q}}{p(1-x) + qx} x^2 + \frac{[n]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q} + \beta)^2} x + \frac{\alpha^2}{([n]_{p,q} + \beta)^2}.$

Proof. Proof of part (i) is obvious.

$$\begin{aligned} L_{n,p,q}^{\alpha,\beta}(t; x) &= \sum_{k=0}^n \left(\frac{p^{n-k} [k]_{p,q} + \alpha}{[n]_{p,q} + \beta} \right) b_{p,q}^{k,n}(t) \\ &= \frac{[n]_{p,q}}{[n]_{p,q} + [\beta]} L_{p,q}^n(t; x) + \frac{[\alpha]}{[n]_{p,q} + [\beta]} L_{p,q}^n(1; x). \end{aligned}$$

So from inequalities (1.6) and (1.7), we get the result.

Proof (iii)

$$\begin{aligned} L_{n,p,q}^{\alpha,\beta}(t^2; x) &= \sum_{k=0}^n \left(\frac{p^{n-k} [k]_{p,q} + \alpha}{[n]_{p,q} + \beta} \right) b_{p,q}^{k,n}(t) \\ &= \frac{1}{([n]_{p,q} + \beta)^2} \left[p^{2n-2k} [k]_{p,q}^2 b_{p,q}^{k,n}(t) \right. \\ &\quad \left. + 2\alpha p^{n-k} [k]_{p,q} b_{p,q}^{k,n}(t) + \alpha^2 b_{p,q}^{k,n}(t) \right] \\ &= \frac{1}{([n]_{p,q} + \beta)^2} [A + B + C]. \end{aligned}$$

$$A = p^{2n} \sum_{k=0}^n \frac{[k]_{p,q}^2}{p^{2k}} \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} t^k (1-t)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-t) + q^{j-1}t\}}$$

$$A = [n]p^{2n} \sum_{k=1}^n \frac{[k]_{p,q}}{p^{2k}} \frac{\begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} u^k}{\prod_{j=1}^n \{p^{j-1} + q^{j-1}u\}}.$$

On shifting the limits and on replacing k by $k+1$, we get

$$\begin{aligned} A &= [n]p^{2n} \sum_{k=1}^n \frac{[k+1]_{p,q}}{p^{2k+2}} \frac{\begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k-1)(n-k-2)}{2}} q^{\frac{k(k+1)}{2}} u^k}{\prod_{j=1}^{n-1} \{p^j + q^j u\}}, \\ &= [n]p^n \frac{u}{u+1} \sum_{k=0}^{n-1} \frac{[k+1]_{p,q}}{p^{k+2}} \frac{\begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k-1)(n-k-2)}{2}} q^{\frac{k(k+1)}{2}} \left(\frac{qu}{p}\right)^k}{\prod_{j=0}^{n-2} \{p^j + q^j \left(\frac{qu}{p}\right)\}}. \end{aligned}$$

Using $[k + 1]_{p,q} = p^k + q[k]_{p,q}$, we get our desired result:

$$\begin{aligned}
 A &= [n]p^n \frac{u}{u+1} \sum_{k=0}^{n-1} \frac{[p^k + q[k]]}{p^{k+2}} \frac{\begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k-1)(n-k-2)}{2}} q^{\frac{k(k+1)}{2}} \left(\frac{qu}{p}\right)^k}{\prod_{j=0}^{n-2} \{p^j + q^j(\frac{qu}{p})\}}, \\
 &= [n]_{p,q} p^{n-1} \frac{u}{u+1} + \frac{q^2 u^2 [n]_{p,q} [n-1]_{p,q}}{(u+1)(p+qu)},
 \end{aligned}$$

equivalently

$$A = [n]_{p,q} p^{n-1} x + \frac{q^2 [n]_{p,q} [n-1]_{p,q}}{(p(1-x) + qx)} x^2.$$

Similarly

$$\begin{aligned}
 B &= 2\alpha p^n \sum_{k=0}^n \frac{[k]}{p^k} \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} t^k (1-t)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-t) + q^{j-1}t\}}, \\
 &= 2\alpha [n]p^n \sum_{k=1}^n \frac{1}{p^k} \frac{\begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} u^k}{\prod_{j=1}^n \{p^{j-1} + q^{j-1}u\}}.
 \end{aligned}$$

After shifting the limits and on replacing k by $k + 1$, we get

$$B = 2\alpha [n]p^n \frac{u}{u+1} \sum_{k=0}^{n-1} \frac{1}{p} \frac{\begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k-1)(n-k-2)}{2}} q^{\frac{k(k-1)}{2}} \left(\frac{qu}{p}\right)^k}{\prod_{j=0}^{n-2} \{p^j + q^j(\frac{qu}{p})\}},$$

which implies

$$B = 2\alpha [n]_{p,q} x.$$

Similarly

$$\begin{aligned}
 C &= \alpha^2 \sum_{k=0}^n \frac{\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} t^k (1-t)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-t) + q^{j-1}t\}}, \\
 &= \alpha^2.
 \end{aligned}$$

□

Theorem 2.1. Let $0 < q_n < p_n \leq 1$ such that $\lim_{n \rightarrow \infty} p_n = 1$, $\lim_{n \rightarrow \infty} q_n = 1$, $\lim_{n \rightarrow \infty} q_n^n = 1$ and for $f \in C[0, 1]$, we have $\lim_n |L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| = 0$.

Proof. Let us recall the following Korovkin's theorem see [20]. Let (T_n) be a sequence of positive linear operators from $C[a, b]$ into $C[a, b]$. Then $\lim \|T_n(f, x) - f(x)\|_{C[a,b]} = 0$, for all $f \in C[a, b]$ if and only if $\lim_n \|T_n(f_i, x) - f_i(x)\|_{C[a,b]} = 0$, for $i = 0, 1, 2$, where $f_0(t) = 1$, $f_1(t) = t$ and $f_2(t) = t^2$.

□

3. THE RATE OF CONVERGENCE

In this section, we compute the rates of convergence of the operators $L_{n,p,q}^{\alpha,\beta}(f; x)$ to the functions f by means of modulus of continuity, elements of Lipschitz class and peetre's K-functional.

Let $f \in C[0, 1]$. The modulus of continuity of f denoted by $\omega(f, \delta)$ is defined as:

$$\omega(f, \delta) = \sup_{y,x \in [0,1], |y-x| < \delta} |f(y) - f(x)|.$$

where $w(f; \delta)$ satisfies the following conditions: for all $f \in C[0, 1]$,

$$\lim_{\delta \rightarrow 0} w(f; \delta) = 0. \quad (3.1)$$

and

$$|f(y) - f(x)| \leq w(f; \delta) \left(\frac{|y-x|}{\delta} + 1 \right). \quad (3.2)$$

Theorem 3.1. Let $0 < q < p \leq 1$, and $f \in C[0, 1]$, and $\delta > 0$, we have

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq 2\omega(f; \delta_n)$$

where

$$\delta_n = \left[\left(\frac{q^2 [n]_{p,q} [n-1]_{p,q}}{([n]_{p,q} + \beta)^2 (p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) + \left(\frac{p^{n-1} [n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right]^{\frac{1}{2}}.$$

Proof. From lemma (4.1) we have

$$|L_{n,p,q}^{\alpha,\beta}(t-x)^2; x| = \left(\frac{q^2[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q} + \beta)^2(p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) x^2 + \left(\frac{p^{n-1}[n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) x + \frac{\alpha^2}{([n]_{p,q} + \beta)^2}. \tag{3.3}$$

For $x \in [0, 1]$, we take

$$|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq w(f; \delta) \left\{ 1 + \frac{1}{\delta} (L_{n,p,q}^{\alpha,\beta}(t-x)^2 : x)^{\frac{1}{2}} \right\},$$

then we get

$$\begin{aligned} \|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} &\leq w(f; \delta) \left\{ 1 + \frac{1}{\delta} (L_{n,p,q}^{\alpha,\beta}(t-x)^2 : x)^{\frac{1}{2}} \right\} \\ &\leq w(f; \delta) \left\{ 1 + \frac{1}{\delta} \left(\frac{1}{([n]_{p,q} + \beta)^2} \frac{q^2[n]_{p,q}[n-1]_{p,q}}{p(1-x) + qx} \right. \right. \\ &\quad \left. \left. - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) + \left(\frac{p^{n-1}[n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right. \right. \\ &\quad \left. \left. + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right)^{\frac{1}{2}} \right\}. \end{aligned}$$

If we choose

$$\begin{aligned} \delta_n &= \left[\left(\frac{q^2[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q} + \beta)^2(p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) \right. \\ &\quad \left. + \left(\frac{p^{n-1}[n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right]^{\frac{1}{2}}. \end{aligned}$$

Then we have

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq 2\omega(f; \delta_n).$$

So we have the desired result. □

Now we compute the approximation order of operator $L_{n,p,q}^{\alpha,\beta}$ in term of the elements of the usual Lipschitz class.

Let $f \in C[0, 1]$ and $0 < \rho \leq 1$. We recall that f belongs to $Lip_M(\rho)$ if the inequality

$$|f(x) - f(y)| \leq M|x - y|^\rho; \text{ for all } x, y \in [0, 1] \quad (3.4)$$

holds.

Theorem 3.2. For all $f \in Lip_M(\rho)$

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq M\delta_n^\rho$$

where

$$\delta_n = \left[\left(\frac{q^2[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q} + \beta)^2(p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) + \left(\frac{p^{n-1}[n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right]^{\frac{1}{2}}$$

and M is a positive constant.

Proof. Let $f \in Lip_M(\rho)$ and $0 < \rho \leq 1$. by (3.4) and linearity and monotonicity of $L_{n,p,q}^{\alpha,\beta}$ then we have

$$\begin{aligned} |L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| &\leq L_{n,p,q}^{\alpha,\beta}(|f(t) - f(x)|; x) \\ &\leq L_{n,p,q}^{\alpha,\beta}(|t - x|^\rho; x). \end{aligned}$$

Applying the Holder inequality with $m = \frac{2}{\rho}$ and $n = \frac{2}{2-\rho}$, we get

$$|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq (L_{n,p,q}^{\alpha,\beta}((t-x)^2; x))^{\frac{\rho}{2}}. \quad (3.5)$$

if we choose $\delta = \delta_n$ as above, then proof is completed.

Finally, we will study the rate of convergence of the positive linear operators $L_{n,p,q}^{\alpha,\beta}$ by means of the Peetre's K-functionals.

$C^2[0, 1]$: The space of those functions f for which $f, f', f'' \in C[0, 1]$. we recall the following norm in the space $C^2[0, 1]$:

$$\|f\|_{C^2[0,1]} = \|f\|_{C[0,1]} + \|f'\|_{C[0,1]} + \|f''\|_{C[0,1]}.$$

We consider the following Peetre's K-functional

$$K(f, \delta) := \inf_{g \in C^2[0,1]} \left\{ \|f - g\|_{C[0,1]} + \delta \|g\|_{C^2[0,1]} \right\}.$$

□

Theorem 3.3. *Let $f \in C[0, 1]$. Then we have*

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq 2K(f; \delta_n)$$

Where $K(f; \delta_n)$ is Peetre's functional and

$$\begin{aligned} \delta_n = & \frac{1}{4} \left(\frac{q^2 [n]_{p,q} [n-1]_{p,q}}{([n]_{p,q} + \beta)^2 (p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) \\ & + \frac{1}{4} \left(\frac{p^{n-1} [n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) + \frac{1}{4} \frac{\alpha^2}{([n]_{p,q} + \beta)^2} + \frac{1}{2} \frac{[\alpha] + [\beta]}{[n]_{p,q} + [\beta]}. \end{aligned}$$

Proof. Let $g \in C^2[0, 1]$. If we use the Taylor's expansion of the function g at $s = x$, we have

$$g(s) = g(x) + (s-x)g'(x) + \frac{(s-x)^2}{2}g''(x).$$

Hence we get

$$\begin{aligned} \|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq & \|L_{n,p,q}^{\alpha,\beta}((s-x); x)\|_{C[0,1]} \|g(x)\|_{C^2[0,1]} \\ & + \frac{1}{2} \|L_{n,p,q}^{\alpha,\beta}((s-x)^2; x)\|_{C[0,1]} \|g(x)\|_{C^2[0,1]}. \end{aligned} \tag{3.6}$$

From the lemma (2.1) we have

$$\|L_{n,p,q}^{\alpha,\beta}((s-x); x)\|_{C[0,1]} \leq \frac{[\alpha] + [\beta]}{[n]_{p,q} + [\beta]}. \tag{3.7}$$

So if we use (3.3) and (3.7) in (3.6), then we get

$$\begin{aligned} \|L_{n,p,q}^{\alpha,\beta}(g; x) - g(x)\|_{C[0,1]} \leq & \left[\frac{1}{2} \left(\frac{q^2 [n]_{p,q} [n-1]_{p,q}}{([n]_{p,q} + \beta)^2 (p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) \right. \\ & + \left(\frac{1}{2} \frac{p^{n-1} [n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) + \frac{1}{2} \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \tag{3.8} \\ & \left. + \frac{[\alpha] + [\beta]}{[n] + [\beta]} \right] \|g(x)\|_{C[0,1]}. \end{aligned} \tag{3.9}$$

On the other hand, we can write

$$\begin{aligned} |L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| &\leq |L_{n,p,q}^{\alpha,\beta}(f - g; x)| + |L_{n,p,q}^{\alpha,\beta}(g; x) - g(x)| \\ &\quad + |f(x) - g(x)|. \end{aligned}$$

If we take the maximum on $[0, 1]$, we have

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq 2\|f - g\|_{C[0,1]} + \|L_{n,p,q}^{\alpha,\beta}(g; x) - g(x)\|_{C[0,1]}. \quad (3.10)$$

If we consider (3.8) in (3.10), we obtain

$$\begin{aligned} \|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} &\leq 2\|f - g\|_{C[0,1]} + \left[\frac{1}{4} \left(\frac{q[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q} + \beta)^2(p(1-x) + qx)} \right. \right. \\ &\quad \left. \left. - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) + \left(\frac{1}{4} \frac{p^{n-1}[n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) \right. \\ &\quad \left. + \frac{1}{4} \frac{\alpha^2}{([n]_{p,q} + \beta)^2} + \frac{1}{2} \frac{[\alpha] + [\beta]}{[n]_{p,q} + [\beta]} \right] \|g(x)\|_{C^2[0,1]}. \end{aligned}$$

If we choose

$$\begin{aligned} \delta_n &= \frac{1}{4} \left(\frac{q^2[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q} + \beta)^2(p(1-x) + qx)} - \frac{[n]_{p,q} - \beta}{[n]_{p,q} + \beta} \right) \\ &\quad + \left(\frac{1}{4} \frac{p^{n-1}[n]_{p,q} - 2\alpha\beta}{([n]_{p,q} + \beta)^2} \right) + \frac{1}{4} \frac{\alpha^2}{([n]_{p,q} + \beta)^2} + \frac{1}{2} \frac{[\alpha] + [\beta]}{[n]_{p,q} + [\beta]}, \end{aligned}$$

then we get

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq 2 \left\{ \|f - g\|_{C[0,1]} + \delta_n \|g(x)\|_{C^2[0,1]} \right\}. \quad (3.11)$$

Finally, one can observe that if we take the infimum of both side of above inequality for the function $g \in C^2[0, 1]$, we can find

$$\|L_{n,p,q}^{\alpha,\beta}(f; x) - f(x)\|_{C[0,1]} \leq 2K(f, \delta_n).$$

□

4. THE RATES OF STATISTICAL CONVERGENCE

At this point, let us recall the concept of statistical convergence. The statistical convergence which was introduced by Fast [41] in 1951, is an important research area in approximation theory. In [41], Gadjiev and Orhan used the concept of statistical convergence in approximation theory. They proved a Bohman-Korovkin type theorem for statistical convergence.

Recently, statistical approximation properties of many operators are investigated in [4, 25, 26, 29, 30].

A sequence $x = (x_k)$ is said to be statistically convergent to a number L if for every $\epsilon > 0$,

$$\delta\{K \in \mathbf{N} : |x_k - L| \geq \epsilon\} = 0,$$

where $\delta(K)$ is the natural density of the set $K \subseteq \mathbf{N}$.

The density of subset $K \subseteq N$ is defined by

$$\delta(K) := \lim_n \frac{1}{n} \{\text{the number } k \leq n : k \in K\}$$

whenever the limit exists.

For instance, $\delta(\mathbf{N}) = 1$, $\delta\{2K : k \in \mathbf{N}\} = \frac{1}{2}$ and $\delta\{k^2 : K \in \mathbf{N}\} = 0$.

To emphasize the importance of the statistical convergence, we have an example: The sequence

$$X_k = \begin{cases} L_1; & \text{if } k = m^2, \\ L_2; & \text{if } k \neq m^2. \end{cases} \quad \text{where } m \in \mathbf{N} \quad (4.1)$$

is statistically convergent to L_2 but not convergent in ordinary sense when $L_1 \neq L_2$. We note that any convergent sequence is statistically convergent but not conversely.

Now we consider sequences $q = q_n$ and $p = p_n$ such that:

$$st - \lim_n q_n = 1, \quad st - \lim_n p_n = 1, \quad \text{and} \quad st - \lim_n q_n^n = 1. \quad (4.2)$$

Gadjiev and Orhan [41] gave the following theorem for linear positive operators which is about statistically Korovkin type theorem. Now, we recall this theorem.

Theorem 4.1. If A_n be the sequence of linear positive operators from $C[a, b]$ to $C[a, b]$ satisfies the conditions

$$st - \lim_n \|A_n((t^\nu; x)) - (x)^\nu\|_{C[0, 1]} = 0 \text{ for } \nu = 0, 1, 2.$$

then for any function $f \in C[a, b]$,

$$st - \lim_n \|A_n(f; \cdot) - f\|_{C[a, b]} = 0.$$

Now we will discuss the rates of statistical convergence of $L_{n,p,q}^{\alpha,\beta}$ operators.

Remark 4.2. For $q \in (0, 1)$ and $p \in (q, 1]$, it is obvious that

$$\lim_{n \rightarrow \infty} [n]_{p,q} = \begin{cases} 0, & \text{when } p, q \in (0, 1) \\ \frac{1}{1-q}, & \text{when } p = 1 \text{ and } q \in (0, 1). \end{cases}$$

In order to reach to convergence results of the operator $L_{p,q}^n(f; x)$, we take a sequence $q_n \in (0, 1)$ and $p_n \in (q_n, 1]$ such that $\lim_{n \rightarrow \infty} p_n = 1$, $\lim_{n \rightarrow \infty} q_n = 1$. So we get $\lim_{n \rightarrow \infty} [n]_{p_n, q_n} = \infty$.

Theorem 4.3. Let $L_{n,p,q}^{\alpha,\beta}$ be the sequence of operators and the sequences $p = p_n$ and $q = q_n$ satisfies Remark 4.2 then for any function $f \in C[0, 1]$

$$st - \lim_n \|L_{n,p_n,q_n}^{\alpha,\beta}(f, \cdot) - f\| = 0. \quad (4.3)$$

Proof. Clearly for $\nu = 0$,

$$L_{n,p_n,q_n}^{\alpha,\beta}(1, x) = 1,$$

which implies

$$st - \lim_n \|L_{n,p_n,q_n}^{\alpha,\beta}(1; x) - 1\| = 0.$$

For $\nu = 1$

$$\begin{aligned} \|L_{n,p_n,q_n}^{\alpha,\beta}(t; x) - x\| &\leq \left| \frac{[n]_{p_n,q_n}}{[n]_{p_n,q_n} + \beta} x + \frac{\alpha}{[n]_{p_n,q_n} + \beta} - x \right| \\ &= \left| \left(\frac{[n]_{p_n,q_n}}{[n]_{p_n,q_n} + \beta} - 1 \right) x + \frac{\alpha}{[n]_{p_n,q_n} + \beta} \right| \\ &\leq \left| \frac{[n]_{p_n,q_n}}{[n]_{p_n,q_n} + \beta} - 1 \right| + \left| \frac{\alpha}{[n]_{p_n,q_n} + \beta} \right|. \end{aligned}$$

For a given $\epsilon > 0$, let us define the following sets.

$$\begin{aligned}
 U &= \{n : \|L_{n,p_n,q_n}^{\alpha,\beta}(t; x) - x\| \geq \epsilon\} \\
 U' &= \left\{n : 1 - \frac{[n]_{p_n,q_n}}{[n]_{p_n,q_n} + \beta} \geq \epsilon\right\} \\
 U'' &= \left\{n : \frac{\alpha}{[n]_{p_n,q_n} + \beta} \geq \epsilon\right\}.
 \end{aligned}$$

It is obvious that $U \subseteq U'' \cup U'$,

So using

$$\delta\{k \leq n : 1 - \frac{[n]_{p_n,q_n}}{[n]_{p_n,q_n} + \beta} \geq \epsilon\},$$

then we get

$$st - \lim_n \|L_{n,p_n,q_n}^{\alpha,\beta}(t; x) - x\| = 0. \tag{4.4}$$

Lastly for $\nu = 2$, we have

$$\begin{aligned}
 \|L_{n,p_n,q_n}^{\alpha,\beta}(t^2; x) - x^2\| &\leq \left| \frac{q^2 [n]_{p_n,q_n} [n-1]_{p_n,q_n}}{p(1-x) + qx} \frac{1}{([n]_{p_n,q_n} + \beta)^2} - 1 \right| \\
 &\quad + \left| \frac{[n]_{p_n,q_n} (2\alpha + p^{n-1})^2}{[n]_{p_n,q_n} + \beta} x \right| + \left| \frac{\alpha^2}{([n]_{p_n,q_n} + \beta)^2} \right|.
 \end{aligned}$$

If we choose

$$\begin{aligned}
 \alpha_n &= \frac{q^2 [n]_{p_n,q_n} [n-1]_{p_n,q_n}}{p(1-x) + qx} \frac{1}{([n]_{p_n,q_n} + \beta)^2} - 1 \\
 \beta_n &= \frac{[n]_{p_n,q_n} (2\alpha + p^{n-1})^2}{[n]_{p_n,q_n} + \beta} \\
 \gamma_n &= \frac{\alpha^2}{([n]_{p_n,q_n} + \beta)^2}.
 \end{aligned}$$

Then

$$st - \lim_n \alpha_n = st - \lim_n \beta_n = st - \lim_n \gamma_n = 0.$$

Now given $\epsilon > 0$, we define the following four sets:

$$\begin{aligned}
 U &= \|L_{n,p_n,q_n}^{\alpha,\beta}(t^2; x) - x^2\| \geq \epsilon, \\
 U_1 &= \left\{n : \alpha_n \geq \frac{\epsilon}{3}\right\}, \\
 U_2 &= \left\{n : \beta_n \geq \frac{\epsilon}{3}\right\},
 \end{aligned}$$

$$U_3 = \{n : \gamma_n \geq \frac{\epsilon}{3}\}.$$

It is obvious that $U \subseteq U_1 \cup U_2 \cup U_3$. Thus we obtain

$$\begin{aligned} & \delta\{K \leq n : \|L_{n,p,q}^{\alpha,\beta}(t^2 : x) - x^2\| \geq \epsilon\} \\ & \leq \delta\{K \leq n : \alpha_n \geq \frac{\epsilon}{3}\} + \delta\{K \leq n : \beta_n \geq \frac{\epsilon}{3}\} + \delta\{K \leq n : \gamma_n \geq \frac{\epsilon}{3}\}. \end{aligned}$$

So the right hand side of the inequalities is zero.

Then

$$st - \lim_n \|L_{n,p_n,q_n}^{\alpha,\beta}(t; x) - x\| = 0$$

holds and thus the proof is completed. □

ACKNOWLEDGEMENT

The authors are thankful to the learned referees for their valuable comments and suggestions leading to improvement of the paper.

REFERENCES

1. T. Acar, (p, q) -Generalization of Szász-Mirakyan operators, *Mathematical Methods in the Applied Sciences*, DOI: 10.1002/mma.3721.
2. T. Acar, A. Aral, S. A. Mohiuddine, On Kantorovich modification of (p, q) -Baskakov operators, *Journal of Inequalities and Applications*, **98**(2016).
3. A. Aral, V. Gupta, R. P. Agarwal, *Applications of q -Calculus in Operator Theory*, Springer, New York, 2013.
4. O. H. H. Edely, S. A. Mohiuddine, Abdullah K. Noman, Korovkin type approximation theorems obtained through generalized statistical convergence, *Appl. Math. Lett.*, **23**(11), (2010), 1382-1387.
5. M. R. Eslahchi, S. Amani, The best uniform polynomial approximation of two classes of rational functions, *Iranian Journal of Mathematical Sciences and Informatics* Vol. 7, No. 2 (2012), pp 93-102.
6. S. N. Bernstein, constructive proof of Weierstrass approximation theorem, *Comm. Kharkov Math. Soc.* (1912)
7. P.E. Bèzier, *Numerical Control-Mathematics and applications*, John Wiley and Sons, London, 1972.
8. L.-W. Han, Y. Chu, Zh.-Y. Qiu, Generalized Bèzier curves and surfaces based on Lupaş q -analogue of Bernstein operator, *Jour. Comput. Appl. Math.* **261**, (2014), 352-363.
9. Q.-B. Cai, G. Zhou, On (p, q) -analogue of kantorovich type Bernstein-stancu-schurer operators, *Appl. Math. Comput.*, **276**, (2016), 1220.
10. M. Norbert Hounkonnou, Joseph Désiré Bukweli Kyemba, $\mathcal{R}(p, q)$ -calculus: differentiation and integration, *SUT Journal of Mathematics*, **49**(2), (2013), 145-167.

11. R. Jagannathan, K. Srinivasa Rao, Two-parameter quantum algebras, twin-basic numbers and associated generalized hypergeometric series, *Proceedings of the International Conference on Number Theory and Mathematical Physics*, (2005), 20-21.
12. U. Kadak, On weighted statistical convergence based on (p, q) -integers and related approximation theorems for functions of two variables, *Journal of Mathematical Analysis and Applications*, (2016), DOI: 10.1016/j.jmaa.2016.05.062.
13. U. Kadak, Naim L. Braha, H. M. Srivastava, Statistical weighted B-summability and its applications to approximation theorems, *Appl. Math. Comput.*, **302**, (2017), 80-96.
14. U. Kadak, Generalized weighted invariant mean based on fractional difference operator with applications to approximation theorems for functions of two variables, *Results in Mathematics*, (2016), doi:10.1007/s00025-016-0634-8.
15. U. Kadak, Weighted statistical convergence based on generalized difference operator involving (p, q) -Gamma function and its applications to approximation theorems, *Journal of Mathematical Analysis and Applications*, **448**(2), (2017), 1633-1650.
16. Kh. Khan, D. K. Lobiyal, A. Kilicman, A de casteljau algorithm for bernstein type polynomials based on (p, q) -integers, *Appl. Appl. Math.*, **13**(2) (2018).
17. Kh. Khan, D.K. Lobiyal, Bézier curves based on lupas (p, q) -analogue of Bernstein functions in CAGD, *Jour. Comput. Appl. Math.*, **317**, (2017), 458-477.
18. Kh. Khan, D.K. Lobiyal, Bézier curves and surfaces based on modified Bernstein polynomials, *Azerb. j. Math.*, **9**(1), 2019.
19. V. Kac, P. Cheung, *Quantum Calculus*, in: Universitext Series, vol.IX, Springer-Verlag, (2002).
20. P. P. Korovkin, Linear operators and approximation theory, *Hindustan Publishing Corporation*, Delhi, 1960.
21. A. Lupaş, A q -analogue of the Bernstein operator, Seminar on Numerical and Statistical Calculus, *University of Cluj-Napoca*, **9**(1987), 85-92.
22. N. I. Mahmudov, P. Sabancigil, Some approximation properties of Lupaş q -analogue of Bernstein operators, *arXiv:1012.4245v1 [math.FA]* 20 Dec 2010.
23. V. N. Mishra, S. Pandey, On (p, q) Baskakov-Durrmeyer-Stancu Operators, *Advances in Applied Clifford Algebras*, (2016), DOI: 10.1007/s00006-016-0738-y.
24. V. N. Mishra, S. Pandey; On Chlodowsky variant of (p, q) Kantorovich-Stancu-Schurer operators, *International Journal of Analysis and Applications*, **11**(1), (2016), 28-39.
25. N. L. Braha, H. M. Srivastava, S. A. Mohiuddine, A Korovkin's type approximation theorem for periodic functions via the statistical summability of the generalized de la Valle Poussin mean. *Appl. Math. Comput.*, **228**, (2014), 162-169.
26. C. Belen, S. A. Mohiuddine: Generalized weighted statistical convergence and application. *Appl. Math. Comput.*, **219**(18), (2013), 9821-9826 .
27. M. Mursaleen, K. J. Ansari, A. Khan, On (p, q) -analogue of Bernstein Operators, *Appl. Math. Comput.*, **266**(2015) 874-882, [Erratum: 278 (2016) 70-71].
28. M. Mursaleen, Md. Nasiruzzaman, A. Nurgali, Some approximation results on Bernstein-Schurer operators defined by (p, q) -integers, *Journal of Inequalities and Applications*, **249**, (2015), DOI 10.1186/s13660-015-0767-4.
29. M. Mursaleen, A. Khan, Statistical Approximation Properties of Modified q -Stancu-Beta Operators, *Bull. Malays. Math. Sci. Soc.*, (2) **36**(3) (2013), 683690.
30. M. Mursaleen, F. Khan, A. Khan, Statistical approximation for new positive linear operators of Lagrange type, *Appl. Math. Comput.*, **232**(2014) 548558.

31. M. Mursaleen, K. J. Ansari, A. Khan, Some Approximation Results by (p, q) -analogue of Bernstein-Stancu Operators, *Appl. Math. Comput.*, **264**, (2015), 392-402.
32. M. Mursaleen, K. J. Ansari, A. Khan, Some approximation results for Bernstein-Kantorovich operators based on (p, q) -calculus, *U.P.B. Sci. Bull., Series A*, **78**(4), (2016).
33. M. Mursaleen, Md. Nasiruzzaman, A. Khan, K. J. Ansari, Some approximation results on Bleimann-Butzer-Hahn operators defined by (p, q) -integers, *Filomat*, **30**(3), (2016), 639-648, DOI 10.2298/FIL1603639M.
34. M. Mursaleen, A. Khan, Generalized q -Bernstein-Schurer Operators and Some Approximation Theorems, *Journal of Function Spaces and Applications* **2013**, Article ID 719834, 7 pages <http://dx.doi.org/10.1155/2013/719834>
35. M. Mursaleen, F. Khan, A. Khan, Approximation by (p, q) -Lorentz polynomials on a compact disk, *Complex Anal. Oper. Theory*, DOI: 10.1007/s11785-016-0553-4.
36. M. Mursaleen, Md. Nasiruzzaman, F. Khan, A. Khan, On (p, q) -analogue of divided differences and Bernstein operators, *Journal of Nonlinear Functional Analysis*, **2017**, (2017), Article ID 25, pp. 1-13.
37. S. Ostrovska, On the Lupaş q -analogue of the Bernstein operator, *Rocky mountain journal of mathematics*, **36**(5), 2006.
38. G. M. Phillips, Bernstein polynomials based on the q -integers, The heritage of P.L.Chebyshev, *Ann. Numer. Math.*, **4**, (1997), 511-518.
39. G. M. Phillips, A generalization of the Bernstein polynomials based on the q -integers *ANZIAMJ* **42**, (2000), 79-86.
40. A. Rababah, S. Manna, Iterative process for G2-multi degree reduction of Bèzier curves, *Appl. Math. Comput.*, 217 (2011) 8126-8133.
41. A. D. Gadjevi, C. Orhan, Some approximation theorems by statistical convergence, *Rocky Mountain J.math.*, **32**, (2002) 129-138.
42. A. Wafi, N. Rao, Bivariate-Schurer-Stancu operators based on (p, q) -integers, *Filomat*, **32**(4), (2018), 1251-1258.
43. A. Wafi, N. Rao, (p, q) -Bivariate-Bernstein-Chlowdosky Operators, *Filomat*, **32**(2), (2018), 369378.
44. A. Wafi, N. Rao, Approximation properties of (p, q) -variant of Stancu-Schurer, *Boletim da Sociedade Paranaense de Matematica* **37**(4), (2017).
45. A. Wafi, N. Rao, A generalization of Szasz-type operators which preserves constant and quadratic test functions, *cogent mathematics*, **3**(1), 2016, 1227023.
46. A. Wafi, N. Rao Modified Szasz operators involving Charlier polynomials based on two parameters, *Thai J. of Maths.*, 2016.
47. A. Wafi, N. Rao Approximation properties by generalized Baskakov Kantorovich Stancu type operators, *Appl. Maths Inf. Sci. Lett.*, **4**(3) (2016), PP:111-118.
48. A. Wafi, N. Rao, Stancu-variant of generalized Baskakov operators, *Filomat*, **31**(9), (2017).