Iranian Journal of Mathematical Sciences and Informatics Vol. 18, No. 1 (2023), pp 109-129 DOI: 10.52547/ijmsi.18.1.109

# An Optimal Algorithm for the $\delta$ -ziti Method to Solve Some Mathematical Problems

L. Bsiss<sup>\*</sup>, C. Ziti

Department of Mathematics, University Moulay Ismaïl, Faculty of Sciences, BP 11201 Zitoune, Meknès 50000, Morocco

> E-mail: lyrbi01@gmail.com E-mail: chziti@gmail.com

ABSTRACT. The numerical approximation methods of the differential problems solution are numerous and various. Their classifications are based on several criteria: Consistency, precision, stability, convergence, dispersion, diffusion, speed and many others. For this reason a great interest must be given to the construction and the study of the associated algorithm: indeed the algorithm must be simple, robust, less expensive and fast. In this paper, after having recalled the  $\delta$ -ziti method, we reformulat it to obtain an algorithm that does not require as many calculations as many nodes knowing that they are counted by thousands. We have, therefore, managed to optimize the number of iterations by passing for example from 10<sup>3</sup> at 10 iterations.

**Keywords:** Algorithm, Meshing,  $\delta$ -ziti, Optimal, Operations number.

## 2000 Mathematics subject classification: 65D15, 65N06, 65Q10.

## 1. INTRODUCTION

109

The  $\delta$ -ziti is a new method of approximation, which allows:

(1) to interpolate, approach and integrate a function of one or more variables,

[ Downloaded from ijmsi.ir on 2025-07-04

<sup>\*</sup>Corresponding Author

Received 02 April 2018; Accepted 08 April 2021 ©2023 Academic Center for Education, Culture and Research TMU

#### L. Bsiss, C. Ziti

- (2) to solve a differential equation of any order,
- (3) to solve an equation or a system with partial derivatives.

Generally, in concrete problems, one confronts with singularities that generate horrible instabilities in the form of oscillations where the classical numerical methods are very sensitive [5](ie improper integrals (generalized), functions which are not sufficiently regular or distributions as Dirac type, singular differential problem, presence of natural phenomena such as shocks, boundary layers, Blow-up, turbulence and others ...). The  $\delta$ -ziti method has shown its effectiveness in solving such problems. When presenting the  $\delta$ -ziti method in [1, 2, 3, 4], we did several tests on several mathematical models. We have established algorithms without considering the cost of calculation and their complications. In this work, we will show how to minimize the cost of calculation by a very small number of iterations. First, we will recall, in the following paragraph, the main lines of the  $\delta$ -ziti method.

#### 2. Overview of the $\delta$ -ziti method.

As it was presented in [3], the method is based, essentially on the famous function

$$\phi(x) = \begin{cases} exp\left(\frac{1}{|x|^2 - R^2}\right) & ; if \ |x|^2 := \sum_{i=1}^n x_i^2 < R, \\ 0 & otherwise. \end{cases}$$
(2.1)

where R is a positive constant.

This function is very often encountered in Numerical Analysis, especially in distributions and functional analysis. It is characterized by its power to approach the Dirac measurement:

indeed, if we put,

$$\varphi_{\epsilon}(x) = \frac{C}{\epsilon^n} \phi\left(\frac{x}{\epsilon}\right) \quad \forall \epsilon > 0, \tag{2.2}$$

where,  $C := \frac{1}{\int_{\mathbb{R}^n} \phi(x) dx}$ , then this sequence  $\varphi_{\epsilon}$  converges to Dirac in the sense of distributions.

The Dirac distribution is a very significant symbol of a singularity, which made us think of using  $\phi(x)$  to deal with singularities. At first, we are interested in the one dimensional case (n = 1). For the multidimensional case, we will see how we can reduce ourselves to the dimensional one. Consider, for the moment, an interval bounded [a, b]. We consider a uniform meshing, the interval [a, b] is subdivided into (m + 1) points (where m is the number of subintervals: (m + 1) nodes) with a constant step h:  $x_1 = a$ ,  $x_{m+1} = b$ ,  $x_i = a + (i - 1)h$  for  $i = 1, \ldots, m + 1$ .

where,  $h = \frac{b-a}{m}$ .

At each  $x_i$ , we associate the function  $\varphi_i$  defined by:

$$\varphi_i(x) = \frac{C}{h} \phi\left(\frac{x - x_i}{h}\right) \text{ for } x \in [x_{i-1}, x_{i+1}] \ (i = 2, \dots, m), \qquad (2.3)$$

An Optimal Algorithm for the  $\delta$ -ziti Method to Solve Some Mathematical Problems 111

$$\varphi_1(x) = \frac{C}{h} \cdot \phi\left(\frac{x-x_1}{h}\right) \text{ for } x \in [x_1, x_2], \qquad (2.4)$$

and,

$$\varphi_{m+1}(x) = \frac{C}{h} \cdot \phi\left(\frac{x - x_{m+1}}{h}\right) \quad \text{for } x \in [x_m, x_{m+1}]. \tag{2.5}$$

This family of functions is represented in the following figures:



FIGURE 1. Superposition of some elements of the family  $(\varphi_i)_{2 \le i \le m}$  for a=-4,b=4,h=1 and C=2.252283621.



FIGURE 2. Superposition of the consecutive elements  $\varphi_{i-1}$ ,  $\varphi_i$  and  $\varphi_{i+1}$  for a=-4,b=4,h=1 and C=2.252283621.



FIGURE 3. Graphical representation of extreme elements  $\varphi_1$  and  $\varphi_{m+1}$  for a=-4,b=4,h=1 and C=2.252283621.

By choosing a suitable Prehilbert space with the scalar product  $\langle , \rangle$ , the norm associated is  $\|.\|$ , we use the Gram-Schmidt process, to construct the orthogonal family  $(\tilde{\psi}_i)_i$ .

We have shown that the family  $\left(\tilde{\psi}_i\right)_i$  satisfies the following recurrence relation:

$$\begin{cases}
\tilde{\psi}_{1}(x) = \varphi_{1}(x) \text{ for all } x \in [x_{1}, x_{m+1}] \\
\tilde{\psi}_{i}(x) = \varphi_{i}(x) + \lambda_{i-1}\tilde{\psi}_{i-1}(x) \text{ for all } x \in [x_{1}, x_{m+1}], \ 2 \leq i \leq m+1 \\
supp\tilde{\psi}_{i} = [x_{1}, x_{i+1}] \text{ pour tout indice } i, \\
with : \lambda_{i-1} = -\frac{\langle \varphi_{i}, \tilde{\psi}_{i-1} \rangle}{\|\tilde{\psi}_{i-1}\|^{2}},
\end{cases}$$
(2.6)

and that

 $-1 < \lambda_k < 0$  for  $1 \le k \le m$ ,

From this relation of recurrence, one can develop  $\tilde{\psi}_i$  according to  $\varphi_k$   $(1 \le k \le i)$ :  $\tilde{\psi}_i(x) = \varphi_i(x) + \lambda_{i-1}\varphi_{i-1}(x) + \lambda_{i-1}\lambda_{i-2}\varphi_{i-2}(x) + \dots + \lambda_{i-1}\lambda_{i-2}\dots\lambda_1\varphi_1(x).$ We construct an algorithm with (2.6) to compute  $\tilde{\psi}_i$ ,  $\lambda_i$  and  $< \tilde{\psi}_i, \tilde{\psi}_i > .$ The functions  $(\tilde{\psi}_i)_i$  are represented in the following figures: Finally, we orthonormalize  $(\tilde{\psi}_i)$  to obtain the family  $(\psi_i)$ :

$$\psi_i(x) = \frac{\tilde{\psi}_i(x)}{\left\|\tilde{\psi}_i\right\|} \text{ pour } 1 \le i \le m+1$$
(2.7)

*Remark* 2.1. The algorithm we are going to build aims to calculate  $\lambda_i$ ,  $\psi_i$  and roots  $r_i$  and to simulate the results.

An Optimal Algorithm for the  $\delta\textsc{-ziti}$  Method to Solve Some Mathematical Problems  $\quad$  113



FIGURE 4. Superposition of some elements of the orthogonal family  $\left(\tilde{\psi}_i\right)_{1 \le i \le m+1}$  for a=-4,b=4,h=1 and c=0.1330861208.



FIGURE 5. Graphical representation of  $\tilde{\psi}_6$  for a=-4,b=4,h=1 and c=0.1330861208.

## 2.1. algorithm 1.

a = -4, b = 4; ( the border of the interval [a, b]) m = 800; (number of intervals after subdivision of [a, b])  $h = \frac{b-a}{m}$ ; (the step of discretization) **Discretization of [a, b]** 

$$\begin{cases} for \ i = 1 : m+1 \\ x_i = a + (i-1)h \\ end \ i \end{cases}$$

Construction of  $(\varphi_i)$ 

$$\begin{split} \phi(x) &= \left\{ \begin{array}{l} \exp\left(\frac{1}{|x|^2 - R^2}\right) &; if \ |x|^2 := \sum_{i=1}^n x_i^2 < R, \\ 0 & otherwise. \end{array} \right. \\ \varphi_1(x) &= \frac{C}{h} \cdot \phi\left(\frac{x - x_1}{h}\right) \ for \ x \in [x_1, x_2], \\ \left\{ \begin{array}{l} for \ i = 2 : m \\ \varphi_i(x) = \frac{C}{h} \cdot \phi\left(\frac{x - x_{m+1}}{h}\right) \ for \ x \in [x_{i-1}, x_{i+1}] \\ end \ i \end{array} \right. \\ \varphi_{m+1}(x) &= \frac{C}{h} \cdot \phi\left(\frac{x - x_{m+1}}{h}\right) \ for \ x \in [x_m, x_{m+1}] \end{aligned} \\ \end{split} \\ \\ \begin{aligned} \text{Construction of } \left(\tilde{\psi_i}\right) \\ \left\{ \begin{array}{l} for \ i = 1 \\ \tilde{\psi_1}(x) = \varphi_1(x), \\ \|\tilde{\psi_1}\|^2 = \int_a^b \tilde{\psi_1}(x)\tilde{\psi_1}(x)dx, \\ \lambda_1 &= \frac{\int_a \varphi_1(x)\varphi_2(x)dx}{\|\tilde{\psi_1}\|^2} \\ \left\{ \begin{array}{l} for \ i = 2 : m \\ \tilde{\psi_i}(x) = \varphi_i(x) + \lambda_{i-1}\tilde{\psi}_{i-1}(x), \\ \|\tilde{\psi_i}\|^2 = \int_a^b \tilde{\psi_i}(x)\tilde{\psi_i}(x)dx, \\ \lambda_i &= \frac{\int_a^b \varphi_i(x)\tilde{\psi_i}(x)dx, \\ \lambda_i &= \frac{\int_a^b \varphi_i(x)\varphi_{i+1}(x)dx}{\|\tilde{\psi_i}\|^2}, \\ end \ i \\ \end{array} \\ \\ \begin{cases} for \ i &= m+1 \\ \tilde{\psi_{m+1}}(x) &= \varphi_{m+1}(x) + \lambda_m \tilde{\psi}_m(x), \end{array} \end{aligned} \end{aligned}$$

$$\begin{cases} \psi_{m+1}(x) = \psi_{m+1}(x) + \lambda_m \psi_m(x), \\ \|\tilde{\psi}_{m+1}\|^2 = \int_a^b \tilde{\psi}_{m+1}(x) \tilde{\psi}_{m+1}(x) dx, \end{cases}$$

Calculating the roots of  $\left( \tilde{\psi}_i \right)$ 

$$r = Roots(\tilde{\psi}_i(x), x = a..b)$$

 $Remark\ 2.2.$  To compute any integral, we use any approximation's method as Simpson-method for example.

i	$\lambda_i$	$r_i$
1	-0.508960181	-3.996667413
2	-0.292344636	-3.987512758
3	-0.274934054	-3.977583248
4	-0.273624280	-3.967588576
5	-0.273526252	-3.957588975
6	-0.273518918	-3.947589005
7	-0.273518369	-3.937589007
8	-0.273518328	-3.927589007
9	-0.273518325	-3.917589007
10	-0.273518325	-3.907589007
11	-0.273518325	-3.897589007
12	-0.273518325	-3.887589007
13	-0.273518325	-3.877589007
14	-0.292344636	-3.987512758
795	-0.273518325	3.942410993
796	-0.273518325	3.952410993
797	-0.273518325	3.962410993
798	-0.273518325	3.972410993
799	-0.273518325	3.982410993
800	-0.273518325	3.992410993
801		4

TABLE 1. Results obtained by the first algorithm with 801 nodes.

## 2.2. Roots of $\psi_i$ .

The  $\delta$ -ziti method mainly uses the roots  $(r_i)$  of the functions  $(\psi_i)_i$  instead of the mesh point  $(x_i)$ . It is not a question of replacing  $(x_i)$  by the roots  $(r_i)$  but

it is about a radical transformation of the approximation.

We studied minorly, the roots  $(r_i)$  and we showed that

- (1) the roots of  $\psi_i$  is, also, root of  $\psi_k$  for:  $k \ge i 1$ , in particular are also roots of  $\psi_{m+1}$
- (2)  $\psi_i$  admits (i-1) distinct real root(s)  $r_k$   $(k = 1; \dots; i-1)$  in the interval  $]x_1, x_i[$  precisely  $r_k \in ]x_k, x_{k+1}[$  for  $k = 1, \dots, i-1,$
- (3)  $r_j \in \left] x_j, x_j + \frac{h}{2} \right[,$
- (4)

$$\lambda_{i-1} = -\frac{\varphi_{i+1}(r_i)}{\varphi_i(r_i)}.$$
(2.8)

What allowed us to write

$$r_k = x_k + h.X_k,\tag{2.9}$$

where:  $X_k$  is solution of the following polynom

$$\Lambda_k X^4 - 2\Lambda_k X^3 - \Lambda_k X^2 + 2(\Lambda_k - 1)X + 1 = 0, \qquad (2.10)$$

where: 
$$\Lambda_k = Ln(-\lambda_k)$$
 and  $\lambda_{i-1} = -\frac{\langle \varphi_i, \varphi_{i-1} \rangle}{\|\tilde{\psi}_{i-1}\|^2}$ .

*Remark* 2.3. A second algorithm differs from the previous one can be built using the results (2.8), (2.9) and (2.10) but still with always the same number of iterations as the number of nodes.

The functions  $(\psi_i)$  and the position of their roots are represented in the following figures:



FIGURE 6. Superposition of some elements of the family  $(\psi_k)_{1 \le k \le m+1}$  and position of their roots for a=-4,b=4,h=1 and c=0.1330861208.



FIGURE 7. Superposition of consecutive elements  $\psi_{i-1}$ ,  $\psi_i$ ,  $\psi_{i+1}$  and position of its roots for a=-4,b=4,h=1 and c=0.1330861208.



FIGURE 8. Representation of element  $\psi_5$  and position of its roots for a=-4, b=4, h=1 and C=2.252283621.



FIGURE 9. The position of the roots  $r_i$  of  $\psi_{m+1}$ .

#### 2.3. Description of $\delta$ -ziti method.

The  $\delta$ -ziti method starts, like the finite element method [6, 7, 8], with a variational formulation to have an adequate functional framework. To avoid the mesh  $(x_i)$  and its complications in the variational formulation, we used the roots  $(r_i)$  of the extreme function  $\psi_{m+1}$  which made it possible to give a better approximation of a function and an integral one by a simple formula.

Remark 2.4. The use of only  $\psi_{m+1}$  and not the others  $\psi_i$  is because every root of  $\psi_i$  is also root of  $\psi_{m+1}$ .

2.3.1. Interpolation and approximation of a function with a single variable. For a given function f, we approached it as follows:

$$f(x) \simeq \sum_{i=1}^{i=m+1} \alpha_i \psi_i(x).$$
 (2.11)

where,

$$\alpha_i = \langle f, \psi_i \rangle,$$

We showed that,

$$\alpha_i = \frac{f(r_i)}{\psi_i(r_i)}.$$

2.3.2. The numerical integration.

We used the following approximation:

$$\int_{a}^{b} f(x)\psi_{i}(x)dx \simeq \frac{f(r_{i})}{\psi_{i}(r_{i})} \text{ for } i = 1...m + 1.$$
(2.12)

$$\int_{a}^{b} f(x)dx \simeq \sum_{i=1}^{i=m+1} \frac{f(r_i)}{\psi_i^2(r_i)}.$$
(2.13)

2.3.3. Function with several variables.

Let us take,  $\Omega$  in the form:  $\Omega = [a, b] \times [c, d]$ , and use an uniform mesh of the interval [a, b] and [c, d] with the step  $h_1 = \frac{b-a}{m_1}$  and  $h_2 = \frac{d-c}{m_2}$  where  $m_1, m_2$  are integers such  $x_1 = a, x_{m_1+1} = b$  and  $y_1 = c, y_{m_2+1} = d, x_k = a + (k-1)h_1$  for:  $k = 1, \ldots, m_1 + 1, y_k = c + (l-1)h_2$  for:  $l = 1, \ldots, m_2 + 1$ .

From (2.6) and (2.7), we construct two orthonormal families:

$$(\psi_k^1)_{1 \le k \le m+1}$$
 and  $(\psi_k^2)_{1 \le k \le m+1}$ 

Our strategy is to take:

$$\psi_{i,j}(x,y) = \psi_i^1(x)\psi_j^2(y) \text{ for } (x,y) \in \Omega.$$
 (2.14)

## i: Approximation of a function of two variables.

Let f be a function of two variables, f(x, y) can be approximated by:

$$f(x,y) \simeq \sum_{i=1}^{m_1+1} \sum_{j=1}^{m_2+1} \alpha_{ij} \psi_i^1(x) \psi_j^2(y).$$
(2.15)

where,

$$\alpha_{ij} = \frac{f(r_i^1, r_j^2)}{\psi_i^1(r_i^1)\psi_j^2(r_j^2)}.$$

#### ii: The numerical integration.

We have approached a multiple integral as follows:

$$\int_{a}^{b} \int_{c}^{d} f(x,y)\psi_{i}^{1}(x)\psi_{j}^{2}(y)dxdy \simeq \frac{f(r_{i}^{1},r_{j}^{2})}{\psi_{i}^{1}(r_{i}^{1})\psi_{j}^{2}(r_{j}^{2})},$$
$$\int_{a}^{b} \int_{c}^{d} f(x,y)dxdy \simeq \sum_{i=1}^{m_{1}+1} \sum_{j=1}^{m_{2}+1} \frac{f(r_{i}^{1},r_{j}^{2})}{\left(\psi_{i}^{1}(r_{i}^{1})\psi_{j}^{2}(r_{j}^{2})\right)^{2}}.$$

Remark 2.5. The extension of the  $\delta$ -ziti method with several variables is done in the same way:

## i: Approximation of a function.

$$f(y_1,\cdots,y_n) \simeq \sum_{1 \le i_1 \le m_1+1} \cdots \sum_{1 \le i_n \le m_n+1} \alpha_{i_1\cdots i_n} \psi_{i_1,\cdots,i_n}(y_1,\cdots,y_n),$$

where,

$$\alpha_{i_1\cdots i_n} = \frac{f(r_{i_1}^1, \cdots, r_{i_n}^n)}{\psi_{i_1, \cdots, i_n}(r_{i_1}^1, \cdots, i_{i_n}^n)}.$$

## ii: The numerical integration.

We have approached a multiple integral as follows:

$$\int_{a_1}^{b_1} \cdots \int_{a_n}^{b_n} f(x_1, \cdots, x_n) \psi_{i_1}^1(x_1) \cdots \psi_{i_n}^n(x_n) dx_1 \cdots dx_n \simeq \frac{f(r_{i_1}^1, \cdots, r_{i_n}^n)}{\psi_i^1(r_i^1) \cdots \psi_i^n(r_{i_n}^n)},$$

$$\int_{a_1}^{b_1} \cdots \int_{a_n}^{b_n} f(x_1, \cdots, x_n) dx_1 \cdots dx_n \simeq \sum_{1 \le i_1 \le m_1 + 1} \cdots \sum_{1 \le i_n \le m_n + 1} \frac{f(r_{i_1}^1, \cdots, r_{i_n}^n)}{\left(\psi_i^1(r_i^1) \cdots \psi_i^n(r_{i_n}^n)\right)^2},$$
where,  $r_{i_j}^j$  is the  $(i_j)^{th}$  root of  $\psi_{m+1}^j$  on  $[a_j, b_j]$  (for  $i_j = 1, \dots, m_j + 1, j = 1, \dots, n$ ).

## 2.3.4. *Differential equation and partial differential equation.* We consider, therefore, an ordinary differential equation of order 1,

$$(\mathcal{P}) \begin{cases} u'(x) = f(x, u(x)) ; x \in [a, b], & (1) \\ u(a) = \xi_0, & (2) \end{cases}$$
(2.16)

where,  $\xi_0$  is a given constant,  $u : \mathbb{R} \to \mathbb{R}$  is the unknown function and  $f : \mathbb{R}^2 \to \mathbb{R}$  is a given function.

The variational formulation of  $(\mathcal{P})$  consists to multiply the differential equation by  $\psi_k$  and integrating:

$$\int_{a}^{b} u'(x).\psi_{i}(x)dx = \int_{a}^{b} f(x,u(x)).\psi_{i}(x)dx \text{ for } i = 1,\dots,m+1.$$
 (2.17)

The use of the previous integration formula (2.15) made it possible to ignore the points of meshing  $x_i$  and to take into account only the roots  $r_i$ . From the integration formula (2.15) we have:

$$\int_{a}^{b} u'(x) \cdot \psi_i(x) dx \simeq \frac{u'(r_i)}{\psi_i(r_i)}$$

and,

$$\int_{a}^{b} f(x, u(x)) \cdot \psi_i(x) dx \simeq \frac{f(r_i, u(r_i))}{\psi_i(r_i)}.$$

Therefore,

$$u'(r_i) = f(r_i, u(r_i)) \text{ for } i = 1, \dots, m+1,$$
 (2.18)

Remark 2.6. For the differential equations of order one, the  $\delta$ -ziti method resembles, at first sight, the collocation method (moreover the finite element method also coincides with the collocation method in dimension one), which is not the case for the order greater than two.

If we approach the derivative  $u'(r_k)$  in (2.18) by (for example) the following finite difference approximations:

$$u'(r_i) \simeq \frac{u(r_{i+1}) - u(r_i)}{r_{i+1} - r_i}$$
 for  $i = 1, \dots, m$ ,

and let's use the fact that,

$$u(r_i) = \alpha_i \psi_i(r_i) \text{ for } i = 1, \dots, m,$$

then the  $(\alpha_i)$  will be solutions of the following discrete problem:

$$(\mathcal{S}) \begin{cases} \alpha_1 = \frac{\xi_0}{\psi_1(r_1)} \\ \alpha_{i+1} = \frac{\alpha_i \psi_i(r_i) + (r_{i+1} - r_i) f(r_i, \alpha_i \psi_i(r_i))}{\psi_{i+1}(r_{i+1})} \text{ for } i = 1, \dots, m. \end{cases}$$
(2.19)

In all of the above, the calculations of  $r_i$  and  $\lambda_i$  are essential to solve (S) but with what cost?

In all our work where we used the  $\delta$ -ziti method we obtained very satisfactory results and even surprising in comparison with the exact results or with the well-known numerical results. We performed the calculation by following the formulas found mathematically without taking into account their complications and the cost (for example the number of iterations). In the next paragraph we will reduce the cost considerably.

3. Reformulation of  $\lambda_i$  and  $r_i$ .

## 3.1. Calculation of $\lambda_i$ ..

We recall that,

$$\alpha = <\varphi_1, \varphi_2 >, \ \beta = <\varphi_1, \varphi_1 > .$$

The following theorem shows that  $(\lambda_i)$  follows a simple iterative process:

**Theorem 3.1.**  $\lambda_i$  verifies the following recurrence relation:

$$\lambda_{1} = -\frac{\alpha_{\beta}}{\beta},$$
  

$$\lambda_{i+1} = f(\lambda_{i}) \text{ pour } 1 \leq i \leq m-1,$$
  
where  $f(x) = \frac{\lambda_{1}}{2-\lambda_{1}x}.$ 

Proof. We have,

$$\lambda_1 = -\frac{\langle \varphi_1, \varphi_2 \rangle}{\left\|\tilde{\psi}_1\right\|^2} = -\frac{\langle \varphi_1, \varphi_2 \rangle}{\left\|\varphi_1\right\|^2} = -\frac{\alpha}{\beta},$$

and,

$$\lambda_{i} = -\frac{\langle \varphi_{i}, \varphi_{i+1} \rangle}{\left\| \tilde{\psi}_{i} \right\|^{2}},$$

then,

$$<\tilde{\psi}_i, \tilde{\psi}_i> = -\frac{<\varphi_i, \varphi_{i+1}>}{\lambda_i},$$
(3.1)

but,

$$\lambda_{i+1} = -\frac{\langle \varphi_{i+1}, \varphi_{i+2} \rangle}{\left\| \tilde{\psi}_{i+1} \right\|^2},$$

from (2.6) we have,

$$\langle \varphi_{i+1}, \tilde{\psi}_i \rangle = \langle \varphi_{i+1}, \varphi_i \rangle,$$

and

$$\tilde{\psi}_{i+1}(x) = \varphi_{i+1}(x) + \lambda_i \tilde{\psi}_i(x).$$

Therefore,

$$\begin{array}{lll} <\tilde{\psi}_{i+1},\tilde{\psi}_{i+1}> &=& <\varphi_{i+1},\varphi_{i+1}>+2\lambda_i<\varphi_{i+1},\tilde{\psi}_i>+\lambda_i^2<\tilde{\psi}_i,\tilde{\psi}_i>\\ &=& <\varphi_{i+1},\varphi_{i+1}>+2\lambda_i<\varphi_{i+1},\varphi_i>+\lambda_i^2<\tilde{\psi}_i,\tilde{\psi}_i>, \end{array}$$

and

$$\lambda_{i+1} = -\frac{\langle \varphi_{i+1}, \varphi_{i+2} \rangle}{\langle \varphi_{i+1}, \varphi_{i+1} \rangle + 2\lambda_i \langle \varphi_{i+1}, \varphi_i \rangle + \lambda_i^2 \langle \tilde{\psi}_i, \tilde{\psi}_i \rangle}.$$
(3.2)

By injecting (3.1) into (3.2), we obtain,

$$\lambda_{i+1} = -\frac{\langle \varphi_{i+1}, \varphi_{i+2} \rangle}{\langle \varphi_{i+1}, \varphi_{i+1} \rangle + \lambda_i \langle \varphi_{i+1}, \varphi_i \rangle},\tag{3.3}$$

according to [3], for every index i, we have (in fact  $\varphi_i$  is a translation of  $\varphi_2)$ 

$$\langle \varphi_{i+1}, \varphi_i \rangle = \langle \varphi_1, \varphi_2 \rangle = \alpha \ et \ \langle \varphi_{i+1}, \varphi_{i+1} \rangle = 2 \langle \varphi_1, \varphi_1 \rangle = 2\beta$$

(3.3) then becomes

$$\lambda_{i+1} = -\frac{\alpha}{2\beta + \lambda_i \alpha} = \frac{-\frac{\alpha}{\beta}}{2 + \lambda_i \frac{\alpha}{\beta}} = \frac{\lambda_1}{2 - \lambda_1 \lambda_i}.$$

The following result will make it possible to estimate  $|\lambda_{i+1} - \lambda_i|$ .

**Corollary 3.2.** For  $\lambda_1 = -\frac{\alpha}{\beta}$  we have,

(1)

$$0 < \left(\frac{\lambda_1}{2 + \lambda_1}\right)^2 < 1. \tag{3.4}$$

(2)

For all *i* such that 
$$i \ge \left[\frac{\ln\left(\frac{\epsilon(2-\lambda_1^2)}{\lambda_1^3-\lambda_1}\right)}{\ln\left(\frac{\lambda_1}{2+\lambda_1}\right)^2}\right] + 1,$$
 (3.5)

we have

 $|\lambda_{i+1} - \lambda_i| < \epsilon$ , for any arbitrary smal real  $\epsilon > 0$ ,

where [x] is the floor function of x.

*Proof.* (1) We know that

$$\lambda_1 = -\frac{\alpha}{\beta} and -1 < \lambda_1 < 0,$$

 $\mathrm{so},$ 

$$0 < \frac{-\lambda_1}{2+\lambda_1} < 1.$$

Therefore,

$$0 < \left(\frac{\lambda_1}{2 + \lambda_1}\right)^2 < 1.$$

(2) Let f be the function defined by:

$$f(X) = \frac{\lambda_1}{2 - \lambda_1 \cdot X} \ avec \ -1 < X < 0,$$

it's easy to verify that

 $f^{''}(X) < 0 \text{ pour tout } X \in ]-1, 0[.$ 

Therefore,  $f^{'}$  is strictly decreasing on ]-1,0[, a simple calculation shows that

$$0 < \frac{\lambda_{1}^{2}}{4} = f^{'}(0) < f^{'}(X) < f^{'}(-1) = \frac{\lambda_{1}^{2}}{\left(2 + \lambda_{1}\right)^{2}},$$

and

$$f^{'}(X) < \left(\frac{\lambda_{1}}{2+\lambda_{1}}\right)^{2}$$
 for all  $X \in ]-1,0[$ .

By applying the theorem of finite increments on  $]\lambda_i, \lambda_{i+1}[$ , we obtain,

$$\begin{aligned} |\lambda_{i+1} - \lambda_i| &= |f(\lambda_i) - f(\lambda_{i-1})|, \\ &= |f'(\xi)| |\lambda_i - \lambda_{i-1}| \xi \text{ entre } \lambda_{i+1} \text{ et } \lambda_i, \end{aligned}$$

Therefore,

$$\begin{aligned} |\lambda_{i+1} - \lambda_i| &< \left(\frac{\lambda_1}{2+\lambda_1}\right)^2 |f(\lambda_{i-1}) - f(\lambda_{i-2})| \\ &< \dots \end{aligned}$$

[ Downloaded from ijmsi.ir on 2025-07-04 ]

so,

$$|\lambda_{i+1} - \lambda_i| < \left(\frac{\lambda_1}{2+\lambda_1}\right)^{2(i-1)} \frac{\lambda_1(\lambda_1^2 - 1)}{2-\lambda_1^2}.$$

For the condition  $|\lambda_{i+1} - \lambda_i| \leq \epsilon$  to be satisfactory, it suffices to take,

$$\left(\frac{\lambda_1}{2+\lambda_1}\right)^{2(i-1)}\frac{\lambda_1(\lambda_1^2-1)}{2-\lambda_1^2} \leq \epsilon,$$

that is to say

$$\left(\frac{\lambda_1}{2+\lambda_1}\right)^{2(i-1)} \le \frac{\epsilon}{\frac{\lambda_1(\lambda_1^2-1)}{2-\lambda_1^2}}.$$
(3.6)

Using the function Ln in (3.6) and the fact that

$$(i-1)ln\left(\left(\frac{\lambda_1}{2+\lambda_1}\right)^2\right) \le ln\left(\frac{\epsilon(2-\lambda_1^2)}{\lambda_1(\lambda_1^2-1)}\right),$$

 $\mathbf{so},$ 

$$0 < \left(\frac{\lambda_1}{2 + \lambda_1}\right)^2 < 1,$$

(3.6) becomes

$$i \ge \frac{\ln\left(\frac{\epsilon(2-\lambda_1^2)}{\lambda_1^3 - \lambda_1}\right)}{\ln\left(\frac{\lambda_1}{2+\lambda_1}\right)^2} + 1.$$

So  $|\lambda_{i+1} - \lambda_i| < \epsilon$ , starting from

$$N_0 = \left[\frac{ln\left(\frac{\epsilon(2-\lambda_1^2)}{\lambda_1^3 - \lambda_1}\right)}{ln\left(\frac{\lambda_1}{2+\lambda_1}\right)^2}\right] + 1.$$

where [x] is the floor function of x.

In practice, we take  $\epsilon = 10^{-N}$ , we then show the following result:

## Corollary 3.3. Once

$$i \ge \left[\frac{-Nln(10) + ln\left(\frac{(2-\lambda_1^2)}{\lambda_1^3 - \lambda_1}\right)}{ln\left(\frac{\lambda_1}{2+\lambda_1}\right)^2}\right] + 1,$$
(3.7)

we have

$$|\lambda_{i+1} - \lambda_i| < 10^{-N}$$

where [x] is the floor function of x.

3.1.1. Idea on some estimates of  $N_0$ . For

$$\begin{split} m &= 1000, \\ \alpha &= <\varphi_1, \varphi_2 >= 21,47547348, \\ \beta &= <\varphi_1, \varphi_1 >= 42,1948008, \\ \lambda_1 &= -\frac{\alpha}{\beta} = -050896018165347, \\ N_0 &= \left[\frac{-Nln(10) + ln\left(\frac{(2-\lambda_1^2)}{\lambda_1^3 - \lambda_1}\right)}{ln\left(\frac{\lambda_1}{2 + \lambda_1}\right)^2}\right] + 1. \end{split}$$

Precision $10^{-N}$	iteration number $N_0$
$1 \le N \le 10$	N
$11 \le N \le 24$	N+1
$25 \le N \le 38$	N+2
$39 \le N \le 52$	N+3
$53 \le N \le 66$	N+4
$67 \le N \le 80$	N+5
$81 \le N \le 94$	N+6
$95 \le N \le 108$	N+7
$109 \le N \le 122$	N+8
$123 \le N \le 136$	N+9

TABLE 2. Examples the calculation of  $N_0$ : to reduce the iteration number.

## **Corollary 3.4.** $(\lambda_i)$ is stationary from $N_0$ .

*Proof.* if  $i \ge N_0$  we have

$$|\lambda_{i+1} - \lambda_i| < 10^{-N},$$

 $\mathbf{SO}$ 

$$\lambda_{N_0+p} - \lambda_{N_0}| < |\lambda_{N_0+p} - \lambda_{N_0+p-1}| + ... + |\lambda_{N_0+1} - \lambda_{N_0}|$$

Therefore,

$$|\lambda_{N_0+p} - \lambda_{N_0}| < p10^{-N},$$

where,  $p = m - N_0$ .

EXAMPLE 3.5. For  $m = 10^3$ ,  $N = N_0 = 10$ ,

$$|\lambda_{N_0+p} - \lambda_{N_0}| < 0,999.10^{-7} < 10^{-7}.$$

3.2. Calculation of  $r_i$ .

The theorem makes it possible to calculate the values of the roots  $r_i$ .

**Theorem 3.6.** For i = 2, ..., m+1 et k = 1, ..., i-1 if we put  $r_k = x_k+h.X_k$ , then,  $X_k$  is the solution of ,

$$\Lambda_k X^4 - 2\Lambda_k X^3 - \Lambda_k X^2 + 2(\Lambda_k - 1)X + 1 = 0, \qquad (3.8)$$

where:  $X_k \in \left]0, \frac{1}{2}\right[$  and  $\Lambda_k = Ln(-\lambda_k)$ .

Remark 3.7.

• The fact that the function:

$$F(X) = \Lambda_k X^4 - 2\Lambda_k X^3 - \Lambda_k X^2 + 2(\Lambda_k - 1)X + 1,$$

is strictly decreasing and concave on  $[0, \frac{1}{2}]$ , then any numerical approximation method of the root of F is valid.

• The fact that  $\Lambda_k = Ln(-\lambda_k)$  and since, from  $N_0$  the sequence  $\lambda_i$  is considered as a stationary sequence  $(|\lambda_{i+1} - \lambda_i| < 10^{-N_0})$ , then all the functions F(X) are the same for  $k \ge N_0$ .

All the calculations that were done with the previous algorithm work with a very large number of nodes m (m = 10000) which gave very large tables (table1) as a database to do the following.

Now, we will reduce the number of calculations: from m to  $N_0$  depending on the precision required.

## 4. $\delta$ -ZITI ALGORITHM.

We give ourselves a precision  $\epsilon = 10^{-N}$  ( $\epsilon$  or N), We take the step of subdivision h of the interval [a,b], (a,b,m), We compute

$$\begin{array}{rcl}
\alpha &=& <\varphi_{1},\varphi_{2}>, \\
\beta &=& <\varphi_{1},\varphi_{1}>, \\
\lambda_{1} &=& -\frac{\alpha}{\beta}, \\
N_{0} &=& \left[\frac{-Nln(10)+ln\left(\frac{(2-\lambda_{1}^{2})}{\lambda_{1}^{2}-\lambda_{1}}\right)}{ln\left(\frac{\lambda_{1}}{2+\lambda_{1}}\right)^{2}}\right]+1. \\
\left\{\begin{array}{rcl}
For \ i=1, N_{0}-1 \\
\lambda_{i+1} &=& \frac{\lambda_{1}}{2-\lambda_{1}\lambda_{i}}. \\
End \ i
\end{array}\right.$$
(4.1)

For  $i \geq N_0 + 1$ 

$$\lambda_i \simeq \lambda_{N_0}$$

L. Bsiss, C. Ziti

$$\begin{cases} For \ k=1:N_0\\ Calculation \ of \ the \ root \ X_k \ of \ the \ function\\ F(X)=AX^4-2AX^3-AX^2+2(A-1)X+1, \ on \ \big]0, \frac{1}{2}\big[\\ where \ A=ln(-\lambda_k),\\ r_k=x_k+h.X_k. \end{cases}$$
end k  
For  $k\geq N_0+1\\ X_k=X_{N_0},\\ r_k=x_k+hX_k, \ r_k=r_{k-1}+h. \end{cases}$ 

## 4.1. Application.

We consider the following Cauchy problem:

$$\begin{cases} x^2 \cdot u'(x) = 4 \cdot u(x) \ x \in [-4, 4], & (1) \\ u(-4) = 1, & (2) \end{cases}$$
(4.2)

According to (2.19) the following algorithm has been obtained:

$$\begin{cases} \alpha_1 = \frac{1}{\psi_1(r_1)} \\ \alpha_{k+1} = \frac{\alpha_k \cdot \psi_k(r_k)(4(r_{k+1} - r_k) + r_k^2)}{r_k^2 \cdot \psi_{k+1}(r_{k+1})} \text{ for } k = 1, \dots, m. \end{cases}$$

$$(4.3)$$

This algorithm of the  $\delta$ -ziti method that we have to build aims to calculate  $\lambda_i$  and roots  $r_i$ .

m = 1000, a = -4, b = 4.precision  $\epsilon = 10^{-N}$  with N = 8. (from table2 we have  $N_0 = N = 8$ ) We calculate

## Calculation of $\lambda_i$ . Calculation of $r_i$ .

For

For example, the secant method is used to solve:

$$\begin{aligned} k &= 1: N_0 = 8 \\ Calculation of the root X_k of the function \\ F(X) &= AX^4 - 2AX^3 - AX^2 + 2(A-1)X + 1, \text{ on } \left]0, \frac{1}{2}\right| \\ where A &= ln(-\lambda_k), \\ r_k &= x_k + h.X_k. \end{aligned}$$

$\lambda_1$	-0.508960181653470	
$\lambda_2$	-0.292344636312607	
$\lambda_3$	-0.274934054384423	
$\lambda_4$	-0.273624280111297	
$\lambda_5$	-0.273526252120255	
$\lambda_6$	-0.273518918194867	
$\lambda_7$	-0.273518369525952	
$\lambda_8$	-0.273518328478775	
$if N > N_0, \ \lambda_N = \lambda_{N_0}$	-0.273518325407943	

An Optimal Algorithm for the  $\delta$ -ziti Method to Solve Some Mathematical Problems 127

TABLE 3. Results using the optimal algorithm.

$X_1$	0.333258836280244
$X_2$	0.248728737840737
$X_3$	0.241680753511322
$X_4$	0.241148006382392
$X_5$	0.241108118629783
$X_6$	0.241105134358228
$X_7$	0.241104911097089
$X_8$	0.241104894394410
$if N > N_0, \ X_N = X_{N_0}$	0.241104894394410

TABLE 4. Results using the optimal algorithm.

The two graphs in Fig.10 present the comparison between the exact and approximate solution of (4.2).

L. Bsiss, C. Ziti

$r_1 = x_1 + hX_1$	-3.997333929309758
$r_2 = x_2 + hX_2$	-3.990010170097274
$r_3 = x_3 + hX_3$	-3.982066553971909
$r_4 = x_4 + hX_4$	-3.974070815948941
$r_5 = x_5 + hX_5$	-3.966071135050962
$r_6 = x_6 + hX_6$	-3.958071158925134
$r_7 = x_7 + hX_7$	-3.950071160711223
$r_8 = x_8 + hX_8$	-3.942071160844845
$if N > N_0, r_N = r_{N_0} + h$	
$r_{m+1} = b$	4

TABLE 5. Results using the optimal algorithm.



FIGURE 10. Comparison the approximate solution obtained by  $\delta$ -ziti's scheme with the exact solution in presence a singularity.

## 5. Conclusion

The method has shown its effectiveness in comparison with other results (exact or obtained with classical methods). In this work, we have established two algorithms that have resulted in the same results. Table 1 obtained by the first algorithm required as many iterations as number of nodes (likely to be very large, for example  $10^6$ ).

Table 2 obtained by the second algorithm deduces from a careful study (reformulation of  $\lambda_i$ ,  $\psi_i$  and  $r_i$ ) to reduce the number of iterations with an acceptable stopping test for example for a test of  $10^{-5}$  on number of iterations is reduced to 8 iterations.

#### Acknowledgments

The authors thank the referees for their valuable comments.

#### References

- L. Bsiss, C. Ziti, A New Approximation (Ziti's δ-scheme) of the Entropic (Admissible) Solution of the Hyperbolic Problems in One and Several Dimensions, Applications to Convection, Burgers, Gas dynamics and Some Biological Problems, *Turkish Journal of* Analysis and Number Theory, 4(4), (2016), 98-108.
- L. Bsiss, C. Ziti, The δ-ziti's Method to Detect the Blow-up in Finite Time in Some Models of Chemotaxis, *Ponte Journal*, 73(2), (2017), 245-260.
- L. Bsiss, C. Ziti, A New Numerical Method for the Integral Approximation and Solving the Differential Problems: Non-oscillating Scheme, Detecting the Singularity in one and Several Dimension, *Ponte Journal*, **73**(6), (2017), 126-172.
- L. Bsiss, C. Ziti, A New Entropic Riemann Solver of Conservation Law of Mixed Type Including Ziti's δ-Method with some Experimental Tests, Applied and Computational Mathematics Journal, 6(5), (2017), 222-232.
- G. Dhatt, G. Touzot, Une Présentation de la Méthode Des éléments Finis, MALOINE S.A Editeur Paris et les Presses de l'université Laval Quebec, 1981.
- B. Meyer, C. Baudoin, Méthodes de Programmation, Collection de la direction des Etudes et Recherches d'Elictricit de France Edition Eyrolles, 1984.
- B. Demidovitch, T. Maron, *Eléments de Calcul Numérique*, Edition MIR de MOSCOU, 1979.
- 8. M. Sibony, J-CI. Mardon, Analyse Numérique I et II, HERMANN, Editions des sciences et des Arts, 1984.