

A Necessary Condition for Zero Divisors in Complex Group Algebra of Torsion-Free Groups

Alireza Abdollahi^a, Meisam Soleimani Malekan^{b*}

^aDepartment of Pure Mathematics, Faculty of Mathematics and Statistics,
University of Isfahan, Isfahan, Iran 81746-73441

^bDepartment of Mathematics, Faculty of Science, Shahid Rajaee Teacher
Training University, Tehran, 16785-136, Iran

E-mail: a.abdollahi@math.ui.ac.ir

E-mail: msmalekan@gmail.com

ABSTRACT. It is proved that if $\sum_{g \in G} a_g g$ is a non-zero zero divisor element of the complex group algebra $\mathbb{C}G$ of a torsion-free group G then $2 \sum_{g \in G} |a_g|^2 < (\sum_{g \in G} |a_g|)^2$.

Keywords: Hilbert space $\ell^2(G)$, Complex group algebras, Zero divisor conjecture, Torsion-free groups.

2020 Mathematics subject classification: 46C07, 46L10, 20C07, 16S34.

1. INTRODUCTION AND RESULTS

Let G be any group and $\mathbb{C}G$ be the complex group algebra of G , i.e. the set of finitely supported complex functions on G . We may represent an element α in $\mathbb{C}G$ as a formal sum $\sum_{g \in G} a_g g$, where $a_g \in \mathbb{C}$ is the value of α in g . The multiplication in $\mathbb{C}G$ is defined by

$$\alpha\beta = \sum_{g,h \in G} a_g b_h gh = \sum_{g \in G} \left(\sum_{x \in G} a_{gx^{-1}} b_x \right) x$$

*Corresponding Author

for $\alpha = \sum_{g \in G} a_g g$ and $\beta = \sum_{g \in G} b_g g$ in $\mathbb{C}G$. We shall say that α is a *zero divisor* if there exists $0 \neq \beta \in \mathbb{C}G$ such that $\alpha\beta = 0$. If there is a non-zero $\beta \in \ell^2(G)$ such that $\alpha\beta = 0$, then we may say that α is *analytical zero divisor*. If $\alpha\beta \neq 0$ for all $0 \neq \beta \in \mathbb{C}G$, then we say that α is *regular*. The following conjecture is called the *zero divisor conjecture*.

Conjecture 1.1. Let G be a torsion-free group. Then all elements in $\mathbb{C}G$ are regular.

Conjecture 1.1 is still open; it has been proven affirmative when G belongs to special classes of groups; ordered groups ([11] and [12]), supersolvable groups [6], polycyclic-by-finite groups ([1] and [5]) and unique product groups [2]. Delzant [3] deals with group rings of word-hyperbolic groups and proves the conjecture for certain word-hyperbolic groups. Let \mathcal{C} be the smallest class of groups which contains all free groups and is closed under directed unions and extensions with elementary amenable quotients. Let G be a torsion-free group in \mathcal{C} then G satisfies Conjecture 1.1 [8].

The map $\langle \cdot, \cdot \rangle : \mathbb{C}G \times \mathbb{C}G \rightarrow \mathbb{R}$ defined by

$$\langle \alpha, \beta \rangle := \sum_{g \in G} a_g \bar{b}_g \quad (\alpha, \beta \in \mathbb{C}G)$$

is an inner product on $\mathbb{C}G$, so $\|\alpha\|_2 = \langle \alpha, \alpha \rangle^{\frac{1}{2}}$ becomes a norm, called 2-norm; the completion of $\mathbb{C}G$ w.r.t. 2-norm is the Hilbert space $\ell^2(G)$. Indeed, we have

$$\ell^2(G) = \left\{ \alpha : G \rightarrow \mathbb{C} : \sum_{g \in G} \|\alpha(g)\|^2 < \infty \right\}.$$

In [9], Linnell formulated an analytic version of the zero divisor conjecture.

Conjecture 1.2. Let G be a torsion-free group. If $0 \neq \alpha \in \mathbb{C}G$ and $0 \neq \beta \in \ell^2(G)$, then $\alpha\beta \neq 0$.

In [7], it is shown that Since $\mathbb{C}G \subset \ell^2(G)$, the second conjecture implies the first one. In [4], it is proved that for finitely generated amenable groups, the two conjectures are actually equivalent. We prove this is true for all amenable torsion-free groups.

The so-called 1-norm is defined on $\mathbb{C}G$ by

$$\|\alpha\|_1 = \sum_{g \in G} |a_g|, \quad \text{for } \alpha = \sum_{g \in G} a_g g \text{ in } \mathbb{C}G.$$

The *adjoint* of an element $\alpha = \sum_{g \in G} a_g g$ in $\mathbb{C}G$, denoted by α^* , is $\alpha^* = \sum_{g \in G} \bar{a}_g g^{-1}$. We call an element $\alpha \in \mathbb{C}G$ *self-adjoint* if $\alpha^* = \alpha$, and use $(\mathbb{C}G)_s$ to denote the set of self-adjoint elements of $\mathbb{C}G$. It is worthy of mention that

if $\alpha = \sum_{g \in G} a_g g$ is self-adjoint then a_1 should be a real number. For $\alpha \in \mathbb{C}G$, β and γ in $\ell^2(G)$, the following equalities hold:

$$\langle \alpha\beta, \gamma \rangle = \langle \beta, \alpha^*\gamma \rangle.$$

The goal of this paper is to give a criterion for an element in a complex group algebra to be regular:

Theorem 1.1. *Let G be a torsion free group. Then $\alpha \in \mathbb{C}G$ is regular if $2\|\alpha\|_2^2 \geq \|\alpha\|_1^2$.*

2. PRELIMINARIES

In this section we provide some preliminaries needed in the following.

Let G be a group. The *support* of an element $\alpha = \sum_{g \in G} a_g g$ in $\mathbb{C}G$, $\text{supp}(\alpha)$, is the finite subset $\{g \in G : a_g \neq 0\}$ of G .

Let H be a subgroup of G , and T be a right transversal for H in G . Then every element $\alpha \in \mathbb{C}G$ (resp. $\alpha \in \ell^2(G)$) can be written uniquely as a finite sum of the form $\sum_{t \in T} \alpha_t t$ with $\alpha_t \in \mathbb{C}H$ (resp. $\alpha_t \in \ell^2(H)$).

For $S \subset G$, we denote by $\langle S \rangle$, the subgroup of G generated by S . We have the following key lemma:

Lemma 2.1. *Let G be a group, $\alpha \in \mathbb{C}G$ and $H = \langle \text{supp}(\alpha) \rangle$. Then α is regular in $\mathbb{C}G$ iff α is regular in $\mathbb{C}H$.*

Proof. Suppose that α is a zero divisor. Among elements $0 \neq \gamma$ in $\mathbb{C}G$ which satisfy $\alpha\gamma = 0$ consider an element β such that $1 \in \text{supp}(\beta)$ and $|\text{supp}(\beta)|$ is minimal, then one can easily show that $\beta \in \mathbb{C}H$, and this proves the result of the lemma. \square

An immediate consequence of this lemma is:

Corollary 2.2. *A group G satisfies Conjecture 1.1 iff all its finitely generated subgroups satisfy the Conjecture 1.1.*

By Lemma 2.1 in hand, we can generalize the main theorem of [4]:

Theorem 2.3. *Let G be an amenable group. If $0 \neq \alpha \in \mathbb{C}G$, $0 \neq \beta \in \ell^2(G)$ and $\alpha\beta = 0$, then there exists $0 \neq \gamma \in \mathbb{C}G$ such that $\alpha\gamma = 0$.*

The above theorem along with results in [10] provides another proof for [7, Theorem 2].

For a normal subgroup N of a group G , we denote the natural quotient map by $q_N : G \rightarrow G/N$. We continue to show that:

Lemma 2.4. *Let N be a normal subgroup of a group G satisfying Conjecture 1.1. Consider a non-torsion element $q_N(t)$, $t \in G$, in the quotient group. Then $\alpha + \beta t$ is regular, for all $\alpha, \beta \in \mathbb{C}N \setminus \{0\}$.*

Proof. Suppose that $\alpha + \beta t$ is a zero divisor for non zero elements $\alpha, \beta \in \mathbb{C}N$. Applying Lemma 2.1 and multiplying by a suitable power of t , we can assume that there are non zero elements γ_k , $k = 0, 1, \dots, n$, such that

$$(\alpha + \beta t) \sum_{k=0}^n \gamma_k t^k = 0.$$

In particular, $0 = \beta t \gamma t^n = (\beta t \gamma_n t^{-1}) t^{n+1}$, whence $\beta t \gamma_n t^{-1} = 0$, a contradiction, because $t \gamma_n t^{-1}$ is a non zero element of $\mathbb{C}N$. \square

Proposition 2.5. *Let N be an amenable normal subgroup of a group G satisfying Conjecture 1.1. Consider a non-torsion element $q_N(t)$, $t \in G$, in the quotient group. Then there is no $0 \neq \gamma \in \ell^2(G)$ such that $(\alpha + \beta t)\gamma = 0$. In particular, $a + bg$ is an analytical zero divisor, for all non-torsion element $g \in G$ and non zero complex numbers a, b .*

Proof. The group $\langle N, t \rangle$ is amenable. Hence Lemma 2.4 together with Theorem 2.3 yields the result. \square

3. A CONE OF REGULAR ELEMENTS

The result of the Proposition 2.5 is true if we replace \mathbb{C} by an arbitrary field \mathbb{F} . The field of complex numbers allows us to define inner product on the group algebra; with the help of inner product, we can construct new regular elements from the ones we have:

Proposition 3.1. *Let G be a group and \mathcal{F} be a finite non-empty subset of $\mathbb{C}G$. If $\sum_{\alpha \in \mathcal{F}} \alpha^* \alpha$ is an analytical zero divisor then all elements of \mathcal{F} are analytical zero divisors. In particular, $\alpha \in \mathbb{C}G$ is an analytical zero divisor if and only if $\alpha^* \alpha$ is an analytical zero divisor.*

Proof. Let $\tilde{\alpha} := \sum_{\alpha \in \mathcal{F}} \alpha^* \alpha$ and $\tilde{\alpha}\beta = 0$ for some $\beta \in \ell^2(G)$. Then

$$0 = \langle \tilde{\alpha}\beta, \beta \rangle = \sum_{\alpha \in \mathcal{F}} \langle \alpha^* \alpha \beta, \beta \rangle = \sum_{\alpha \in \mathcal{F}} \langle \alpha \beta, \alpha \beta \rangle = \sum_{\alpha \in \mathcal{F}} \|\alpha \beta\|_2^2,$$

whence $\alpha \beta = 0$ for all $\alpha \in \mathcal{F}$. This completes the proof. \square

A *cone* in a vector space \mathfrak{X} is a subset \mathfrak{K} of \mathfrak{X} such that $\mathfrak{K} + \mathfrak{K} \subset \mathfrak{K}$ and $\mathbb{R}_+ \mathfrak{K} \subset \mathfrak{K}$. We proceed by introducing a cone of regular elements in $\mathbb{C}G$. First a definition:

Definition 3.2. Let G be a group and $(\mathbb{C}G)_s$ be the set of self adjoint elements $\alpha \in \mathbb{C}G$, we define a function $\Upsilon : (\mathbb{C}G)_s \rightarrow \mathbb{R}$ by

$$\Upsilon(\alpha) := a_1 - \sum_{g \neq 1} |a_g|.$$

We call an element $\alpha \in (\mathbb{C}G)_s$ **golden** if $\Upsilon(\alpha) \geq 0$. The set of all golden elements in $(\mathbb{C}G)_s$ is denoted by $(\mathbb{C}G)_{\text{gold}}$.

What is important about golden elements is:

Proposition 3.3. *For a torsion free group G , $(\mathbb{C}G)_{gold}$ is a cone of regular elements.*

Proof. It is obvious that if α is golden then so is $r\alpha$ for any $r > 0$. The triangle inequality for \mathbb{C} shows that if α and γ are golden then so is $\alpha + \gamma$. For $\alpha \in (\mathbb{C}G)_s$, we have

$$\begin{aligned}\alpha &= \frac{1}{2}(\alpha + \alpha^*) \\ &= a_1 + \frac{1}{2} \sum_{g \neq 1} (\bar{a}_g g^{-1} + a_g g) \\ &= \Upsilon(\alpha) + \frac{1}{2} \sum_{g \neq 1} (2|a_g| + \bar{a}_g g^{-1} + a_g g) \\ &= \Upsilon(\alpha) + \frac{1}{2} \sum_{g \neq 1} |a_g| \left(\frac{\bar{a}_g}{|a_g|} + g \right)^* \left(\frac{a_g}{|a_g|} + g \right)\end{aligned}$$

Hence, by Lemma 2.5 and Proposition 3.1, α is regular. \square

Now, we are ready to prove our main result:

Proof of Theorem 1.1. For $\alpha = \sum_{g \in G} a_g g$ in $\mathbb{C}G$, $\alpha^* \alpha$ is self-adjoint, and one can easily show that

$$\Upsilon(\alpha^* \alpha) \geq 2\|\alpha\|_2^2 - \|\alpha\|_1^2.$$

Hence, by Proposition 3.3, the result of the Theorem is proved. \square

ACKNOWLEDGMENTS

The author wish to thank the referee for the helpful suggestions and comments.

REFERENCES

1. K. A. Brown, On Zero Divisors in Group Rings, *Bull. Lond. Math. Soc.*, **8**, (1976), 251-256.
2. J. M. Cohen, Zero Divisors in Group Rings, *Comm. Algebra*, **2**, (1974), 1-14.
3. T. Delzant, Sur l'anneau Dun Groupe Hyperbolique, *C. R. Acad. Sci. Paris Ser. I Math.*, **324**(4), (1997), 381-384.
4. G. Elek, On the Analytic Zero Divisor Conjecture of Linnell, *Bulletin of the London Mathematical Society*, **35**(2), (2003), 236-238.
5. D. R. Farkas, R. L. Snider, K_0 and Noetherian Group Rings, *J. Algebra*, **42**, (1976), 192-198.
6. E. Formanek, The Zero Divisor Question for Supersolvable Groups, *Bull. Aust. Math. Soc.*, **73c**, (1973), 67-71.
7. P. A. Linnell, Zero Divisors and Group von Neumann Algebras, *Pacific J. Math.*, **149**, (1991), 349-363.
8. P. A. Linnell, Division Rings and Group von Neumann Algebras, *Forum Math.*, **5**(6), (1993), 561-576.

9. P. A. Linnell, *Analytic Versions of the Zero Divisor Conjecture*, In Geometry and cohomology in group theory (Durham 1994), 252. London Math. Soc. Lecture Note, 1998.
10. P. A. Linnell, P. H. Kropholler, J. A. Moody, Applications of a New K -theoretic Theorem to Soluble Group Rings, *Proc. Amer. Math. Soc.*, **104**, (1988), 675-684.
11. A. I. Malcev, On Embedding of Group Algebras in a Division Algebra (in russian), *Dokl. Akad. Nauk.*, **60**, 1948, 1499-1501.
12. B. H. Neumann, On Ordered Division Rings, *Trans. Amer. Math. Soc.*, **66**, (1949), 202-252.