

## Labeling Subgraph Embeddings and Cordiality of Graphs

Zhen-Bin Gao<sup>a</sup>, Ruo-Yuan Han<sup>a</sup>, Sin-Min Lee<sup>b</sup>,  
 Hong-Nan Ren<sup>a</sup>, Gee-Choon Lau<sup>c,\*</sup>

<sup>a</sup>College of Science, Harbin Engineering University, Harbin,  
 150001, P. R. China.

<sup>b</sup>1403, North First Avenue, Upland, CA 91786, USA.

<sup>c</sup>Faculty of Computer and Mathematical Sciences,  
 Universiti Teknologi MARA (Segamat Campus), 85000 Johor, Malaysia.

E-mail: gaozhenbin@aliyun.com

E-mail: 3213358692@qq.com

E-mail: sinminlee@gmail.com

E-mail: 1114912080@qq.com

E-mail: geeclau@yahoo.com

**ABSTRACT.** Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ , a vertex labeling  $f : V(G) \rightarrow \mathbb{Z}_2$  induces an edge labeling  $f^+ : E(G) \rightarrow \mathbb{Z}_2$  defined by  $f^+(xy) = f(x) + f(y)$ , for each edge  $xy \in E(G)$ . For each  $i \in \mathbb{Z}_2$ , let  $v_f(i) = |\{u \in V(G) : f(u) = i\}|$  and  $e_{f^+}(i) = |\{xy \in E(G) : f^+(xy) = i\}|$ . A vertex labeling  $f$  of a graph  $G$  is said to be friendly if  $|v_f(1) - v_f(0)| \leq 1$ . The friendly index set of the graph  $G$ , denoted by  $FI(G)$ , is defined as  $\{|e_{f^+}(1) - e_{f^+}(0)| : \text{the vertex labeling } f \text{ is friendly}\}$ . The full friendly index set of the graph  $G$ , denoted by  $FFI(G)$ , is defined as  $\{e_{f^+}(1) - e_{f^+}(0) : \text{the vertex labeling } f \text{ is friendly}\}$ . A graph  $G$  is cordial if  $-1, 0$  or  $1 \in FFI(G)$ . In this paper, by introducing labeling subgraph embeddings method, we determine the cordiality of a family of cubic graphs which are double-edge blow-up of  $P_2 \times P_n, n \geq 2$ . Consequently, we completely determined friendly index and full product cordial index sets of this family of graphs.

**Keywords:** Vertex labeling, Full friendly index set, Cordiality,  $P_2$ -embeddings,  $C_4$ -embeddings.

**2000 Mathematics Subject Classification:** 05C78, 05C25.

---

\*Corresponding Author

Received 05 July 2016; Accepted 03 March 2018

©2019 Academic Center for Education, Culture and Research TMU

## 1. INTRODUCTION

Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . Let  $A$  be an abelian group. A labeling  $f : V(G) \rightarrow A$  induces an edge labeling  $f^+ : E(G) \rightarrow A$  defined by  $f^+(x, y) = f(x) + f(y)$ , for each edge  $(x, y) \in E(G)$ . For  $i \in A$ , let  $v_f(i) = |\{v \in V(G) : f(v) = i\}|$  and  $e_{f^+}(i) = |\{e \in E(G) : f^+(e) = i\}|$ . Let  $c(f) = \{|e_{f^+}(i) - e_{f^+}(j)| : (i, j) \in A \times A\}$ . A labeling  $f$  of a graph  $G$  is said to be  $A$ -friendly if  $|v_f(i) - v_f(j)| \leq 1$  for all  $(i, j) \in A \times A$ . If  $c(f)$  is  $(i, j)$ -matrix for some  $A$ -friendly labeling  $f$ , then  $f$  is said to be  $A$ -cordial. The notion of  $A$ -cordial labelings was first introduced by Hovey [5], who generalized the concept of cordial graphs of Cahit [1].

In this paper, we will exclusively focus on  $A = \mathbb{Z}_2$ , and drop the reference to the group. A vertex  $v$  is called a  $k$ -vertex if  $f(v) = k$ ,  $k \in \{0, 1\}$ , an edge  $e$  is called a  $k$ -edge if  $f^+(e) = k$ ,  $k \in \{0, 1\}$ . A vertex labeling  $f$  of a graph  $G$  is said to be friendly if  $|v_f(1) - v_f(0)| \leq 1$ .

In [4] the following concept was introduced.

**Definition 1.1.** The *friendly index set*  $FI(G)$  of a graph  $G$  is defined as  $\{|e_{f^+}(1) - e_{f^+}(0)| : \text{the vertex labeling } f \text{ is friendly}\}$ .

The following result was established in [6]:

**Theorem 1.2.** For any graph  $G$  with  $q$  edges, the friendly index set  $FI(G) \subseteq \{0, 2, \dots, q\}$  if  $q$  is even, and  $FI(G) \subseteq \{1, 3, \dots, q\}$  if  $q$  is odd.

For more details of known results and open problems on friendly index sets, the reader can see [7, 8, 9].

Shiu and Kwong [12] extended  $FI(G)$  to  $FFI(G)$ .

**Definition 1.3.** The *full friendly index set*  $FFI(G)$  of a graph  $G$  is defined as  $\{e_{f^+}(1) - e_{f^+}(0) : \text{the vertex labeling } f \text{ is friendly}\}$ .

Hence, a graph  $G$  is cordial if  $-1, 0$  or  $1 \in FFI(G)$ . Moreover, the cordiality of  $G$  can be determined by finding the  $FI(G)$  or  $FFI(G)$ . Shiu and Kwong [12] determined  $FFI(P_2 \times P_n)$ . Shiu and Lee [13] determined the full friendly index sets of twisted cylinders. Shiu and Wong [15] determined the full friendly index sets of cylinder graphs. Shiu and Ho [10] determined the full friendly index sets of some permutation Petersen graphs, they also determined the full friendly index sets of slender and flat cylinder graphs [11]. Shiu and Ling [14] determined the full friendly index sets of Cartesian products of two cycles. Sinha and Kaur [16] studied the full friendly index sets of some graphs such as  $K_n$ ,  $C_n$ , fans  $F_n$ ,  $F_{2,m}$  and  $P_3 \times P_n$ . Interested readers may refer to [2] for more results on cordiality of graphs. In general, it is difficult to obtain the full friendly index sets of graphs. The problem on the full friendly index sets and cordiality of general cubic graphs is still beyond our reach at this moment.

**Definition 1.4.** Let  $G$  and  $H$  be two graphs such that  $u$  and  $v$  are two particular vertices of  $H$ . An edge  $xy$  of  $G$  is *blown-up* by  $H$  at  $u$  and  $v$  if  $xy$  is replaced by  $H$  by identifying  $x$  and  $u$ , and  $y$  and  $v$  respectively.

**Definition 1.5.** Let  $P_2 \times P_n$  ( $n \geq 2$ ) be the ladder graph of order  $2n$  and size  $3n-2$  with vertex set  $V = \{u_i, v_i : 1 \leq i \leq n\}$  and edge set  $E = \{u_i u_{i+1}, v_i v_{i+1} : 1 \leq i \leq n-1\} \cup \{u_i v_i : 1 \leq i \leq n\}$ . Let  $K_4^- = K(1, 1, 2)$  be the complete tripartite graph with degree 2 vertices  $u, v$ . The *double-edge blow-up* graph of  $P_2 \times P_n$ , denoted  $DB(n-2)$ , is obtained by blowing up the edges  $u_1 v_1$  and  $u_n v_n$  by a  $K_4^-$  at  $u$  and  $v$  respectively such that a cubic graph is obtained.

In what follow, we let  $m = n - 2 \geq 0$  so that  $|V(DB(m))| = 2(m+4)$ ,  $|E(DB(m))| = 3m+12$ .

EXAMPLE 1.6. The graph  $DB(2)$  is illustrated in Figure 1.




FIGURE 1. Graph  $DB(2)$

In this paper, we introduce a labeling subgraph embeddings method to obtain the full friendly index sets of  $DB(m)$ . Consequently, the cordiality of  $DB(m)$  is determined.

## 2. PRELIMINARIES

We now present some derived results and prove some results which will be used to obtain our main results.


**Theorem 2.1.** (Shiu and Wong[15]) *Let  $f$  be a labeling of a graph  $G$  that contains a cycle  $C$  as its subgraph. If  $C$  contains a 1-edge, then the number of 1-edges in  $C$  is a positive even number.*

**Theorem 2.2.** *In any friendly labeling  $f$  of  $DB(m)$ , if any two vertex labels are exchanged, then  $e_{f+}(1)$  changes by  $-6, -4, -2, 0, 2, 4$ , or  $6$ .*

*Proof.* Since the graph  $DB(m)$  is cubic, any vertex  $u$  is adjacent to three vertices  $u_1, u_2$ , and  $u_3$ . In a friendly labeling of  $DB(m)$ , suppose that the vertices  $u, u_1, u_2$ , and  $u_3$  are labeled by  $x, x_1, x_2$ , and  $x_3$  ( $x, x_1, x_2, x_3 \in \{0, 1\}$ ) respectively. When we change the label of  $u$  to  $1-x$ , the number of 1-edges changes by  $-3, -1, 1$ , or  $3$ .

For any two vertices of  $u$  and  $v$  in  $DB(m)$ , there are five cases that three of them are listed in Figure 2.

Exchange the labels of  $u$  and  $v$ ,  $e_{f+}(1)$  changes by  $-6, -4, -2, 0, 2, 4$ , or  $6$  in (a), and by  $-4, -2, 0, 2$ , or  $4$  in (b) and (c). In the fourth case,  $u$  and  $v$

FIGURE 2. Three possible structures of  $DB(m)$ 

are the two degree 3 vertices of a  $K_4^-$  subgraph that gives no change to  $e_{f^+}(1)$ . In the fifth case,  $u$  and  $v$  are the two degree 2 vertices of a  $K_4^-$  subgraph with  $e_{f^+}(1)$  changes by 0 or 2. Hence, if any two vertex labels are exchanged, then  $e_{f^+}(1)$  changes by  $-6, -4, -2, 0, 2, 4$ , or  $6$ .  $\square$

**Theorem 2.3.** *If  $f_1$  and  $f_2$  be two friendly labelings of  $DB(m)$  such that  $f_2$  is obtained from  $f_1$  by exchanging two distinct vertex labels under  $f_1$ , then  $(e_{f_1^+}(1) - e_{f_1^+}(0)) - (e_{f_2^+}(1) - e_{f_2^+}(0)) \equiv 0 \pmod{4}$ .*

*Proof.* Since  $|V(DB(m))| = 2(m+4)$ , any friendly labeling  $f_1$  of  $DB(m)$  gives  $v_f(1) = v_f(0)$ . Hence, exchanging two vertex labels under  $f_1$  gives a new friendly labeling  $f_2$ .

Since  $e_{f^+}(1) - e_{f^+}(0) = 2e_{f^+}(1) - |E|$ , we have  $(e_{f_1^+}(1) - e_{f_1^+}(0)) - (e_{f_2^+}(1) - e_{f_2^+}(0)) = 2(e_{f_1^+}(1) - e_{f_2^+}(1)) \equiv 0 \pmod{4}$ , by Theorem 2.2.  $\square$

An edge  $uv$  is called an  $(i, j)$ -edge if it is incident with an  $i$ -vertex and an  $j$ -vertex ( $i, j \in \{0, 1\}$ ). In the following discussions, all numbers are integer. When the context is clear, we shall also drop the subscript  $f, f^+$ . Below are three necessary notations.

**(1). Labeling graph:** A graph  $G$  with a friendly labeling  $f$  such that  $e(1) - e(0) = a$  is called a labeling graph of  $G$ , denoted by  $G(a)$ . For easy reading, the  $P_2 = a_1a_2$  labeling subgraph is denoted by  $\begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$ , and the induced  $C_4 = b_1b_2b_4b_3b_1$  (or induced  $P_4 = b_1b_2b_4b_3$  or  $b_2b_1b_3b_4$ ) labeling subgraph is denoted by  $\begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}$ .

Throughout this paper, unless stated otherwise, every labeling subgraph  $\begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}$  denotes an induced  $C_4$  or an induced  $P_4$  subgraph with vertices as stated above. Let  $G(a)$  be a labeling graph having an edge with end-vertex labels  $b_1$  and  $b_2$  respectively, and another edge with end-vertex labels  $b_3$  and  $b_4$  respectively.

**(2).  $P_2$ -embedding:** A  $P_2$ -embedding onto  $G(a)$  at  $(b_1, b_2)$ -edge and  $(b_3, b_4)$ -edge is obtained by replacing  $uv$  and  $u'v'$  by a length 2 path  $uxv$  and  $u'x'v'$  respectively and embedding an edge  $xx'$  with corresponding end-vertex labels  $a_1$

and  $a_2$  such that a new labeling graph  $G(b)$  with three extra edges is obtained. Such a  $P_2$ -embedding is denoted by

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} + \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}.$$

**(3).  $C_4$ -embedding:** A  $C_4$ -embedding onto  $G(a)$  at  $(b_1, b_2)$ -edge and  $(b_3, b_4)$ -edge is obtained by replacing  $uv$  and  $u'v'$  by a length 3 path  $uxyv$  and  $u'x'y'v'$  respectively and embedding an edge  $xx'$  with corresponding end-vertex labels  $a_1$  and  $a_3$ , and another edge  $yy'$  with corresponding end-vertex labels  $a_2$  and  $a_4$  such that a new labeling graph  $G(b)$  with extra 6 edges is obtained. Such a  $C_4$ - or  $P_4$ -embedding is denoted by

$$\begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} + \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}.$$

EXAMPLE 2.4. In Figure 3, a  $P_2$ -embedding onto  $P_2 \times P_4$  with  $e(1) - e(0) = -6$  gives a labeling graph of  $P_2 \times P_5$  with  $e(1) - e(0) = -7$ . The embedding is denoted by

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}.$$

Similarly, a  $C_4$ -embedding onto the same  $P_2 \times P_4$  gives a labeling graph of  $P_2 \times P_6$  with  $e(1) - e(0) = -4$ . The embedding is denoted by

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}.$$




FIGURE 3. A  $P_2$ - and a  $C_4$ -embedding onto a  $P_2 \times P_4$ , respectively

### 3. FULL FRIENDLY INDEX SETS OF $DB(m)$

In the following discussions, all  $P_2$ - or  $C_4$ -embeddings onto  $DB(m)$  only apply to edges  $u_i u_{i+1}$  and  $v_i v_{i+1}$  for some  $i \in \{0, 1, \dots, n-1\}$ .

**Lemma 3.1.** *For any friendly labeling  $f$  of  $DB(m)$ , we have*

- (1)  $e(1) \leq 10$  if  $m = 0$ ;
- (2)  $e(1) \leq 3m + 8$  if  $m > 0$ .

*Proof.* Given any friendly labeling  $f$  of  $DB(m)$ , it is clear that each  $K_4^-$  subgraph contains at least one 0-edge. Hence,  $e(0) \geq 2$ . Consequently,  $e(1) \leq 10$  if  $m = 0$ . Suppose  $m \geq 1$ . By Theorem 2.1, each  $K_4^-$  subgraph contains at least one 0-edge. In order to get  $\min\{e(0)\}$ , each  $K_4^-$  must contain exactly one 0-edge. Assuming  $m = 1$ . It is easy to verify that  $e(0) \geq 4$ . Assuming  $m \geq 2$  and  $e(0) = 3$ . Without loss of generality, we may assume  $f(u_i) = f(v_j) = x$  for odd  $i$  and even  $j$ , whereas  $f(u_i) = f(v_j) = 1 - x$  for even  $i$  and odd  $j$ . As  $f$  is friendly, the four degree 3 vertices of both  $K_4^-$  must be assigned with two  $x$  and two  $1 - x$ ,  $x \in \{0, 1\}$ . Consequently,  $e(0) = 4$ , which is a contradiction. Hence,  $e(0) \geq 4$  and  $e(1) \leq 3m + 8$  if  $m > 0$ .  $\square$

**Lemma 3.2.** *For any friendly labeling  $f$  of  $DB(m)$ , we have*

- (1)  $e(1) \geq 2$  if  $m \geq 0$  is even;
- (2)  $e(1) \geq 3$  if  $m > 0$  is odd.

*Proof.* Let  $x_1, x_2$  (respectively,  $x_3, x_4$ ) be the 2 common neighbors of  $u_1, v_1$  (respectively,  $u_n, v_n$ ). By Theorem 2.1, any largest induced cycle of  $DB(m)$ , say  $C$ , contains at least two 1-edges. When  $m$  is even,  $e(1) = 2$  can be attained by labeling  $x_1, x_2, u_i$  and  $v_i$  ( $1 \leq i \leq (m+2)/2$ ) by 1 and the remaining vertices by 0. Assume  $m$  is odd. Since  $v(1) = v(0) = m + 4$  and  $C$  has  $2m + 6 \geq 6$  vertices, we have  $|v(1) - v(0)| = 0$  or 2. Note that the two  $K_4^-$  subgraphs may contain 0, 1 or 2 1-edges in  $C$ . In each possibility, we can verify that  $e(1) \geq 3$ . The equality can be attained by labeling  $x_1, x_2, u_i, u_{(m+3)/2}$  and  $v_i$  ( $1 \leq i \leq (m+1)/2$ ) by 1 and the remaining vertices by 0. The lemma holds.  $\square$

**Lemma 3.3.**  $FFI(DB(0)) = \{-4 + 4i : -1 \leq i \leq 3\}$ .

*Proof.* By Theorems 1.2, 2.3 and Lemmas 3.1, 3.2,  $FFI(DB(0)) \subseteq \{-4 + 4i : -1 \leq i \leq 3\}$ . The labeling graphs in Figure 4 show that the equality holds.  $\square$

**Lemma 3.4.** *For even  $m \geq 2$ ,  $FFI(DB(m)) = \{-3m - 8 + 4i : 0 \leq i \leq \frac{3(m+2)}{2}\}$ .*

*Proof.* By Theorems 1.2, 2.3 and Lemmas 3.1, 3.2,  $FFI(DB(m)) \subseteq \{-3m - 8 + 4i : 0 \leq i \leq \frac{3(m+2)}{2}\}$ . We prove the equality holds by induction on  $m$ . Suppose  $m = 2$ . We show that there exist labeling graphs of  $DB(2)(-14 + 4i)$ ,  $0 \leq i \leq 6$ , by doing the following seven  $C_4$ -embeddings:




FIGURE 4. Labeling graphs of  $DB(0)$  with  $e(1) - e(0) \in \{-4 + 4i : -1 \leq i \leq 3\}$

**Case (1).** In  $DB(0)(-8)$ , embed  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 6. Hence, we obtain  $DB(2)(-14)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ .

**Case (2).** In  $DB(0)(-8)$ , embed  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(2)(-10)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Case (3).** In  $DB(0)(-4)$ , embed  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(2)(-6)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Case (4).** In  $DB(0)(0)$ , embed  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(2)(-2)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Case (5).** In  $DB(0)(4)$ , embed  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(2)(2)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Case (6).** In  $DB(0)(8)$ , embed  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(2)(6)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Case (7).** In  $DB(0)(8)$ , embed  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  to increase  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(2)(10)$ , and in this labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

The labeling graphs of  $DB(2)$  that we have obtained are illustrated in Figure 5.




FIGURE 5. Labeling graphs of  $DB(2)$  with  $e(1) - e(0) \in \{-14 + 4i : 0 \leq i \leq 6\}$

Hence,  $FFI(DB(2)) = \{-14 + 4i : 0 \leq i \leq 6\}$ .

Note that in the labeling graphs in Figure 5 (except  $DB(2)(-14)$  and  $DB(2)(10)$ ), there exist the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ . In  $DB(2)(-14)$ , there exists the labeling subgraph  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ . In  $DB(2)(10)$ , there exists the labeling subgraph  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

Since the embedding  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  decreases  $e(1) - e(0)$  by 2, we do this embedding onto  $DB(2)(-14 + 4i)$ ,  $1 \leq i \leq 5$ , to obtain  $DB(4)(-16 + 4i)$ ,  $1 \leq i \leq 5$ , and in these labeling graphs, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

Similarly, we do the embeddings  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  onto  $DB(2)(-14)$  to decrease  $e(1) - e(0)$  by 2 and 6 respectively. Thus, we obtain  $DB(4)(-16)$  and  $DB(4)(-20)$  with the labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ , respectively.

Next, we do the embeddings  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  onto  $DB(2)(10)$  so that  $e(1) - e(0)$  is decreased by 2 and is increased by 6, respectively. Thus, we obtain  $DB(4)(8)$  and  $DB(4)(16)$  with the labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ , respectively.

In  $DB(2)(6)$ , we do the embedding  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  to increase  $e(1) - e(0)$  by 6. Hence, we obtain  $DB(4)(12)$ , and there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

Hence,  $FFI(DB(4)) = \{-20 + 4i : 0 \leq i \leq 9\}$ .

Now, assume that for even  $k \geq 6$ ,  $FFI(DB(k)) = \{-3k - 8 + 4i : 0 \leq i \leq \frac{3(k+2)}{2}\}$  such that the labeling graph  $DB(k)(-3k - 8 + 4i)$ , ( $1 \leq i \leq \frac{3k+4}{2}$ ), has a labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ , and that in  $DB(k)(-3k - 8)$  and  $DB(k)(3k + 4)$ ,

there exist the labeling subgraphs  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ , respectively.

Hence, we do the embedding  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  onto  $DB(k)(-3k - 8 + 4i)$ ,  $1 \leq i \leq \frac{3k+4}{2}$ , to obtain  $DB(k+2)(-3k - 10 + 4i)$ ,  $1 \leq i \leq \frac{3k+4}{2}$ , and in these labeling graphs, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

Similarly, we do the embeddings  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$  onto  $DB(k)(-3k - 8)$  to obtain  $DB(k+2)(-3k - 10)$  and  $DB(k+2)(-3k - 14)$  respectively, and there exist the labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ , respectively.

Next, we do the embeddings  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  onto  $DB(k)(3k + 4)$  such that  $e(1) - e(0)$  is decreased by 2 and is increased by 6 respectively. Thus, we obtain  $DB(k+2)(3k + 2)$  and  $DB(k+2)(3k + 10)$  having the labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ , respectively.

Finally, in  $DB(k)(3k)$ , we do the embedding  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  to increase  $e(1) - e(0)$  by 6. Hence, we obtain  $DB(k+2)(3k + 6)$ , and there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

By mathematical induction, we have  $FFI(DB(k+2)) = \{-3(k+2) - 8 + 4i : 0 \leq i \leq \frac{3(k+4)}{2}\}$ . The proof is complete.  $\square$

By Lemmas 3.3 and 3.4, we have

**Theorem 3.5.** For even  $m$ ,  $FFI(DB(m)) =$

- (1)  $\{-4 + 4i : -1 \leq i \leq 3\}$  if  $m = 0$ ;
- (2)  $\{-3m - 8 + 4i : 0 \leq i \leq \frac{3(m+2)}{2}\}$  if  $m \geq 2$ .

We now consider odd  $m \geq 1$ .

**Theorem 3.6.** For odd  $m \geq 1$ ,  $FFI(DB(m)) = \{-3m - 6 + 4i : 0 \leq i \leq \frac{3m+5}{2}\}$ .

*Proof.* By Theorems 1.2, 2.3 and Lemmas 3.1, 3.2, we have  $FFI(DB(m)) \subseteq \{-3m - 6 + 4i : 0 \leq i \leq \frac{3m+5}{2}\}$ . We prove the equality holds by induction on  $m$ . The labeling graphs in Figures 6 and 7 show that the equality holds for  $m = 1, 3$ .




FIGURE 6. Labeling graphs of  $DB(1)$  with  $e(1) - e(0) \in \{-9 + 4i : 0 \leq i \leq 4\}$




FIGURE 7. Labeling graphs of  $DB(3)$  with  $e(1) - e(0) \in \{-15 + 4i : 0 \leq i \leq 7\}$

Note that in the labeling graphs in Figure 7 (except the labeling graphs  $DB(3)(-15)$  and  $DB(3)(13)$ ), there exist the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  or

$$\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}.$$

Moreover, in  $DB(3)(-15)$ , there exist a  $C_4$ -embedding onto two induced  $P_4$  labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ . In  $DB(3)(-11)$ , there exists a  $C_4$ -embedding onto two induced  $P_4$  labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ . In  $DB(3)(9)$ , there exists the labeling subgraph  $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ . In  $DB(3)(13)$ , there exist two labeling subgraphs  $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$  (which is an induced  $P_4$  subgraph) and  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

Now, assume that for odd  $m = k \geq 5$ ,  $FFI(DB(k)) = \{-3k - 6 + 4i : 0 \leq i \leq \frac{3k+5}{2}\}$  such that the labeling graphs  $DB(k)(-3k - 6 + 4i)$ ,  $1 \leq i \leq \frac{3k+3}{2}$ , has a labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  or  $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$ . Moreover, in  $DB(k)(-3k - 6)$ , there exist two labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ . In  $DB(k)(-3k - 2)$ , there exist two labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ . In  $DB(k)(3k)$ , there exists the labeling subgraph  $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ . In  $DB(k)(3k + 4)$ , there exist two labeling subgraphs  $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

We consider the following 6 cases.

**Case (1).** In  $DB(k)(-3k - 6 + 4i)$ ,  $1 \leq i \leq \frac{3k+3}{2}$ , we do the embeddings  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$  or  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 2. Hence, we obtain  $DB(k+2)(-3k - 8 + 4i)$ ,  $1 \leq i \leq \frac{3k+3}{2}$ , and in these labeling graphs, there exists the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Case (2).** In  $DB(k)(-3k - 2)$ , we do the embedding  $\begin{bmatrix} 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 6. We obtain  $DB(k+2)(-3k - 8)$  and in the labeling graph, there exist the labeling subgraph  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

**Case (3).** In  $DB(k)(-3k-6)$ , we do the embedding  $\begin{bmatrix} 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$  and  $\begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  to decrease  $e(1) - e(0)$  by 6. We obtain  $DB(k+2)(-3k-12)$ , and in the labeling graph, there exist the labeling subgraphs  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ .

**Case (4).** In  $DB(k)(3k)$ , we do the embedding  $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$  to increase  $e(1) - e(0)$  by 2. We obtain  $DB(k+2)(3k+2)$ , and in the labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ .

**Case (5).** In  $DB(k)(3k+4)$ , we do the embedding  $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$  to increase  $e(1) - e(0)$  by 2. We obtain  $DB(k+2)(3k+6)$ , and in the labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ .

**Case (6).** In  $DB(k)(3k+4)$ , we do the embedding  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  to increase  $e(1) - e(0)$  by 6. We obtain  $DB(k+2)(3k+10)$ , and in the labeling graph, there exists the labeling subgraph  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

By mathematical induction, we have  $FFI(DB(m)) = \{-3m - 6 + 4i : 0 \leq i \leq \frac{3m+5}{2}\}$ . The proof is complete.  $\square$

Hence, the cordiality and friendly index sets of  $DB(m)$  are determined as follows.

**Corollary 3.7.** For odd  $m \geq 1$ ,  $DB(m)$  is cordial. For even  $m \geq 0$ , we have

- (1)  $DB(m)$  is cordial if  $m \equiv 0 \pmod{4}$ , and
- (2)  $DB(m)$  is not cordial if  $m \equiv 2 \pmod{4}$ .

*Proof.* By Theorem 3.6, there exists  $|e(1) - e(0)| = 1$  for odd  $m \geq 1$ . So  $DB(m)$  is cordial. By Theorem 3.5, there exists  $|e(1) - e(0)| = 0$  for  $m \equiv 0 \pmod{4}$ , but for  $m \equiv 2 \pmod{4}$ , the minimum value of  $|e(1) - e(0)| = 2$ . Hence,  $DB(m)$  is cordial for  $m \equiv 0 \pmod{4}$ , but not for  $m \equiv 2 \pmod{4}$ .  $\square$

**Corollary 3.8.** For odd  $m$ ,  $FI(DB(m)) = \{2i + 1 : 0 \leq i \leq \frac{3m+5}{2}\}$ . For even  $m \geq 0$ , we have

- (1)  $FI(DB(m)) = \{4i : 0 \leq i \leq \frac{3m+8}{4}\}$  if  $m \equiv 0 \pmod{4}$ , and
- (2)  $FI(DB(m)) = \{4i + 2 : 0 \leq i \leq \frac{3m+6}{4}\}$  if  $m \equiv 2 \pmod{4}$ .

*Proof.* By Theorem 3.6, for odd  $m \geq 1$ ,  $FFI(DB(m)) = \{-3m - 6, -3m - 2, -3m + 2, \dots, 3m + 4\}$ . Hence,  $FI(DB(m)) = \{1, 3, \dots, 3m + 6\} = \{2i + 1 :$

$0 \leq i \leq (3m+5)/2\}$ . By Theorem 3.5,  $FFI(DB(0)) = \{-8, -4, 0, 4, 8\}$ , and for even  $m \geq 2$ ,  $FFI(DB(m)) = \{-3m-8, -3m-4, -3m, \dots, 3m+4\}$ . Hence, for  $m \equiv 0 \pmod{4}$ ,  $FI(DB(m)) = \{0, 4, 8, \dots, 3m+8\} = \{4i : 0 \leq i \leq (3m+8)/4\}$ ; and for  $m \equiv 2 \pmod{4}$ ,  $FI(DB(m)) = \{2, 6, 10, \dots, 3m+8\} = \{4i+2 : 0 \leq i \leq (3m+6)/4\}$ .  $\square$

Shiu and Wong [15] introduced the full product-cordial index set of  $G$ .

**Definition 3.9.** Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ , a vertex labeling  $f : V(G) \rightarrow \mathbb{Z}_2$  induces an edge labeling  $f^* : E(G) \rightarrow \mathbb{Z}_2$  defined by  $f^*(xy) = f(x)f(y)$ , for each edge  $xy \in E(G)$ . For each  $i \in \mathbb{Z}_2$ , let  $v_f(i) = |\{u \in V(G) : f(u) = i\}|$  and  $e_{f^*}(i) = |\{xy \in E(G) : f^*(xy) = i\}|$ . The full product-cordial index set of  $G$ , denoted  $FPCI(G)$ , is defined as  $\{e_{f^*}(1) - e_{f^*}(0) : \text{the vertex labeling } f \text{ is friendly}\}$ .

They obtained the following result.

**Lemma 3.10.** (Shiu and Wong[15]) Let  $f$  be a friendly labeling of  $G$ ,  $f^*$  be a product labeling of  $G$ . If  $G$  is an  $r$ -regular graph of even order, then  $e_{f^*}(1) - e_{f^*}(0) = -\frac{1}{2}(|E| + e_{f^+}(1) - e_{f^+}(0))$ .

Since  $DB(m)$  is 3-regular graph,  $|V| = 2(m+4)$ ,  $|E| = 3m+12$ . By Lemma 3.10 and Theorems 3.5, 3.6, we have

**Corollary 3.11.** For even  $m$ ,  $FPCI(DB(m)) =$

- (1)  $\{-2i : 1 \leq i \leq 5\}$  if  $m = 0$ ,
- (2)  $\{-2i : 1 \leq i \leq \frac{3m+8}{2}\}$  if  $m \geq 2$ .

**Corollary 3.12.** For odd  $m \geq 1$ ,  $FPCI(DB(m)) = \{-2i-1 : 1 \leq i \leq \frac{3m+7}{2}\}$ .

#### 4. CONCLUSION

In this paper, we determined the cordiality of a family of cubic graphs by the labeling subgraph embeddings method. The results in [3] show that this method may be used to determine the cordiality of all families of graphs that can be constructed by repeated subgraph embeddings and that the full friendly indices form an arithmetic sequence.

#### ACKNOWLEDGMENTS

The authors thank the referee for the valuable comments that improve the clarity of some proofs. The work is supported by NSFC Grant No.11371109. The authors wish to thank the referee for the valuable comments.

## REFERENCES

1. I. Cahit, Cordial graphs: a weaker version of graceful and harmonious graphs, *Ars Combin.*, **23**, (1987), 201-207.
2. J. A. Gallian, A dynamic survey of graph labeling, *Elect. J. Combin.*, (2015) #DS6.
3. Zh.-B. Gao, H.-N. Ren, S.-M. Lee, R.-Y. Han, G.-Ch. Lau, A new method in finding full friendly indices, *Bull. Mal. Math. Sci. Soc.*, (2016). doi:10.1007/s40840-016-0373-8.
4. Y. S. Ho, S.-M. Lee, H. K. Ng, On friendly index sets of root-unions of stars by cycles, *J. Combin. Math. Combin. Comput.*, **62**, (2007), 97-120.
5. M. Hovey, A-cordial graphs, *Discrete Math.* **93**, (1991), 183-194.
6. H. Kwong, S.-M. Lee, On friendly index sets of generalized books, *J. Combin. Math. Combin. Comput.*, **66**, (2008), 43-58.
7. H. Kwong, S.-M. Lee, H. K. Ng, On friendly index sets of 2-regular graphs, *Discrete Math.*, **308**, (2008), 5522-5532.
8. S.-Min Lee, H. K. Ng, On friendly index sets of cycles with parallel chords, *Ars Combin.*, **97A**, (2010), 253-267.
9. S.-M. Lee H.K. Ng, On friendly index sets of Prisms and Möbius ladder, *J. Combin. Math. Combin. Comput.*, **90**, (2014), 59-74.
10. W. C. Shiu, M. H. Ho, Full friendly index sets and full product-cordial index sets of some permutation Petersen graphs, *J. Comb. Number Theory*, **5(3)**, (2013), 227-244.
11. W. C. Shiu, M.H. Ho, Full friendly index sets of slender and flat cylinder graphs, *Trans. Combin.*, **2(4)**, (2013), 63-80.
12. W. C. Shiu, H. Kwong, Full friendly index sets of  $P_2 \times P_n$ , *Discrete Math.*, **308**, (2008), 3688-3693.
13. W. C. Shiu, S.-M. Lee, Full friendly index sets and full product-cordial index sets of twisted cylinders, *J. Comb. Number Theory*, **3(3)**, (2012), 209-216.
14. W. C. Shiu, M. H. Ling, Full friendly index sets of Cartesian products of two cycles, *Acta Math. Sin.*, **26**, (2010), 1233-1244.
15. W. C. Shiu, F. S. Wong, Full friendly index sets of cylinder graphs, *Australas J. Combin.*, **52**, (2012), 141-162.
16. D. Sinha, J. Kaur, Full friendly index set-I, *Discrete Appl. Math.*, **161**, (2013), 1262-1274.